

Statistica Sinica Preprint No: SS-2016-0285.R2

Title	Aggregated Expectile Regression by Exponential Weighting
Manuscript ID	SS-2016-0285.R2
URL	http://www.stat.sinica.edu.tw/statistica/
DOI	10.5705/ss.202016.0285
Complete List of Authors	Yuwen Gu and Hui Zou
Corresponding Author	Hui Zou
E-mail	zouxx019@umn.edu
Notice: Accepted version subject to English editing.	

AGGREGATED EXPECTILE REGRESSION BY EXPONENTIAL WEIGHTING

Yuwen Gu and Hui Zou

University of Connecticut and University of Minnesota

Abstract: Various estimators have been proposed to estimate conditional expectiles, including those from the multiple linear expectile regression, local polynomial expectile regression, boosted expectile regression, and so on. It is common practice that several plausible candidate estimators are fitted and a final estimator is selected from the candidate list. In this article, we advocate the use of an exponential weighting scheme to adaptively aggregate the candidate estimators into a final estimator. We show oracle inequalities for the aggregated estimator. Simulations and real data examples demonstrate that the aggregated estimator could have substantial gain in accuracy under both the squared and asymmetric squared errors.

Key words and phrases: Cross-validation, expectile regression, oracle inequality, model aggregation.

1. Introduction

Expectiles (Newey and Powell, 1987) are informative location measures of probability distributions. For each $\tau \in (0, 1)$, the τ th expectile of a

probability distribution F is defined as the quantity e_τ that satisfies

$$\int_{-\infty}^{e_\tau} |x - e_\tau| dF(x) = \tau \int_{-\infty}^{\infty} |x - e_\tau| dF(x). \quad (1.1)$$

Denote \mathcal{E}^τ the expectile operator at level τ such that $\mathcal{E}^\tau(F) = e_\tau$. For any random variable $Y \sim F$, we will also write $\mathcal{E}^\tau(Y) = \mathcal{E}^\tau(F)$. It can be shown that the 0.5th expectile coincides with the mean, $\mathcal{E}^{0.5}(F) = \int_{-\infty}^{\infty} x dF(x)$, and moreover, all the expectiles exist as long as the mean is finite. Expectiles have a clear financial meaning: $\mathcal{E}^\tau(F)$ is the amount of money that should be added to a position in order to have a pre-specified gain-loss ratio (Bernardo and Ledoit, 2000). Specifically, suppose $Y \sim F$ and let $x_+ = \max(x, 0)$ and $x_- = \max(-x, 0)$. From (1.1), one has

$$\frac{\mathbb{E}(Y - e_\tau)_+}{\mathbb{E}(Y - e_\tau)_-} = \frac{1 - \tau}{\tau}, \quad (1.2)$$

where $\mathbb{E}(Y - e_\tau)_+$ and $\mathbb{E}(Y - e_\tau)_-$ can be interpreted as the expected magnitudes of the gain and loss respectively and $(1 - \tau)/\tau$ the targeted gain-loss ratio. When $\int_{-\infty}^{\infty} x^2 dF(x) < \infty$, expectiles can be obtained via $\mathcal{E}^\tau(F) = \arg \min_{a \in \mathbb{R}} \int_{-\infty}^{\infty} \Psi_\tau(x - a) dF(x)$, where $\Psi_\tau(u) = |\tau - I(u < 0)|u^2$ is the asymmetric squared error loss and $I(\cdot)$ represents the indicator function. The special role of expectiles in risk management has been recognized by

many researchers recently (Kuan et al., 2009; Taylor, 2008; Bellini et al., 2014; Bellini and Di Bernardino, 2017). In risk management, Value at Risk (VaR) and expected shortfall (ES) are the two most popular risk measures in use. However, it has been well known that VaR lacks the desired property of coherence (Artzner et al., 1999). Specifically, VaR is not sub-additive, which contradicts the diversification principle that merging portfolios together should reduce the risk. ES is coherent (Acerbi and Tasche, 2002), but nevertheless fails to enjoy elicibility (Gneiting, 2011), another desired property of risk measures for which meaningful point forecasts and forecast performance comparisons are possible. Expectiles are the only risk measure that is both coherent and elicible (Ziegel, 2016; Bellini and Bigozzi, 2015).

Expectile regression estimates the conditional expectiles of a response variable given a set of covariates and is a useful extension to the mean regression. It has been widely applied to finance, demography, and education (see Taylor, 2008; Schnabel and Eilers, 2009a; Sobotka et al., 2013b). Since its advent (Aigner et al., 1976), a variety of expectile regression methods have been proposed. The multiple linear expectile regression was systematically studied in Newey and Powell (1987). Nonparametric and semi-parametric expectile estimation methods have also been considered in the literature to allow for more flexibility. Among others, Yao and Tong (1996) provided

kernel smoothing estimators of the conditional expectiles based on local polynomial regression. Their work was followed and extended by Guo and Härdle (2012), in which simultaneous confidence bands were established for the expectile functions. A nonparametric expectile estimation method based on spline smoothing was introduced in Schnabel and Eilers (2009b), and similar to that, Sobotka et al. (2013a) proposed a semi-parametric expectile estimation approach using splines. Yang and Zou (2015) proposed a nonparametric multiple expectile regression method using gradient boosting with regression tree base learners.

With the availability of various expectile regression methods, a practical problem is to choose the right method for the data at hand. The topic of model selection in the context of mean regression has been heavily studied in the literature. For example, lots of work has been devoted to the so-called model selection information criteria such as AIC (Akaike, 1974) and BIC (Schwarz, 1978). To our knowledge, there is no AIC- or BIC-like model selection criterion for expectile regression that has been justified theoretically. Moreover, these information criteria are often not applicable when comparing a parametric model with a nonparametric alternative. As such, cross-validation has been widely applied in practice and of course can be used in the context of expectile regression. The model selection process

by information criteria or cross-validation is always stochastic. Consequently, the uncertainty in model selection is inherently a part of the stochastic error in the final chosen model. Therefore, when the model selection uncertainty is large, the selected model tends to suffer.

When several plausible expectile regression estimators are available, instead of trying to select the best one, a good alternative approach is aggregation. In the literature, this idea is also known as model averaging or model combining. One may use these three names interchangeably wherever no confusion arises. There are multiple ways to do aggregation. One popular approach is the Bayesian model averaging. We refer the interested readers to a review article by Hoeting et al. (1999) on Bayesian model averaging. In this article, we take an exponential weighting scheme to combine different expectile regression estimators. Our estimator is a weighted average of these candidate estimators and the weight of each candidate estimator is inversely proportional to the exponential of its cumulative empirical prediction risk. Such an exponential weighting scheme has a solid information-theoretic justification in the context of conditional mean regression. See, for example, Yang (2001, 2004) and Catoni and Picard (2004). We prove an oracle inequality for the aggregated expectile regression estimator by exponential weighting in terms of both prediction risk and squared error risk. The theory

implies that the aggregated expectile regression estimator at least behaves like the best candidate expectile regression estimator. We further compare the aggregated estimator and the cross-validated estimator by extensive simulations. It is shown that the aggregated estimator significantly outperforms the cross-validated estimator when there is selection uncertainty.

The article is organized as follows. In the next section, we present the aggregated expectile regression estimator and study its theoretical properties. The applications of the aggregated expectile regression are introduced in Section 3 through several simulation examples. We apply the aggregated expectile regression to real personal computer data and S&P 500 Index data in Section 4. The technical proofs are given in the supplementary file.

2. Aggregated Expectile Regression by Exponential Weighting

2.1. Setup and notation

Consider the standard regression setting with i.i.d. observations (y_i, \mathbf{x}_i) , $i = 1, \dots, n$, where $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})^\top$ are p -dimensional covariate vectors and y_i are scalar responses. Assume these observations are realizations of the random pair (Y, \mathbf{X}) , where $\mathbf{X} = (X_1, \dots, X_p) \in \mathbb{R}^p$ and $Y \in \mathbb{R}$. Let $m(\mathbf{x}) = \mathbb{E}(Y|\mathbf{X} = \mathbf{x})$ and $\sigma^2(\mathbf{x}) = \text{var}(Y|\mathbf{X} = \mathbf{x})$ be respectively the conditional mean and variance functions. Assume both $m(\mathbf{X})$ and $\sigma(\mathbf{X})$ exist and $\sigma(\mathbf{X}) > 0$ almost surely. Define $\varepsilon = (Y - m(\mathbf{X}))/\sigma(\mathbf{X})$. It follows

immediately that $\mathbb{E}(\varepsilon|\mathbf{X}) = 0$ and $\text{var}(\varepsilon|\mathbf{X}) = 1$ almost surely. For ease of exposition, let us write Y in terms of \mathbf{X} and ε as

$$Y = m(\mathbf{X}) + \sigma(\mathbf{X})\varepsilon. \quad (2.1)$$

It should be noted that (2.1) by no means implies that we restrict ourselves to additive models only, although additive models are obviously included. As a matter of fact, a model with multiplicative error, for instance, $Y = f(\mathbf{X})\varepsilon$, can be easily cast into (2.1) as long as the conditional mean and variance functions of Y given \mathbf{X} exist. Denote the τ th conditional expectile by $e_\tau(\mathbf{x}) = \mathcal{E}^\tau(Y|\mathbf{X} = \mathbf{x})$, $0 < \tau < 1$. The goal of the expectile regression is to estimate $e_\tau(\mathbf{x})$. For an estimator $\hat{e}_\tau(\mathbf{x})$ of the expectile function $e_\tau(\mathbf{x})$, define the prediction risk and squared error risk of $\hat{e}_\tau(\mathbf{x})$ by $\mathbb{E}\Psi_\tau(Y - \hat{e}_\tau(\mathbf{X}))$ and $\mathbb{E}(\hat{e}_\tau(\mathbf{X}) - e_\tau(\mathbf{X}))^2$, respectively. If the τ th conditional expectile of ε given \mathbf{X} is $b_\tau(\mathbf{X})$, then $e_\tau(\mathbf{X}) = m(\mathbf{X}) + \sigma(\mathbf{X})b_\tau(\mathbf{X})$. When $e_\tau(\mathbf{x})$ is approximately linear in \mathbf{x} , the linear expectile regression is expected to perform quite well. However, when complicated nonlinear pattern exists in $e_\tau(\mathbf{x})$, the linear expectile regression can result in very large bias. As a remedy, nonparametric expectile regression methods can be used to accommodate the non-linearity. Of course, nonparametric expectile regression often has higher estimation

variance than the linear expectile regression.

2.2. The aggregation algorithm

Suppose we have a sequence of estimating procedures $\Delta = \{\delta_j, j \geq 1\}$, all of which can provide estimates of $e_\tau(\mathbf{x})$. Specifically, let us denote the estimate of $e_\tau(\mathbf{x})$ from procedure $\delta_j \in \Delta$ fitted on data with sample size n by $\hat{e}_{\tau,j,n}(\mathbf{x})$, $j \geq 1$. Note that we allow the number of procedures to be either finite or countably infinite. Following Yang (2001), we impose no special assumptions on the procedures and they can be either model-based or non-model-based. The goal is to construct an estimating procedure δ_a by adaptively aggregating this sequence of candidate estimating procedures in the hope of achieving a small estimation risk. The algorithm for this aggregation is displayed in Algorithm 1.

In Algorithm 1, n_0 is often chosen such that both n_0 and $n - n_0$ are of the same order as n . See more discussion in the next subsection. The tuning parameter λ is a properly chosen constant which controls the effect of the performance of the candidate estimators on the weights. On one hand, when λ is very small, Algorithm 1 will assign almost equal weights to the candidate estimators. In the extreme case $\lambda = 0$, Algorithm 1 is merely a simple average of the candidate estimators. On the other hand, when λ is large enough, Algorithm 1 will put almost all the weights on the procedure

Algorithm 1: The aggregated expectile regression by exponential weighting (AEREW) – Single split.

1. Randomly split the data into two parts. Without loss of generality, denote the two parts by $D^{(0)} = (y_i, \mathbf{x}_i)_{i=1}^{n_0}$ and $D^{(1)} = (y_i, \mathbf{x}_i)_{i=n_0+1}^n$ respectively, where $D^{(0)}$ is used for training and $D^{(1)}$ is used for evaluation.
2. For each procedure δ_j , obtain the estimate $\hat{e}_{\tau,j,n_0}(\mathbf{x}_i)$ of $e_{\tau}(\mathbf{x}_i)$ for every $\mathbf{x}_i \in D^{(1)}$ based on the training data $D^{(0)}$, $n_0 + 1 \leq i \leq n$, $j \geq 1$.
3. Set $W_{j,n_0+1} = \pi_j$ such that $\pi_j \geq 0$, $j \geq 1$, and $\sum_{j=1}^{\infty} \pi_j = 1$. For $j \geq 1$ and $n_0 + 2 \leq i \leq n$, calculate the weights

$$W_{j,i} = \frac{\pi_j \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j,n_0}(\mathbf{x}_k)) \right\}}{\sum_{j'=1}^{\infty} \pi_{j'} \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j',n_0}(\mathbf{x}_k)) \right\}}.$$

Obtain the aggregating procedure δ_a which estimates $e_{\tau}(\mathbf{x})$ by

$$\hat{e}_{\tau,\cdot,n}(\mathbf{x}) = \sum_{j=1}^{\infty} \left(\sum_{i=n_0+1}^n \frac{W_{j,i}}{n - n_0} \right) \hat{e}_{\tau,j,n_0}(\mathbf{x}).$$

with best performance upon evaluation on $D^{(1)}$. We comment on the choice of λ in Remark 3 in the next subsection.

Note that in Algorithm 1, the weights $W_{j,i}$, $n_0 + 1 \leq i \leq n$, $j \geq 1$, depend on the order of the observations from the random partition. Multiple splits can be carried out as shown in Algorithm 2 to avoid large variance in the weights. From the computational point of view, these multiple splits can be carried out in parallel to accelerate the computation. We recommend Algorithm 2 for practical use, and according to our empirical studies, the algorithm often works quite well when the number of splits B is taken to be several hundred.

Algorithm 2: The aggregated expectile regression by exponential weighting (AEREW) – Multiple splits.

1. Randomly split the data into two parts. Without loss of generality, denote the two parts by $D^{(0)} = (y_i, \mathbf{x}_i)_{i=1}^{n_0}$ and $D^{(1)} = (y_i, \mathbf{x}_i)_{i=n_0+1}^n$ respectively, where $D^{(0)}$ is used for training and $D^{(1)}$ is used for evaluation.
2. For each procedure δ_j , obtain the estimate $\hat{e}_{\tau,j,n_0}(\mathbf{x}_i)$ of $e_{\tau}(\mathbf{x}_i)$ for every $\mathbf{x}_i \in D^{(1)}$ using the training data $D^{(0)}$ for fitting, $n_0 + 1 \leq i \leq n$, $j \geq 1$.
3. Set $W_{j,n_0+1} = \pi_j$ such that $\pi_j \geq 0$, $j \geq 1$, and $\sum_{j=1}^{\infty} \pi_j = 1$. For $j \geq 1$ and $n_0 + 2 \leq i \leq n$, calculate the weights

$$W_{j,i} = \frac{\pi_j \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j,n_0}(\mathbf{x}_k)) \right\}}{\sum_{j'=1}^{\infty} \pi_{j'} \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j',n_0}(\mathbf{x}_k)) \right\}}.$$

4. Repeat the above three steps $(B - 1)$ more times. Denote the estimates and weights from the k th random split by $\hat{e}_{\tau,j,n_0}^{(k)}(\mathbf{x})$ and $W_{j,i}^{(k)}$, $n_0 + 1 \leq i \leq n$, $j \geq 1$, $1 \leq k \leq B$, respectively. Obtain the aggregating procedure δ_a^B which estimates $e_{\tau}(\mathbf{x})$ by

$$\hat{e}_{\tau, \cdot, n}^B(\mathbf{x}) = \sum_{j=1}^{\infty} \sum_{i=n_0+1}^n \sum_{k=1}^B \frac{W_{j,i}^{(k)}}{B(n - n_0)} \hat{e}_{\tau,j,n_0}^{(k)}(\mathbf{x}).$$

2.3. Oracle inequalities for AEREW

We provide the oracle inequalities for AEREW in terms of statistical risk bounds under both squared and asymmetric squared error losses in the following theorem. To facilitate the discussion, let us introduce some notation. We denote $\underline{c} = \min(\tau, 1 - \tau)$ and $\bar{c} = \max(\tau, 1 - \tau)$. For a random variable Z , let us define the sub-exponential norm of Z by $\|Z\|_{\text{SEXP}} \equiv \sup_{k \geq 1} k^{-1} (\mathbb{E}|Z|^k)^{1/k}$. If $\|Z\|_{\text{SEXP}}$ is finite, we call Z a sub-exponential random variable (see, e.g., Vershynin, 2010).

Theorem 1. *Under the general model (2.1), assume that the candidate estimators satisfy the following conditions:*

(C1) *With probability one, $\sup_{i,j} |\hat{e}_{\tau,j,i}(\mathbf{X}) - e_{\tau}(\mathbf{X})| \leq A_{\tau}$ and $|e_{\tau}(\mathbf{X})| \leq B_{\tau}$,*

where $A_{\tau}, B_{\tau} \in (0, \infty)$ are positive constants depending on τ .

(C2) *With probability one, $|\sigma(\mathbf{X})| \leq C_0$, where $C_0 \in (0, \infty)$ is also a positive constant.*

(C3) *With probability one, the sub-exponential norm of ε given \mathbf{X} is bounded by a positive constant $K \in (0, \infty)$.*

Let $K_{\tau} = 2\bar{c}(K + B_{\tau})$ and $D_{\tau} = 4eK_{\tau}$, where $e = \exp(1)$. Define the two functions $\mathcal{M}_0(t) = 2 \exp(2e^2 K_{\tau}^2 t^2)$ and $\mathcal{M}_2(t) = 16\sqrt{2} \exp(4e^2 K_{\tau}^2 t^2)$. When

the tuning parameter λ is chosen such that

$$\lambda \leq \min \left\{ \frac{1}{2C_0 A_\tau D_\tau}, \frac{\underline{c} \exp(-\bar{c} A_\tau (C_0 D_\tau)^{-1})}{C_0^2 \mathcal{M}_2(D_\tau^{-1}) + 16\bar{c}^2 A_\tau \mathcal{M}_0(D_\tau^{-1})} \right\}, \quad (2.2)$$

the risk of the aggregated estimator by AEREW (Algorithm 1 and Algorithm 2) under loss Ψ_τ has the following upper bound

$$\mathbb{E} \Psi_\tau(Y - \hat{e}_{\tau, \cdot, n}(\mathbf{X})) \leq \inf_{j \geq 1} \left\{ \frac{\log(1/\pi_j)}{\lambda(n - n_0)} + \mathbb{E} \Psi_\tau(Y - \hat{e}_{\tau, j, n_0}(\mathbf{X})) \right\} \quad (2.3)$$

and the risk of the aggregated estimator under the squared error loss satisfies

$$\mathbb{E}(\hat{e}_{\tau, \cdot, n}(\mathbf{X}) - e_\tau(\mathbf{X}))^2 \leq \inf_{j \geq 1} \left\{ \frac{\log(1/\pi_j)}{\lambda \underline{c}(n - n_0)} + \frac{\bar{c}}{\underline{c}} \mathbb{E}(\hat{e}_{\tau, j, n_0}(\mathbf{X}) - e_\tau(\mathbf{X}))^2 \right\}, \quad (2.4)$$

where (\mathbf{X}, Y) is taken to be a random observation from (2.1) that is independent of the observations $(\mathbf{X}_i, Y_i)_{i=1}^n$.

Remark 1. The assumption of conditions (C1) – (C2) is mild and can be easily satisfied with high probability if the mean function $m(\mathbf{X})$ as well as the variance function $\sigma^2(\mathbf{X})$ are bounded almost surely. This assumption is fairly common in related work for aggregation.

Remark 2. The class of sub-exponential random variables covers all random variables for which the moment generating functions exist in a neighborhood of zero and hence is quite large to encompass commonly used error distributions. As a consequence, condition (C3) does not restrict the response Y to be bounded, which is however a condition often assumed in the machine learning literature for simplicity.

Remark 3. The oracle inequality (2.3) tells us that the aggregated estimator achieves a prediction risk that is smaller than the smallest prediction risk offered by the candidate estimators plus an additional risk term. Assume there are M candidate estimators all of which are assigned equal prior weights, the extra risk term in the oracle inequality (2.3) becomes $\log(M)/\{\lambda(n-n_0)\}$. Although λ has an upper bound by (2.2) in order for us to prove the oracle inequalities, there should also be a lower bound for λ in order to make the extra term $\frac{\log(M)}{\lambda(n-n_0)}$ much smaller than $\mathbb{E}\Psi_\tau(Y - e_\tau(\mathbf{X}))$. Otherwise the oracle inequality offers no meaningful conclusions. Typically, $\mathbb{E}\Psi_\tau(Y - \hat{e}_{\tau,j,n_0}(\mathbf{X}))$ converges to $\mathbb{E}\Psi_\tau(Y - e_\tau(\mathbf{X}))$ at rate n_0^{-1} for parametric estimators and at a slower rate than n_0^{-1} for nonparametric estimators. So if one only cares about the absolute prediction risk, then we only need to require $\frac{\log(M)}{\lambda(n-n_0)} \ll 1$. Of course, we often also care about the rate of convergence. If n_0 is chosen such that both n_0 and $n - n_0$ are of order n , then as long as $\lambda \geq \mathcal{O}(\log(M))$

the extra risk term $\frac{\log(M)}{\lambda(n-n_0)}$ does not affect the rate of convergence, i.e., the aggregated estimator by AEREW will achieve the same rate of convergence as the best candidate estimator. We recommend using $\max(1, \lfloor \log(M) \rfloor)$ as the default value for λ . In all of our numerical experiments this default choice works very well and we have also found that the performance of AEREW is insensitive to the choice of λ in a fairly wide range.

2.4. Extension to time series data

Previous discussion mainly explored aggregation of expectile estimators from i.i.d. data. In practice, time series data are also frequently collected. We note that the aggregation algorithm (Algorithm 1) can be readily modified to combine expectile estimators from time series. Specifically, let us consider a time series Y_1, Y_2, \dots and let \mathbf{X}_i be the vector of explanatory variables related to Y_i at time $i \geq 1$. Our goal is to estimate the conditional expectiles of Y_i given \mathbf{X}_i and earlier data $Z^{i-1} = (Y_k, \mathbf{X}_k)_{k=1}^{i-1}$. Assume $Y_i = m_i + \sigma_i \varepsilon_i$, where m_i and σ_i^2 are respectively the conditional mean and variance functions of Y_i given $\mathbf{X}_i = \mathbf{x}_i$ and $Z^{i-1} = z^{i-1}$, and ε_i satisfies $\mathbb{E}(\varepsilon_i | \mathbf{X}_i, Z^{i-1}) = 0$ and $\text{var}(\varepsilon_i | \mathbf{X}_i, Z^{i-1}) = 1$ almost surely. Denote the τ th conditional expectile function of Y_i given $\mathbf{X}_i = \mathbf{x}_i$ and $Z^{i-1} = z^{i-1}$ by $e_{\tau,i}$, $i \geq 1$. Let $\hat{e}_{\tau,j,i}$ be the estimator of $e_{\tau,i}$ from procedure $\delta_j \in \Delta$ using data \mathbf{x}_i and z^{i-1} , $j \geq 1$, $i \geq 1$. We present in Algorithm 3 the aggregation algorithm for combining

expectile estimators for the time series.

Algorithm 3: The aggregated expectile regression by exponential weighting – Time series (AEREW-ts).

1. Spare the first n_0 observations $z^{n_0} = (y_i, \mathbf{x}_i)_{i=1}^{n_0}$ for initial estimation.
2. For $i = n_0 + 1, \dots, n$ and procedure δ_j , $j \geq 1$, obtain the estimate $\hat{e}_{\tau,j,i}$ of $e_{\tau,i}$ using data \mathbf{x}_i and z^{i-1} .
3. Set $\Lambda_{j,n_0+1} = \pi_j$ such that $\pi_j \geq 0$, $j \geq 1$, and $\sum_{j=1}^{\infty} \pi_j = 1$, For $j \geq 1$ and $n_0 + 2 \leq i \leq n$, calculate the weights

$$\Lambda_{j,i} = \frac{\pi_j \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j,k}) \right\}}{\sum_{j'=1}^{\infty} \pi_{j'} \exp \left\{ -\lambda \sum_{k=n_0+1}^{i-1} \Psi_{\tau}(y_k - \hat{e}_{\tau,j',k}) \right\}}.$$

Obtain the aggregating procedure δ_a which estimates $e_{\tau,i}$ by

$$\hat{e}_{\tau,.,i} = \sum_{j=1}^{\infty} \Lambda_{j,i} \hat{e}_{\tau,j,i}, \quad i \geq n_0 + 1.$$

Under conditions modified accordingly to the time series, we can establish the oracle inequalities for the aggregated expectile estimator.

Theorem 2. *Assume that the candidate estimators satisfy the following conditions:*

(C1') *With probability one, $\sup_{i,j} |\hat{e}_{\tau,j,i} - e_{\tau,i}| \leq A_{\tau}$ and $\sup_i |e_{\tau,i}| \leq B_{\tau}$,*

where $A_{\tau}, B_{\tau} \in (0, \infty)$ are positive constants depending on τ .

(C2') *With probability one, $|\sigma_i| \leq C_0$ for all $i \geq 1$, where $C_0 \in (0, \infty)$ is also a positive constant.*

(C3') With probability one, the sub-exponential norm of ε_i given \mathbf{X}_i and Z^{i-1} is bounded by a positive constant $K \in (0, \infty)$ for all $i \geq 1$.

When the tuning parameter λ is chosen to satisfy (2.2), the mean average risk of the aggregated estimator by AEREW-ts (Algorithm 3) under loss Ψ_τ has the following upper bound

$$\begin{aligned} & \frac{1}{n - n_0} \sum_{i=n_0+1}^n \mathbb{E} \Psi_\tau(y_i - \hat{e}_{\tau, \cdot, i}) \\ & \leq \inf_{j \geq 1} \left\{ \frac{\log(1/\pi_j)}{\lambda(n - n_0)} + \frac{1}{n - n_0} \sum_{i=n_0+1}^n \mathbb{E} \Psi_\tau(y_i - \hat{e}_{\tau, j, i}) \right\} \end{aligned} \quad (2.5)$$

and the mean average risk of the aggregated estimator under the squared error loss satisfies

$$\begin{aligned} & \frac{1}{n - n_0} \sum_{i=n_0+1}^n \mathbb{E}(\hat{e}_{\tau, \cdot, i} - e_{\tau, i})^2 \\ & \leq \inf_{j \geq 1} \left\{ \frac{\log(1/\pi_j)}{\lambda \underline{c}(n - n_0)} + \frac{\bar{c}/\underline{c}}{n - n_0} \mathbb{E}(\hat{e}_{\tau, j, i} - e_{\tau, i})^2 \right\}. \end{aligned} \quad (2.6)$$

3. Applications and Simulation Examples

In this section, we demonstrate several useful applications of aggregation in expectile regression. These applications are illustrated through two simulation examples.

3.1. Local expectile regression: bandwidth selection or aggrega-

tion?

When there is a single covariate, nonparametric expectile regression can be done via the local fitting scheme as shown in Yao and Tong (1996). It was argued by Yao and Tong (1996) that the local linear fit automatically corrects the boundary effects inherited from the local constant fit (see also Fan, 1992) and the estimator of the derivative plays an important role in monitoring the reliability of non-linear prediction and in detecting chaos. To be specific, given a random sample $(Y_i, X_i)_{i=1}^n$, the local linear estimators of $e_\tau(x) = \mathcal{E}^\tau(Y|X = x)$ and $e'_\tau(x) = de_\tau(x)/dx$ are defined as

$$\begin{aligned} & (\hat{e}_\tau(x; h), \hat{e}'_\tau(x; h)) \\ &= \arg \min_{a, b \in \mathbb{R}} \sum_{i=1}^n \Psi_\tau(Y_i - a - b(X_i - x)) \frac{1}{h} K\left(\frac{X_i - x}{h}\right), \end{aligned} \quad (3.1)$$

where $K(\cdot)$ is the kernel density and $h > 0$ is the bandwidth. Though many kernel densities are available for the local linear regressions, we choose the Gaussian kernel $K(u) = (2\pi)^{-1/2} \exp(-u^2/2)$ for illustrative purposes.

The choice of h reflects the trade-off between the bias and variance of the estimators and has a high impact on the performance of the prediction. The theoretically optimal bandwidth is $h = C_\tau n^{-1/5}$ where C_τ depends on unknown quantities (Yao and Tong, 1996). In practice, we can select the

bandwidth through cross-validation. See Heidenreich et al. (2013) for a recent review of bandwidth selection methods.

Alternatively, we can combine the kernel estimators at different bandwidths. Specifically, for a sequence of candidate bandwidths h_1, \dots, h_M , we obtain the local linear fit $\hat{e}_\tau(x; h_j)$ for each bandwidth h_j , $1 \leq j \leq M$ and combine these estimators using AEREW.

We illustrate this application through a simulation study. Consider the following heteroscedastic model

$$Y = 0.5\{X + 2 \exp(-16X^2)\} + \{0.4 \exp(-2X^2) + 0.2\}\varepsilon, \quad (3.2)$$

where the scalar covariate X is independent of the random error ε . Moreover, suppose $X \sim \text{Uniform}(-2, 2)$ and $\varepsilon \sim \text{Laplace}(0, 1/\sqrt{2})$. The density of ε is $f_\varepsilon(u) = \exp(-\sqrt{2}|u|)/\sqrt{2}$. Note that ε is a sub-exponential random variable satisfying $\mathbb{E}(\varepsilon) = 0$ and $\text{var}(\varepsilon) = 1$. A similar model to (3.2) was considered in Fan and Yao (1998) under a different error distribution. For the simulation study, a training set of $n = 200$ observations was randomly generated from model (3.2) and local linear regressions (3.1) were fitted to the training data with five candidate bandwidths $\mathbf{h} = (0.1, 0.3, 0.5, 0.7, 0.9) \times (200)^{-1/5}$.

To demonstrate the benefit of aggregation and compare it with cross-

validation in bandwidth selection, we applied a five-fold cross-validation to select the best bandwidth and also combined the five local linear expectile regression estimators using AEREW (Algorithm 2), for which $B = 200$ splits were conducted and the splitting size $n_0 = 160$ and prior weights $\pi_j = 1/5$, $1 \leq j \leq 5$ were chosen. We also set $\lambda = 1$. To compare the estimation performance of different procedures, we independently simulated a test set of $n_1 = 10000$ observations from model (3.2) and calculated the following two performance measures based on the test data. Assume the true expectile function is $e_\tau(\cdot)$ and its estimate from a specific procedure is $\hat{e}_\tau(\cdot)$. The two measures: (i) the estimated prediction risk, and (ii) estimated squared deviation (MSD) for $\hat{e}_\tau(\cdot)$ are respectively defined as

$$\begin{aligned} \text{risk}(\tau) &= \frac{1}{n_1} \sum_{i=1}^{n_1} \Psi_\tau(Y_i - \hat{e}_\tau(X_i)), \quad \text{and} \\ \text{MSD}(\tau) &= \sqrt{\frac{1}{n_1} \sum_{i=1}^{n_1} (\hat{e}_\tau(X_i) - e_\tau(X_i))^2}. \end{aligned} \tag{3.3}$$

For model (3.2), the true expectile function is $e_\tau(x) = 0.5\{x + 2 \exp(-16x^2)\} + \{0.4 \exp(-2x^2) + 0.2\}b_\tau$, where $b_\tau = \mathcal{E}^\tau(\varepsilon)$ is the τ th expectile of the Laplace random error. The simulations were repeated $M = 100$ times under the above setting. For illustrative purposes, we also present the proportion p_{CV} of each candidate estimator being selected by the five-fold cross-validation

among these 100 runs. The results are summarized in Table 1.

Table 1: Estimated prediction risks and MSDs of local linear regressions with five candidate bandwidths, the five-fold cross-validated kernel estimator, and AEREW ($\lambda = 1$) for the heteroscedastic model (3.2). The numbers listed are averages over 100 independent runs with their respective standard errors reported in the parentheses. The proportion of each candidate estimator being selected by the five-fold cross-validation among these 100 runs is reported by p_{CV} . All numbers are of order 10^{-2} except those corresponding to p_{CV} .

τ	Measures	Bandwidth (h)					Cross-validation	Aggregation
		0.0347	0.104	0.173	0.243	0.312		
0.05	risk	3.71 (0.03)	2.98 (0.02)	2.90 (0.02)	2.90 (0.02)	2.92 (0.02)	2.93 (0.02)	2.89 (0.02)
	MSD	25.16 (0.31)	16.56 (0.39)	15.11 (0.36)	15.72 (0.30)	16.48 (0.25)	16.03 (0.33)	14.67 (0.32)
	p_{CV}	0.00	0.23	0.23	0.12	0.42	–	–
0.10	risk	4.98 (0.06)	4.13 (0.03)	4.06 (0.02)	4.12 (0.02)	4.20 (0.02)	4.17 (0.03)	4.07 (0.02)
	MSD	22.30 (0.39)	13.93 (0.37)	13.05 (0.31)	14.42 (0.24)	15.83 (0.20)	14.49 (0.40)	13.04 (0.29)
	p_{CV}	0.03	0.34	0.28	0.14	0.21	–	–
0.25	risk	6.69 (0.06)	5.82 (0.03)	5.85 (0.03)	6.05 (0.03)	6.26 (0.03)	5.88 (0.03)	5.86 (0.03)
	MSD	17.60 (0.39)	10.22 (0.22)	10.71 (0.20)	13.10 (0.18)	15.25 (0.16)	10.99 (0.29)	10.84 (0.21)
	p_{CV}	0.00	0.41	0.46	0.03	0.10	–	–
0.50	risk	7.63 (0.06)	6.72 (0.03)	6.78 (0.03)	7.04 (0.03)	7.37 (0.03)	6.78 (0.04)	6.81 (0.03)
	MSD	16.05 (0.32)	8.96 (0.23)	9.58 (0.21)	12.10 (0.17)	14.55 (0.15)	9.54 (0.25)	9.92 (0.22)
	p_{CV}	0.00	0.54	0.42	0.03	0.01	–	–
0.75	risk	6.76 (0.10)	5.88 (0.03)	5.92 (0.03)	6.13 (0.03)	6.41 (0.04)	5.96 (0.04)	5.94 (0.03)
	MSD	17.62 (0.51)	10.23 (0.25)	10.61 (0.23)	12.77 (0.19)	15.04 (0.16)	10.90 (0.29)	10.81 (0.23)
	p_{CV}	0.01	0.43	0.42	0.12	0.02	–	–
0.90	risk	4.98 (0.06)	4.17 (0.03)	4.09 (0.03)	4.16 (0.03)	4.31 (0.03)	4.15 (0.03)	4.10 (0.03)
	MSD	22.51 (0.35)	14.47 (0.36)	13.08 (0.32)	14.26 (0.26)	16.18 (0.20)	14.02 (0.32)	13.26 (0.31)
	p_{CV}	0.02	0.26	0.42	0.20	0.10	–	–
0.95	risk	3.87 (0.04)	3.08 (0.03)	2.97 (0.03)	3.00 (0.03)	3.08 (0.03)	3.04 (0.03)	2.99 (0.03)
	MSD	26.78 (0.34)	18.41 (0.45)	16.18 (0.46)	16.62 (0.43)	18.20 (0.38)	17.37 (0.45)	16.16 (0.42)
	p_{CV}	0.00	0.29	0.30	0.25	0.16	–	–

In Table 1, the performance measures are calculated by averaging over the 100 replicates and their respective standard errors are reported in the parentheses. It is clear from Table 1 that the optimal bandwidths are different for different expectile levels. Smaller bandwidths are preferred for expectile levels around 0.5 while at extreme expectile levels (τ close to 0 and 1), slightly larger bandwidths are favored. Also it is evident from Table 1 that AEREW compares quite favorably with the five-fold cross-validation. Indeed, AEREW outperforms the cross-validation for all expectile levels other than 0.5 and its performance there is very close to or even better than that of the best candidate estimator. On the other hand, the cross-validation gives slightly better estimation for the mean function ($\tau = 0.5$) than AEREW, but still AEREW performs quite well in this case. From the simulation result, it can be seen that when cross-validation is uncertain about the best estimator (several estimators have reasonably large p_{CV} values), AEREW can outperform the cross-validated estimator.

3.2. Multiple expectile regression: parametric or nonparametric?

We now demonstrate the application of AEREW in multiple expectile regression where more than one covariate is available. For such models, the local linear estimator, such as (3.1), is not very useful in practice when there are more than five covariates. In the current toolbox for multidimensional

expectile regression, we have the multiple linear expectile regression (Newey and Powell, 1987; Efron, 1991) as well as the regression tree based nonparametric gradient boosting (Yang and Zou, 2015). The question one often encounters in practice is which method to use. In fact, it is well known that the nonparametric regression is quite flexible to accommodate non-linearity but loses efficiency when the linear parametric model is correctly specified. In the case of expectile regression, when linear and nonlinear effects coexist in an underlying model, it is also possible that the expectile function is nearly linear at certain expectile levels and becomes highly nonlinear at other levels. Therefore, when multiple expectile levels need to be inspected together, it is beneficial to consider adaptively aggregating both parametric and nonparametric methods for better estimation.

As an illustration, let us consider the following heteroscedastic model with multiple covariates

$$Y = \mathbf{X}^T \boldsymbol{\beta} + 2\varepsilon \exp(-0.35X_2 - 1.1X_4), \quad (3.4)$$

where $\boldsymbol{\beta} = (1.5, 2.5, 1.0, 0.5, 2.0, 1.5)^T$, $\mathbf{X} = (X_1, \dots, X_6)^T \sim N(\mathbf{0}, \mathbf{I}_6)$, $\varepsilon \sim N(0, 1)$ and \mathbf{X} is independent of ε . The true expectile function in this model is $\mathbf{X}^T \boldsymbol{\beta} + 2b_\tau \exp(-0.35X_2 - 1.1X_4)$, where b_τ is the τ th expectile of $N(0, 1)$.

Intuitively, for some expectile levels (such as those near 0.5), the performance of parametric estimators may dominate that of nonparametric estimators, while the opposite is true at the other expectile levels (such as those with large or small τ values).

For the simulation study, we considered three candidate estimators. The multiple linear expectile regression and nonparametric multiple expectile regression via gradient boosting arise as the two natural candidates. For illustrative purposes, we also considered a slightly more complicated version of the linear expectile regression by including an additional interaction term between X_2 and X_4 besides all the main effects X_1, \dots, X_6 . We included such a linear interaction model because we expect it to improve the estimation accuracy of the conditional expectiles at some levels. The simulations were repeated $M = 100$ times. For each simulation, we first generated a training set of $n = 500$ observations and applied the three aforementioned candidate estimators, plus a five-fold cross-validation to select the best procedure, as well as an aggregation of the candidate estimators using AEREW (Algorithm 2) to obtain the estimates. In AEREW, the weights were averaged over $B = 200$ splits with split size $n_0 = 400$. Equal prior weights were assigned for the three candidate estimators for each split. We set $\lambda = 1$. The estimated prediction risks and MSDs, defined in (3.3), are

reported in Table 2 for independent test sets of $n_1 = 10000$ observations from model (3.4). In addition, we also include the proportion p_{CV} of each candidate estimator being selected by the cross-validation among the 100 independent simulations.

It can be seen from Table 2 that none of the three candidate estimator is universally better than the others for all expectile levels. Indeed, when τ is in the middle of its range (around 0.5), the linear expectile regression with only main effects compares favorably with the other two procedures. This is expected since the true expectile function is mainly linear. For extreme expectile levels (τ away from 0.5), the linear expectile regression with interaction and the gradient boosting outperform the linear expectile regression with main effects only. From the patterns of p_{CV} , it can be seen that for expectile levels that are close to 0, 1, or 0.5, there is a clear dominating candidate estimator which is selected by cross-validation with high probability. While for moderate expectile levels, there are usually two competing candidate estimators (see, e.g., $\tau = 0.25$ and 0.75). AEREW outperforms the three candidate estimators as well as the cross-validated estimator at all expectile levels. Moreover, we observe a greater gain by using AEREW for moderate expectile levels when cross-validation experiences difficulty in selecting a clear winner.

Table 2: Estimated prediction risks, MSDs and their respective standard errors (in parentheses) of the linear expectile regression with only main effects, linear expectile regression with interaction, nonparametric expectile regression via boosting, the five-fold cross-validation and AEREW ($\lambda = 1$) for the heteroscedastic model (3.4) over 100 independent runs. The proportion of each candidate estimator being selected by the cross-validation is summarized by p_{CV} .

τ	Measures	Individual			Cross-validation	Aggregation
		Linear	Interaction	Boosting		
0.05	Risk	53.58 (4.20)	49.07 (4.03)	53.39 (4.03)	49.56 (4.06)	47.71 (3.97)
	MSD	12.08 (0.39)	10.88 (0.39)	11.74 (0.30)	10.95 (0.39)	10.39 (0.30)
	p_{CV}	0.06	0.87	0.07	–	–
0.10	Risk	57.71 (2.86)	55.33 (2.84)	60.13 (2.82)	54.91 (2.76)	53.92 (2.74)
	MSD	8.41 (0.21)	7.74 (0.25)	9.13 (0.20)	7.78 (0.21)	7.59 (0.18)
	p_{CV}	0.17	0.71	0.12	–	–
0.25	Risk	64.87 (2.40)	66.14 (3.21)	69.06 (2.18)	65.86 (3.20)	62.99 (2.14)
	MSD	4.49 (0.16)	4.44 (0.24)	5.77 (0.15)	4.45 (0.24)	4.14 (0.13)
	p_{CV}	0.48	0.47	0.05	–	–
0.50	Risk	71.20 (2.41)	72.37 (2.45)	76.07 (2.45)	71.84 (2.44)	71.52 (2.41)
	MSD	1.51 (0.07)	2.01 (0.11)	3.39 (0.10)	1.75 (0.10)	1.67 (0.08)
	p_{CV}	0.71	0.21	0.08	–	–
0.75	Risk	67.78 (3.23)	67.34 (3.18)	70.07 (3.23)	67.62 (3.19)	66.62 (3.20)
	MSD	4.34 (0.11)	4.18 (0.13)	5.06 (0.11)	4.29 (0.12)	4.07 (0.11)
	p_{CV}	0.47	0.43	0.10	–	–
0.90	Risk	54.98 (3.26)	53.48 (3.50)	53.86 (3.09)	51.44 (3.03)	50.56 (3.05)
	MSD	8.40 (0.26)	7.78 (0.31)	8.41 (0.17)	7.64 (0.17)	7.46 (0.17)
	p_{CV}	0.20	0.66	0.14	–	–
0.95	Risk	46.76 (2.93)	43.32 (2.84)	45.91 (2.97)	43.83 (2.95)	42.22 (2.92)
	MSD	11.50 (0.25)	10.40 (0.24)	10.88 (0.22)	10.46 (0.23)	10.00 (0.20)
	p_{CV}	0.12	0.70	0.18	–	–

4. Real Data Examples

4.1. Personal computer data

We apply AEREW to a data set described in Stengos and Zacharias (2006). The data set contains monthly price information of personal computers from January 1993 to November 1995 and was analyzed using a hedonic analysis. There are $N = 6259$ observations with 10 variables. The response variable is *Price*, and the hedonic variables *Speed*, *HD*, *RAM*, *Screen*, *CD*, *Multi*, and *Premium* directly describe the major hedonic characteristics that make up a computer. The other two explanatory variables *ADs* and *Trend* are not directly related to the personal computer characteristics, but are believed to be associated with the price (Stengos and Zacharias, 2006). Therefore, we also include these two variables in our analysis. After inspecting the data, we decided to take the logarithmic transformation on all continuous variables except *Trend*. We considered a hedonic analysis at different price levels using expectile regression. Three candidate models were considered: the multiple linear expectile regression with main effects only, the linear expectile regression with main effects and two-way interactions, and the nonparametric approach via boosting. We applied a five-fold cross-validation to select the best procedure among the three candidates. Finally, we used AEREW to aggregate the three candidate models.

For the analysis, we randomly sampled $n = 3129$ observations from the data to form a training set, on which the three aforementioned candidate estimators were fitted and a five-fold cross-validation was applied to select the best procedure. To aggregate the candidate estimators through AEREW (Algorithm 2), we averaged the weights over $B = 200$ splits with split size $n_0 = 2503$. Equal prior weights were assigned and $\lambda = 1$ was set. The prediction risks of the procedures, as defined in (3.3), were evaluated on the remaining $n_1 = 3130$ observations. This procedure was repeated $M = 100$ times and the results are summarized in Table 3 and Figure 1. The proportions of the candidate estimators being selected by the five-fold cross-validation are reported by p_{CV} .

It can be seen from both Table 3 and Figure 1 that at all expectile levels, the linear expectile regression with interactions outperforms the linear expectile regression with main effects only. The boosting estimator performs better for expectile levels in the middle range, while the linear expectile regression with interactions is better for extreme expectile levels. At 0.10 level, boosting and the linear model with interactions are very comparable to each other. This can be seen from the values of p_{CV} . However, it is also very clear that if we consider multiple expectile levels, there is no clear winner among the candidate estimators. It is worth noting that AEREW

Table 3: Estimated prediction risks of the linear expectile regression with main effects only, the augmented linear expectile regression with interactions, the nonparametric expectile regression via boosting, the five-fold cross-validation, and AEREW ($\lambda = 1$) for the personal computer data. The measures are averaged over 100 random splits of the data and their corresponding standard errors are included in the parentheses. Proportions the candidate estimators being selected by the cross-validation are given by p_{CV} . All numbers are of order 10^{-3} except those corresponding to p_{CV} .

τ	Measure	Individual			Cross-validation	Aggregation
		Linear	Interaction	Boosting		
0.05	risk	2.074 (0.005)	1.905 (0.005)	2.066 (0.009)	1.905 (0.005)	1.787 (0.005)
	p_{CV}	0.00	1.00	0.00	–	–
0.10	risk	3.327 (0.007)	3.069 (0.006)	3.046 (0.012)	3.065 (0.009)	2.778 (0.007)
	p_{CV}	0.00	0.53	0.47	–	–
0.25	risk	5.634 (0.010)	5.166 (0.009)	4.733 (0.015)	4.733 (0.015)	4.519 (0.011)
	p_{CV}	0.00	0.00	1.00	–	–
0.50	risk	6.973 (0.011)	6.294 (0.010)	5.558 (0.017)	5.558 (0.017)	5.427 (0.011)
	p_{CV}	0.00	0.00	1.00	–	–
0.75	risk	5.946 (0.012)	5.236 (0.010)	4.721 (0.018)	4.721 (0.018)	4.581 (0.012)
	p_{CV}	0.00	0.00	1.00	–	–
0.90	risk	3.728 (0.009)	3.203 (0.007)	3.066 (0.012)	3.086 (0.013)	2.896 (0.009)
	p_{CV}	0.00	0.13	0.87	–	–
0.95	risk	2.436 (0.005)	2.070 (0.005)	2.126 (0.010)	2.071 (0.006)	1.916 (0.006)
	p_{CV}	0.00	0.91	0.09	–	–




Figure 1: Estimated prediction risks of the linear expectile regression with main effects only, the augmented linear expectile regression with interactions, the nonparametric expectile regression via gradient boosting, the five-fold cross-validation, and AEREW ($\lambda = 1$) based on 100 independent runs for the personal computer data. On the x -axis of each boxplot, “L” represents the linear expectile regression with main effects only, “I” denotes the augmented linear expectile regression with interactions, and “B” stands for the nonparametric expectile regression via gradient boosting. Each boxplot summarizes the results for one expectile level $\tau \in \{0.05, 0.10, 0.25, 0.50, 0.75, 0.90, 0.95\}$.

gives smaller prediction errors than all candidates and the cross-validated estimator at all expectile levels in Table 3.

4.2. S&P 500 data

We demonstrate the application of the AEREW-ts algorithm in risk management through the S&P 500 data and show how the estimation performance of conditional expectiles can be improved from combining several candidate models. In the financial markets, the expectiles are closely related to gain-loss ratios; see equation (1.2). For example, theoretically, the corresponding gain-loss ratios of expectiles at levels $\tau = 0.01, 0.05,$ and 0.10 are respectively 99, 19 and 9. In our analysis, we examine both estimation risks (in terms of the asymmetric squared error) and realized gain-loss ratios of the estimated expectile functions. Two portfolios under consideration are represented by the S&P 500 Index from October 14, 2008 to October 8, 2010 and from October 8, 2010 to May 1, 2017, respectively. In our analysis we look at the logarithmic returns of the two portfolios, which correspond to two time series containing respectively $n = 500$ and $n' = 1650$ trading days. The candidates are the historical method which uses the sample expectile of the most recent 100 observations to predict the conditional expectile of the current observation (HS100), the linear expectile regression with a lag of 20 observations (Linear), and the boosted expectile regression with a lag of 20

observations (Boosting). Both the linear and boosted expectile regression models can be cast as $\mathcal{E}^\tau(y_t|y_i, i < t) = f(y_{t-1}, y_{t-2}, \dots, y_{t-20})$, where f takes a linear or complicated (nonparametric) form. We took $n_0 = 100$ and applied the AEREW-ts algorithm to aggregate the three candidate models using each of the two time series separately. The results are summarized in Table 4. Since in practice large gain-loss ratios are of main interest, the levels $\tau = 0.01, 0.05$, and 0.10 are considered. The following observations can be made:

- (i) For the first series, the linear expectile regression with lag 20 has the lowest estimation risks among the candidates, while HS100 achieves the highest gain-loss ratios. For the second series, the boosted expectile regression with lag 20 is best in terms of both estimation risk and achieved gain-loss ratio. Thus, there is no universally best method among the three candidates.
- (ii) The AEREW-ts algorithm gives similar results under different λ values. This observation is consistent with our findings in the simulation studies (more details can be found in the supplementary file).
- (iii) AEREW-ts achieves gain-loss ratios that are above the nominal levels in all settings, while at the same time it maintains very small esti-

Table 4: Estimation risks and realized gain-loss ratios (G/L) of the linear expectile regression with lag 20, the boosted expectile regression with lag 20, the historical method and AEREW-ts ($\lambda = 0.1, 1, 10$) for the S&P 500 data.

τ	Measure	Individual			Aggregation (λ)		
		Linear	Boosting	HS100	0.1	1	10
Series 1							
0.01	Risk	0.88	0.97	1.41	0.98	0.98	0.98
	G/L	99.00	80.41	147.51	135.04	135.02	134.81
0.05	Risk	3.06	3.29	3.97	3.22	3.22	3.22
	G/L	19.00	17.03	29.39	23.05	23.05	23.00
0.10	Risk	4.72	5.08	5.70	4.92	4.92	4.91
	G/L	9.00	7.87	13.28	10.22	10.22	10.20
Series 2							
0.01	Risk	0.65	0.76	0.85	0.66	0.66	0.66
	G/L	99.00	115.99	75.31	123.06	123.07	123.15
0.05	Risk	1.78	1.67	2.01	1.70	1.70	1.70
	G/L	19.00	20.81	18.28	20.88	20.88	20.89
0.10	Risk	2.57	2.52	2.80	2.52	2.52	2.52
	G/L	9.00	9.67	8.87	9.47	9.47	9.48

mation risks. This demonstrates the adaptivity of AEREW-ts to the best candidate and can even have performance improvement over the best candidate. Also, note that for time series data, it is not very straightforward to use cross-validation, but our aggregation algorithm can be readily applied.

5. Discussion

It is of interest to discuss the connection and difference between the aggregation procedure considered in our article and those in Shan and Yang (2009) and Dalalyan and Tsybakov (2007). Shan and Yang (2009) focuses on the aggregation of quantile regression models, which also employ an asymmetric loss (the check loss) and have important applications in risk

management. We note that the expectile loss is strongly convex, while the check loss is not. Therefore, we could provide risk bounds under the squared error, but it is hard to do so for the quantile regression aggregation. Moreover, in the oracle inequality of Shan and Yang (2009), the remainder term is of order $(n - n_0)^{-1/2}$, while the one in ours is of order $(n - n_0)^{-1}$.

Note that by setting $\tau = 0.5$ in Theorem 1, we get similar risk bound to that in Theorem 2 of Dalalyan and Tsybakov (2007). We point out that the candidate procedures in Dalalyan and Tsybakov (2007) are assumed to be deterministic, which is different from our aggregation procedure that can fit arbitrary model using training data and hence allows stochastic candidates.

Supplementary Materials

The supplementary file contains proofs of the theorems and additional numerical studies.

Acknowledgements

The authors thank the editor, an associate editor and four reviewers for their insightful comments and suggestions that have helped us improve the quality of the paper substantially. This work is supported in part by NSF grant DMS-1505111.

References

- ACERBI, C. and TASCHE, D. (2002). On the coherence of expected shortfall. *Journal of Banking & Finance* **26**, 1487–1503.
- AIGNER, D. J., AMEMIYA, T. and POIRIER, D. J. (1976). On the estimation of production frontiers: maximum likelihood estimation of the parameters of a discontinuous density function. *International Economic Review* **17**, 377–396.
- AKAIKE, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* **19**, 716–723.
- ARTZNER, P., DELBAEN, F., EBER, J.-M. and HEATH, D. (1999). Coherent measures of risk. *Mathematical Finance* **9**, 203–228.
- BELLINI, F. and BIGNOZZI, V. (2015). On elicitable risk measures. *Quantitative Finance* **15**, 725–733.
- BELLINI, F. and DI BERNARDINO, E. (2017). Risk management with expectiles. *The European Journal of Finance* **23**, 487–506.
- BELLINI, F., KLAR, B., MÜLLER, A. and GIANIN, E. R. (2014). Generalized quantiles as risk measures. *Insurance: Mathematics and Economics* **54**, 41–48.
- BERNARDO, A. E. and LEDOIT, O. (2000). Gain, loss, and asset pricing. *Journal of Political Economy* **108**, 144–172.
- CATONI, O. and PICARD, J. (2004). *Statistical learning theory and stochastic optimization:*

REFERENCES35

- Ecole d'Été de Probabilités de Saint-Flour, XXXI-2001*, vol. 1851. Springer.
- DALALYAN, A. S. and TSYBAKOV, A. B. (2007). Aggregation by exponential weighting and sharp oracle inequalities. In *International Conference on Computational Learning Theory*. Springer.
- EFRON, B. (1991). Regression percentiles using asymmetric squared loss. *Statistica Sinica* **55**, 93–125.
- FAN, J. (1992). Design-adaptive nonparametric regression. *Journal of the American Statistical Association* **87**, 998–1004.
- FAN, J. and YAO, Q. (1998). Efficient estimation of conditional variance functions in stochastic regression. *Biometrika* **85**, 645–660.
- GNEITING, T. (2011). Making and evaluating point forecasts. *Journal of the American Statistical Association* **106**, 746–762.
- GUO, M. and HÄRDLE, W. K. (2012). Simultaneous confidence bands for expectile functions. *AStA Advances in Statistical Analysis* **96**, 517–541.
- HEIDENREICH, N.-B., SCHINDLER, A. and SPERLICH, S. (2013). Bandwidth selection for kernel density estimation: a review of fully automatic selectors. *AStA Advances in Statistical Analysis* **97**, 403–433.
- HOETING, J. A., MADIGAN, D., RAFTERY, A. E. and VOLINSKY, C. T. (1999). Bayesian model averaging: A tutorial. *Statistical Science* **14**, 382–401.

REFERENCES36

- KUAN, C.-M., YEH, J.-H. and HSU, Y.-C. (2009). Assessing value at risk with CARE, the conditional autoregressive expectile models. *Journal of Econometrics* **150**, 261–270.
- NEWKEY, W. K. and POWELL, J. L. (1987). Asymmetric least squares estimation and testing. *Econometrica* **55**, 819–847.
- SCHNABEL, S. K. and EILERS, P. H. (2009a). An analysis of life expectancy and economic production using expectile frontier zones. *Demographic Research* **21**, 109–134.
- SCHNABEL, S. K. and EILERS, P. H. (2009b). Optimal expectile smoothing. *Computational Statistics & Data Analysis* **53**, 4168–4177.
- SCHWARZ, G. (1978). Estimating the dimension of a model. *The Annals of Statistics* **6**, 461–464.
- SHAN, K. and YANG, Y. (2009). Combining regression quantile estimators. *Statistica Sinica* **19**, 1171.
- SOBOTKA, F., KAUERMANN, G., WALTRUP, L. S. and KNEIB, T. (2013a). On confidence intervals for semiparametric expectile regression. *Statistics and Computing* **23**, 135–148.
- SOBOTKA, F., RADICE, R., MARRA, G. and KNEIB, T. (2013b). Estimating the relationship between women’s education and fertility in botswana by using an instrumental variable approach to semiparametric expectile regression. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* **62**, 25–45.
- STENGOS, T. and ZACHARIAS, E. (2006). Intertemporal pricing and price discrimination: a semiparametric hedonic analysis of the personal computer market. *Journal of Applied*

REFERENCES37

- Econometrics* **21**, 371–386.
- TAYLOR, J. W. (2008). Estimating value at risk and expected shortfall using expectiles. *Journal of Financial Econometrics* **6**, 231–252.
- VERSHYNIN, R. (2010). Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027v7* .
- YANG, Y. (2001). Adaptive regression by mixing. *Journal of the American Statistical Association* **96**, 574–588.
- YANG, Y. (2004). Combining forecasting procedures: some theoretical results. *Econometric Theory* **20**, 176–222.
- YANG, Y. and ZOU, H. (2015). Nonparametric multiple expectile regression via ER-Boost. *Journal of Statistical Computation and Simulation* **85**, 1442–1458.
- YAO, Q. and TONG, H. (1996). Asymmetric least squares regression estimation: A nonparametric approach. *Journal of Nonparametric Statistics* **6**, 273–292.
- ZIEGEL, J. F. (2016). Coherence and elicibility. *Mathematical Finance* **26**, 901–918.
- Department of Statistics, University of Connecticut, Storrs, CT 06269, USA.
E-mail: yuwen.gu@uconn.edu
- School of Statistics, University of Minnesota, Minneapolis, MN 55455, USA.
E-mail: zouxx019@umn.edu