

Statistica Sinica Preprint No: SS-2018-0013

Title	A Functional Single Index Model
Manuscript ID	SS-2018-0013
URL	http://www.stat.sinica.edu.tw/statistica/
DOI	10.5705/ss.202018.0013
Complete List of Authors	Fei Jiang Seungchul Baek Jiguo Cao and Yanyuan Ma
Corresponding Author	Jiguo Cao
E-mail	Jiguo_cao@sfu.ca
Notice: Accepted version subject to English editing.	

A Functional Single Index Model

Fei Jiang, Seungchul Baek, Jiguo Cao and Yanyuan Ma

University of Hong Kong, University of South Carolina

Simon Fraser University, Penn State University

Abstract: We propose a semiparametric functional single index model to study the relationship between a univariate response and multiple functional covariates. The parametric part of the model integrates the functional linear regression model and the sufficient dimension reduction structure. The nonparametric part of the model allows the response-index dependence or the link function to be unspecified. The B-spline method is used to approximate the coefficient function, which leads to a dimension folding type model. A new kernel regression method is developed to handle the dimension folding model, which allows the efficient estimation of both the index vector and the B-spline coefficients. We also establish the asymptotic properties and semiparametric optimality for the estimators.

Key words and phrases: B-spline, Dimension reduction, Dimension folding, Functional linear model, Functional data analysis, Kernel estimation.

1. Introduction

The National Morbidity, Mortality and Air Pollution Study (NMMAP) is an important study aiming at addressing the uncertainties regarding the association between pollution and health (Samet et al., 2000). In this study, daily measurements of various air pollutants carbon monoxide (CO), nitrogen dioxide (NO₂), sulfur dioxide (SO₂) and ozone (O₃) are collected in different cities over the course of a year. The annual death rate caused by cardiovascular diseases (CVD) is also collected in these cities. Each pollutant has been studied individually with no significant effect detected on the CVD death rate (Cox and Popken, 2015; Turner et al., 2016).

This motivates us to develop a single air pollution index which combines the pollutants in the way that best describes the severity of the overall air pollution level to the CVD death rate. At the same time, we are also interested in discovering the possible time-varying effect of the single air pollution index to the CVD death rate. To achieve these goals, we propose a functional single index model and proceed to devise a novel class of estimators.

More specifically, the NMMAP data contains the measurements of the daily air pollutant concentrations $\mathbf{X}(t) \equiv \{X_1(t), \dots, X_J(t)\}^T$, a J -dimensional functional covariate of $t \in [0, 1]$, and the annual CVD death rate in the subsequent year as the response Y . To measure the overall severeness of the air pollution,

we define the single air pollution index as

$$W(t) = \boldsymbol{\beta}^T \mathbf{X}(t) = \sum_{i=1}^J \beta_i X_i(t),$$

where $\boldsymbol{\beta} = (\beta_1, \dots, \beta_J)^T$ is the vector of weights to various air pollutants, capturing the relative importance of various pollutants in determining the pollution severity. We assume that Y is linked to $W(t)$ through

$$f\{Y|\mathbf{X}(t)\} = f\left\{Y, \int_0^1 W(t)\alpha(t)dt\right\}, \quad (1.1)$$

where f is a conditional probability density function (pdf) or probability mass function (pmf) and is left unspecified. Here, the functional parameter $\alpha(t)$ captures the time-varying effect of the air pollution index on the annual CVD death rate. Note that unlike the usual single index model as discussed in Chen et al. (2011) and Ma (2016), we make the assumption on the conditional distribution, instead of the conditional mean.

The estimation and inference of the functional single index model (1.1) is not simple. The complexity is due to the unspecified bivariate link function f as well as the unknown coefficient function $\alpha(t)$, in addition to the unknown index vector $\boldsymbol{\beta}$. If $\alpha(t)$ had been known, (1.1) would reduce to a central space estimation problem and various methods exist to estimate $\boldsymbol{\beta}$ (Li, 1991; Cook and Weisberg, 1991; Li and Wang, 2007; Ma and Zhu, 2013). If $\boldsymbol{\beta}$ had been known, (1.1) would reduce to a functional dimension reduction problem (Ferré

and Yao, 2003, 2005, 2007). Qu et al. (2016) handle the unknown $\alpha(t)$ by using the reproducing kernel method, where $\alpha(t)$ is approximated by a function in the reproducing kernel space. Based on a similar idea, we approximate $\alpha(t)$ by a spline function, which facilitates the estimation and inference procedures on $\alpha(t)$.

The functional single index model (1.1) is closely related to sufficient dimension reduction modelling, where a response depends on the covariate vector through its linear transformation (Cook, 1998). To this end, we can view $\int_0^1 \mathbf{X}(t)\alpha(t)dt$ as the covariate vector in the classical sufficient dimension reduction model. Moreover, the proposed method also forms an alternative solution for the dimension folding problem (Li et al., 2010), and our method does not require any conditions on the covariates as other methods in the literature do.

In summary, the new model and estimators have the following features: (i) The model contains a single air pollution index to summarize the pollution severity level; (ii) The time varying coefficient helps to provide timely adjusted health advice to the general public; (iii) The flexible relation between the CVD death rate and the overall pollution effect avoids possible model misspecification; (iv) The model extends the sufficient dimension reduction model to handle multi-functional covariates.

2. Methodology

2.1 Model and Identifiability

The identifiability of Model (1.1) is shown in Proposition 1, which is justified in the supplementary material.

Proposition 1. Assume $\alpha(t)$ is continuous and set $\alpha(0) = 1$ and $\beta_J = 1$, then model (1.1) is identifiable.

Under the assumptions in Proposition 1, we approximate $\alpha(t)$ by a B-spline function $\sum_{k=1}^{d_\gamma} \gamma_k B_{rk}(t)$, where $B_{rk}(t)$ is the k th B-spline basis function with order r ($r \geq 0$). Then the functional single index model (1.1) reduces to the model

$$f(Y, \beta^T \mathbf{Z} \gamma) = f \left\{ Y, \sum_{j=1}^J \sum_{k=1}^{d_\gamma} \beta_j \gamma_k \int_0^1 X_j(t) B_{rk}(t) dt \right\}, \quad (2.2)$$

where \mathbf{Z} is a $J \times d_\gamma$ matrix with the (j, k) th entry $Z_{jk} \equiv \int_0^1 X_j(t) B_{rk}(t) dt$, where f is a unknown density function. We then proceed to estimate β and $\gamma = (\gamma_1, \dots, \gamma_{d_\gamma})^T$ simultaneously. Because the B-spline property ensures $B_{r1}(0) = 1$ and $B_{rk}(0) = 0$ for $k > 1$ and our parameterization fixes $\alpha(0) = 1$, we automatically obtain $\gamma_1 = 1$.

Model (2.2) is in the form of the dimension folding models described in Li et al. (2010) in which the predictors, i.e. \mathbf{Z} 's in our form, are matrix-valued. The

covariate matrix is sandwiched between the left and right coefficient vectors, i.e. β and γ in our setting, to generate a univariate quantity. The dimension folding structure reduces the number of parameters of interest. It has been seen in our setting that we only have $d_\gamma + J - 2$ free coefficients to estimate due to the multiplication from both left and right sides of \mathbf{Z} . The dimension folding model has two main advantages over the standard single index model (Li, 1991; Cook and Weisberg, 1991; Li and Wang, 2007), where the matrix covariates \mathbf{Z} 's are vectorized and $\beta_j \gamma_k$ for $j = 1, \dots, J, k = 1, \dots, d_\gamma$ are Jd_γ new coefficients (Ma, 2016): (1) it automatically takes into account the relation among the coefficients; (2) it reduces the dimension of the unknown coefficients from Jd_γ to $d_\gamma + J - 2$, and hence avoids possible extra high dimensional problems when the number of covariates are merely moderately large.

As an improvement of the dimension folding method, our proposed estimation procedure relaxes the additional constraints on the covariate matrix \mathbf{Z} : our procedure does not require that $E\{\mathbf{X} \mid (\gamma \otimes \beta)^T \text{vec}(\mathbf{X})\}$ is a linear function of $(\gamma \otimes \beta)^T \text{vec}(\mathbf{X})$ and that $\text{var}\{\mathbf{X} \mid (\gamma \otimes \beta)^T \text{vec}(\mathbf{X})\}$ does not depend on $(\gamma \otimes \beta)^T \text{vec}(\mathbf{X})$ as being enforced in Li et al. (2010). Here, \otimes stands for the Kronecker product and $\text{vec}(\mathbf{X})$ is the vector formed by the concatenation of the columns of \mathbf{X} . These constraints may be violated and are not assumed to hold for model (2.2) in general.

2.2 Doubly Robust Local Efficient Score

To estimate the parameters, we first derive the analytic form of the efficient score.

Let

$$\begin{aligned} \mathbf{S}_{\text{eff}\beta}(Y, \mathbf{Z}, \beta, \gamma, g) &= [g(Y, \beta^T \mathbf{Z}\gamma) - E\{g(Y, \beta^T \mathbf{Z}\gamma) \mid \beta^T \mathbf{Z}\gamma\}] \quad (2.3) \\ &\quad \times (\mathbf{I}_{J-1}, \mathbf{0})\{\mathbf{Z} - E(\mathbf{Z} \mid \beta^T \mathbf{Z}\gamma)\}\gamma, \\ \mathbf{S}_{\text{eff}\gamma}(Y, \mathbf{Z}, \beta, \gamma, g) &= [g(Y, \beta^T \mathbf{Z}\gamma) - E\{g(Y, \beta^T \mathbf{Z}\gamma) \mid \beta^T \mathbf{Z}\gamma\}] \\ &\quad \times (\mathbf{0}, \mathbf{I}_{d_\gamma-1})\{\mathbf{Z} - E(\mathbf{Z} \mid \beta^T \mathbf{Z}\gamma)\}^T \beta, \end{aligned}$$

where the function $g(Y, \beta^T \mathbf{Z}\gamma) = f'_2(Y, \beta^T \mathbf{Z}\gamma)/f(Y, \beta^T \mathbf{Z}\gamma)$ and $f'_2(Y, \beta^T \mathbf{Z}\gamma)$ is the partial derivative of f with respect to its second argument. Then the efficient score is $\mathbf{S}_{\text{eff}}(Y_i, \mathbf{Z}_i, \beta, \gamma, g) \equiv \{\mathbf{S}_{\text{eff}\beta}(Y, \mathbf{Z}, \beta, \gamma, g)^T, \mathbf{S}_{\text{eff}\gamma}(Y, \mathbf{Z}, \beta, \gamma, g)^T\}^T$.

Our hope is to use the efficient score to construct estimating equation hence to solve for β, γ from

$$\sum_{i=1}^n \mathbf{S}_{\text{eff}}(Y_i, \mathbf{Z}_i, \beta, \gamma, g) = \mathbf{0}. \quad (2.4)$$

When $g(\cdot)$ and $E(\cdot \mid \beta^T \mathbf{Z}\gamma)$ are correctly specified, \mathbf{S}_{eff} is indeed a function which falls in the space that is orthogonal to the nuisance tangent space induced by the unknown conditional density $f(\cdot)$ defined in Proposition 3. Hence, as shown in Bickel et al. (1993); Tsiatis (2004), \mathbf{S}_{eff} is the efficient score (see Proposition S4.1 in the supplementary document) which yields the optimal estimators

with the smallest asymptotic variances. In addition, S_{eff} is also a doubly robust function so that the estimation consistency holds whenever $E\{g(Y, \beta^T \mathbf{Z}\gamma) \mid \beta^T \mathbf{Z}\gamma\}$ or $E(\mathbf{Z} \mid \beta^T \mathbf{Z}\gamma)$ is correctly specified (Ma and Zhu, 2012, 2013).

In reality, the functional form for the conditional density $f(\cdot)$ is usually unknown, this leads to the difficulty of obtaining $E(\cdot \mid \beta^T \mathbf{Z}\gamma)$ and $f'_2(Y, \beta^T \mathbf{Z}\gamma)/f(Y, \beta^T \mathbf{Z}\gamma)$ in (2.3), hence the efficient score cannot be directly used. To retain the best estimation efficiency without imposing more assumptions, we adopt nonparametric estimation to estimate the unknown components in the efficient score function. Specifically, we use the standard kernel smoothing method in nonparametric regression to estimate $E(\cdot \mid \beta^T \mathbf{Z}\gamma)$, i.e.

$$\widehat{E}\{m(Y_i, \mathbf{Z}_i) \mid \beta^T \mathbf{Z}\gamma\} = \frac{\sum_{i=1}^n m(Y_i, \mathbf{Z}_i) K_h(\beta^T \mathbf{Z}_i \gamma - \beta^T \mathbf{Z}\gamma)}{\sum_{i=1}^n K_h(\beta^T \mathbf{Z}_i \gamma - \beta^T \mathbf{Z}\gamma)}, \quad (2.5)$$

for arbitrary function $m(Y_i, \mathbf{Z}_i)$. Here $K(\cdot)$ is a kernel function and $K_h(\cdot) = h^{-1}K(\cdot/h)$. To estimate $f(Y, \beta^T \mathbf{Z}_i \gamma)$, we use a local linear estimator. Specifically, we obtain the estimators $\widehat{f}(Y, \beta^T \mathbf{Z}_i \gamma) = c_0$, and $\partial \widehat{f}(Y, \beta^T \mathbf{Z}_i \gamma) / \partial (\beta^T \mathbf{Z}_i \gamma) = c_1$ from minimizing

$$\sum_{j=1}^J \{K_b(Y_j - Y) - c_0 - c_1(\beta^T \mathbf{Z}_j \gamma - \beta^T \mathbf{Z}_i \gamma)\}^2 K_h(\beta^T \mathbf{Z}_j \gamma - \beta^T \mathbf{Z}_i \gamma) \quad (2.6)$$

with respect to c_0, c_1 . (2.5) and (2.6) allow us to obtain the unknown quantities in S_{eff} consistently given any parameter β, γ .

It is clearly a profiling procedure, where unknown nuisance components are

estimated as functions of the parameters of interest and then the estimating equations are solved to obtain the final estimator. This procedure yields the optimal estimator for β, γ , while it has the drawback that it requires quite heavy computation, especially in solving (2.6) for each value of $\beta^T \mathbf{Z}_j \gamma$. Thus, when the estimation variability is not of great concern, to ease the computation, we may aim for a possibly non-optimal estimator. Specifically, we may posit a working model for f and f_2 , say f^* and f_2^* . Let $g^* = f_2^*/f^*$, then $\mathbf{S}_{\text{eff}}(Y_i, \mathbf{Z}_i, \beta, \gamma, g^*)$ is a locally efficient score function. Using the locally efficient score function to construct estimating equations guarantees estimation consistency, while it can result in efficient estimator when g^* is the truth.

It is also worth pointing out the issue of obtaining \mathbf{Z}_i 's. Unlike in the usual dimension reduction problems, \mathbf{Z}_i 's are not directly observed and need to be constructed from the observed $X_j(t)$'s. This involves a numerical approximation of the integrals $\int_0^1 B_{rk}(t)X_j(t)dt$. The composite Simpson's rule (Atkinson, 1989) can be used to approximate the numerical integration, which has the form

$$\int_0^1 B_{rk}(t)X_j(t)dt = \frac{1}{3Q} \left[B_{rk}(t_0)X_j(t_0) + 2 \sum_{q=1}^{Q/2-1} \{B_{rk}(t_{2q})X_j(t_{2q})\} + 4 \sum_{q=1}^{Q/2} \{B_{rk}(t_{2q-1})X_j(t_{2q-1})\} + B_{rk}(t_Q)X_j(t_Q) \right],$$

where $t_q = q/Q$, $q = 0, 1, \dots, Q$, and Q is an even number.

The estimation procedures can be summarized as follows:

Step 1: Choose f and f'_2 by minimizing (2.6) or positing f and f'_2 . Denote the choices by $f^*(Y, \beta^T \mathbf{Z}\gamma, \beta, \gamma)$ and $f'^*_2(Y, \beta^T \mathbf{Z}\gamma, \beta, \gamma)$ and let $g^*(Y, \beta^T \mathbf{Z}\gamma, \beta, \gamma) = f'^*_2(Y, \beta^T \mathbf{Z}\gamma, \beta, \gamma) / f^*(Y, \beta^T \mathbf{Z}\gamma, \beta, \gamma)$.

Step 2: Replace g in (2.4) by g^* according to the **Step 1** choice.

Step 3: Let $\widehat{\mathbf{S}}_{\text{eff}\gamma}$ be the version of $\mathbf{S}_{\text{eff}\gamma}$ when replacing $E(\cdot | \beta^T \mathbf{Z}\gamma)$ by $\widehat{E}(\cdot | \beta^T \mathbf{Z}\gamma)$ defined in (2.5). Treating γ as a function of β , denoted by $\gamma(\beta)$, we solve

$$\sum_{i=1}^n \widehat{\mathbf{S}}_{\text{eff}\gamma}[Y_i, \mathbf{Z}_i, \beta, \gamma(\beta), g^*\{Y_i, \beta^T \mathbf{Z}_i \gamma, \beta, \gamma(\beta)\}] = \mathbf{0}$$

for $\gamma(\beta)$, and denote the estimator as $\widehat{\gamma}(\beta)$.

Step 4: Let $\widehat{\mathbf{S}}_{\text{eff}\beta}$ be the version of $\mathbf{S}_{\text{eff}\beta}$ when replacing $E(\cdot | \beta^T \mathbf{Z}\gamma)$ by $\widehat{E}(\cdot | \beta^T \mathbf{Z}\gamma)$.

We solve

$$\sum_{i=1}^n \widehat{\mathbf{S}}_{\text{eff}\beta}[Y_i, \mathbf{Z}_i, \beta, \widehat{\gamma}(\beta), g^*\{Y_i, \beta^T \mathbf{Z}_i \gamma, \beta, \widehat{\gamma}(\beta)\}] = \mathbf{0}$$

for β , and denote the estimator as $\widehat{\beta}$.

In the algorithm, we used the last two arguments in $g^*(Y_i, \beta^T \mathbf{Z}_i \gamma, \beta, \gamma)$ to emphasize its possible dependence on β, γ . Obviously, when we posit a model for f , the functional form does not have to depend on β, γ , while when we estimate the model f , the functional form certainly depends on β, γ as it is the case in all profiling estimators. The resulting estimators are consistent as discussed in Proposition 2, because the expectations of the score functions have zero mean when the parameters are correctly specified. In estimating $\widehat{E}(\cdot | \beta^T \mathbf{Z}\gamma)$, we used

the variance of $\beta^T \mathbf{Z} \gamma$ times $n^{-1/5}$ as bandwidth. We find that the results are robust in the range between half of this bandwidth to twice of it.

2.3 Asymptotic Results

The profiling procedures Step 3 and 4 yield the estimators which are asymptotically equivalent to the ones from solving the estimating equation based on the estimating function $(\widehat{\mathbf{S}}_{\text{eff}\gamma}^T, \widehat{\mathbf{S}}_{\text{eff}\beta}^T)^T$. Hence the estimation consistency readily holds by the following proposition.

Proposition 2. Let $\widehat{\beta}, \widehat{\gamma}$ satisfy

$$\sum_{i=1}^n \{ \widehat{\mathbf{S}}_{\text{eff}\beta}(Y_i, \mathbf{Z}_i, \widehat{\beta}, \widehat{\gamma}, g^*)^T, \widehat{\mathbf{S}}_{\text{eff}\gamma}(Y_i, \mathbf{Z}_i, \widehat{\beta}, \widehat{\gamma}, g^*)^T \}^T = \mathbf{0}$$

where

$$\begin{aligned} & \widehat{\mathbf{S}}_{\text{eff}\beta}(Y_i, \mathbf{Z}_i, \widehat{\beta}, \widehat{\gamma}, g^*) \\ &= \left\{ g^*(Y_i, \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) - \frac{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) g^*(Y_j, \widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma})}{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma})} \right\} \Theta_{\beta} \\ & \times \left\{ \mathbf{Z}_i - \frac{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) \mathbf{Z}_j}{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma})} \right\} \widehat{\gamma}, \\ & \widehat{\mathbf{S}}_{\text{eff}\gamma}(Y_i, \mathbf{Z}_i, \widehat{\beta}, \widehat{\gamma}, g^*) \\ &= \left\{ g^*(Y_i, \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) - \frac{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) g^*(Y_j, \widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma})}{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma})} \right\} \Theta_{\gamma} \\ & \left\{ \mathbf{Z}_i - \frac{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma}) \mathbf{Z}_j}{\sum_{j=1}^J K_h(\widehat{\beta}^T \mathbf{Z}_j \widehat{\gamma} - \widehat{\beta}^T \mathbf{Z}_i \widehat{\gamma})} \right\}^T \widehat{\beta}. \end{aligned}$$

Let β_0 be the true β . Further, let γ_0 be a spline coefficient which satisfies $\sup_{t \in [0,1]} |\mathbf{B}_r(t)^\top \gamma_0 - \alpha_0(t)| = O_p(h_b^q)$ as stated in Condition (A5). Then $\widehat{\beta} - \beta_0 = o_p(1)$, $\sup_{t \in [0,1]} |\mathbf{B}_r(\cdot)^\top \widehat{\gamma} - \mathbf{B}_r(\cdot)^\top \gamma_0| = o_p(1)$.

In Step 3, at the point of convergence, we show that $\widehat{\gamma}(\beta_0)$ achieves the non-parametric spline regression convergence rate, and derive its asymptotic variation as follows.

Theorem 1. Assume Conditions (A1)–(A8) hold, and let $\mathbf{B}_r(\cdot)^\top \widehat{\gamma}(\beta_0)$ satisfy

$$\sum_{i=1}^n \widehat{\mathbf{S}}_{\text{eff}\gamma}(Y_i, \mathbf{Z}_i, \beta_0, \widehat{\gamma}(\beta_0), g^*) = \mathbf{0}.$$

Then $n^{1/2}\{\widehat{\gamma}^-(\beta_0) - \gamma_0^-\} = \mathbf{L} + o_p(\mathbf{L})$, where

$$\begin{aligned} \mathbf{L} = & - \left(\left\{ E \left(\Theta_\gamma [\mathbf{Z}_i - E\{\mathbf{Z}_i | \beta_0^\top \alpha_0(\mathbf{X}_i)\}]^\top \beta_0 \right. \right. \\ & \times \frac{\partial (g^*\{Y_i, \beta_0^\top \alpha_0(\mathbf{X}_i)\} - E[g^*\{Y_i, \beta_0^\top \alpha_0(\mathbf{X}_i)\} | \beta_0^\top \alpha_0(\mathbf{X}_i)])}{\partial \beta_0^\top \alpha_0(\mathbf{X}_i)} \\ & \left. \left. \times \beta_0^\top [\mathbf{Z}_i - E\{\mathbf{Z}_i | \beta_0^\top \alpha_0(\mathbf{X}_i)\}] \Theta_\gamma^\top \right\}^{-1} \right) \left(n^{-1/2} \sum_{i=1}^n [g^*\{Y_i, \beta_0^\top \alpha_0(\mathbf{X}_i)\} \right. \\ & \left. - E\{g^*\{Y_i, \beta_0^\top \alpha_0(\mathbf{X}_i)\} | \beta_0^\top \alpha_0(\mathbf{X}_i)\}] \Theta_\gamma [\mathbf{Z}_i - E\{\mathbf{Z}_i | \beta_0^\top \alpha_0(\mathbf{X}_i)\}]^\top \beta_0 \right). \end{aligned}$$

Here $\gamma_0^- = (\gamma_{02}, \dots, \gamma_{0d_\gamma})^\top$, and $\widehat{\gamma}^-(\beta_0) = (\widehat{\gamma}_2(\beta_0), \dots, \widehat{\gamma}_{d_\gamma}(\beta_0))^\top$. Further, for arbitrary $d_{\gamma-1}$ -dimensional vector with $\|\mathbf{a}\|_2 < \infty$, we have $\mathbf{a}^\top \{\widehat{\gamma}^-(\beta_0) - \gamma_0^-\} = O_p\{(nh_b)^{-1/2}\}$.

In addition, we show $\widehat{\beta}$ from Step 4 is not only root- n consistent, but also efficient and achieve the information lower bound $\{E(\mathbf{S}_{\text{oeff}}^{\otimes 2})\}^{-1}$. Here \mathbf{S}_{oeff} is

the efficient score for β of the original model (1.1), which contains $\alpha(\cdot)$ instead of $\mathbf{B}_r(\cdot)^T \gamma$, hence it is different from $\mathbf{S}_{\text{eff}\beta}$. Its precise expression is given in Proposition S4.2 in the supplementary document.

To show the asymptotic properties of $\hat{\beta}$, we first define

$$\begin{aligned} & \Delta g_c^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} \\ \equiv & \frac{\partial [g^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} - E\{g^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} | \beta_0^T \alpha_0(\mathbf{X}_i)\}]}{\partial \{\beta_0^T \alpha_0(\mathbf{X}_i)\}} \end{aligned}$$

and $\mathbf{w}_0^*(t)$ as a function that satisfies

$$\begin{aligned} & \Theta_\beta E[\alpha_{c0}(\mathbf{X}_i) \Delta g_c^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} \mathbf{X}_{ic}^T(t)] \beta_0 \quad (2.7) \\ = & \int_0^1 E[\beta_0^T \mathbf{X}_{ic}(s) \Delta g_c^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} \mathbf{X}_{ic}^T(t) \beta_0] \mathbf{w}_0^*(s) ds, \end{aligned}$$

where $\alpha_{c0}(\mathbf{X}) \equiv \alpha_0(\mathbf{X}) - E\{\alpha_0(\mathbf{X}) | \beta_0^T \alpha_0(\mathbf{X})\}$. We have the following results.

Theorem 2. Assume Conditions (A1)–(A8) hold, and let $\hat{\beta}$ satisfy

$$\sum_{i=1}^n \hat{\mathbf{S}}_{\text{eff}\beta}(Y_i, \mathbf{Z}_i, \hat{\beta}, \hat{\gamma}(\hat{\beta}), g^*) = \mathbf{0}.$$

Then $\sqrt{n}(\hat{\beta} - \beta_0) = \mathbf{A}^{-1} \mathbf{B} + o_p(1)$, where

$$\begin{aligned} \mathbf{A} = & -E \left[\Delta g_c^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} \{\Theta_\beta \alpha_{c0}(\mathbf{X}_i)\}^{\otimes 2} \right. \\ & \left. - \Delta g_c^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} \int_0^1 \beta_0^T \mathbf{X}_{ic}(t) \mathbf{w}_0^*(t) dt \alpha_{c0}(\mathbf{X}_i)^T \Theta_\beta^T \right] \end{aligned}$$

and

$$\mathbf{B} = n^{-1/2} \sum_{i=1}^n \left\{ g^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} - E [g^* \{Y_i, \beta_0^T \alpha_0(\mathbf{X}_i)\} | \beta_0^T \alpha_0(\mathbf{X}_i)] \right\} \\ \times \left\{ \Theta_{\beta} \alpha_{c0}(\mathbf{X}_i) - \int_0^1 \beta_0^T \mathbf{X}_{ic}(t) \mathbf{w}_0^*(t) dt \right\}.$$

Hence, $\sqrt{n}(\hat{\beta} - \beta_0)$ converges to a normal distribution with mean $\mathbf{0}$ and variance Σ , where $\Sigma \equiv \mathbf{A}^{-1} E(\mathbf{B}^{\otimes 2}) \mathbf{A}^{-1T}$. Here $\mathbf{a}^{\otimes 2} = \mathbf{a} \mathbf{a}^T$ for arbitrary vector or matrix \mathbf{a} . In addition, when $g^* = g$, $\sqrt{n}(\hat{\beta} - \beta_0)$ converges to a normal distribution with mean $\mathbf{0}$ and variance $\{E(\mathbf{S}_{\text{eff}}^{\otimes 2})\}^{-1}$, i.e., $\hat{\beta}$ is the semiparametric efficient estimator of β for model (1.1).

Theorem 2 indicates that $\hat{\beta}$ is the consistent estimator. Further it is semiparametric efficient when g^* is correctly specified, even though the estimation of $\hat{\beta}$ is devised under the approximate model (2.2). Generally, we can replace g^* by a consistent estimator of g . The following corollary ensures the asymptotic efficiency of the resulting $\hat{\beta}$.

Corollary 1. Assume Conditions (A1)–(A8) hold, and let $\hat{\beta}$ satisfy

$$\sum_{i=1}^n \hat{\mathbf{S}}_{\text{eff}\beta}(Y_i, \mathbf{Z}_i, \hat{\beta}, \hat{\gamma}(\hat{\beta}), \hat{g}) = \mathbf{0},$$

where \hat{g} is a uniformly consistent estimator for the true function g , and $\hat{\mathbf{S}}_{\text{eff}\beta}$ is defined in Proposition 2, then $\hat{\beta}$ is semiparametric efficient.

The corollary readily holds by using the result in Theorem 2 and the consistency of \hat{g} . We omit the details. In practice, we can use kernel method to estimate $f(Y, \beta^T \mathbf{Z} \gamma)$ and in turn to obtain \hat{g} , which is guaranteed to be uniformly consistent to g (Mack and Silverman, 1982). Combining the results of Theorems 1 and 2, we are able to establish the theoretical properties of the estimation of $\hat{\alpha}(t)$ in Theorem 3. Specifically, Theorem 3 shows that the spline approximation $\hat{\alpha}(t) = \mathbf{B}_r(t)^T \hat{\gamma}(\hat{\beta})$ with β, γ estimated using the estimating equation set (3.8) indeed achieves the usual nonparametric spline regression convergence rate.

Theorem 3. *Assume Conditions (A1) – (A8) hold, then*

$$\sup_{t \in [0,1]} |\mathbf{B}_r(t)^T \hat{\gamma}(\hat{\beta}) - \alpha_0(t)| = O_p(n^{-1/2} h_b^{-1/2}).$$

The proofs for the theoretic results are elaborated in the supplementary document.

3. Relation to the semiparametric sufficient dimension reduction

Although performed for the functional model and the dimension folding model, the proposed estimation procedure is inline with the semiparametric sufficient dimension reduction techniques discussed in Ma and Zhu (2012). To illustrate the similarity, following Bickel et al. (1993) and Tsiatis (2004), we first develop the nuisance tangent space Λ^\perp in the following proposition, which allows us to

construct estimators of β, γ from various choices of the function \mathbf{f} appeared in the description of Λ^\perp .

Proposition 3. In the Hilbert space \mathcal{H} of all the mean zero finite variance functions associated with (2.2), i.e. $\mathcal{H} = \{\mathbf{a}(\mathbf{Z}, Y) : \int \mathbf{a}(\mathbf{z}, y) f(y, \beta^\top \mathbf{z} \gamma) f_{\mathbf{Z}}(\mathbf{z}) d\mu(\mathbf{z}, y) = \mathbf{0}, \int \mathbf{a}^\top(\mathbf{z}, y) \mathbf{a}(\mathbf{z}, y) f(y, \beta^\top \mathbf{z} \gamma) f_{\mathbf{Z}}(\mathbf{z}) d\mu(\mathbf{z}, y) < \infty, \mathbf{a}(\mathbf{z}, y) \in \mathcal{R}^{d_\gamma + J - 2}\}$, where $\mu(\mathbf{z}, y)$ is the probability measure of (\mathbf{Z}, Y) , $f_{\mathbf{Z}}(\mathbf{z})$ is the pdf of \mathbf{Z} and $f(y, \beta^\top \mathbf{z} \gamma)$ is given in (2.2), the orthogonal complement of the nuisance tangent space is

$$\Lambda^\perp = \{\mathbf{f}(Y, \mathbf{Z}) - E(\mathbf{f} | Y, \beta^\top \mathbf{Z} \gamma) : E(\mathbf{f} | \mathbf{Z}) = E(\mathbf{f} | \beta^\top \mathbf{Z} \gamma), \forall \mathbf{f}\}.$$

The proof of Proposition 3 is given in the supplementary document. Let $\mathbf{f}(Y, \mathbf{Z}) = [\mathbf{g}(Y, \beta^\top \mathbf{Z} \gamma) - E\{\mathbf{g}(Y, \beta^\top \mathbf{Z} \gamma) | \beta^\top \mathbf{Z} \gamma\}] \mathbf{a}(\mathbf{Z})$, where \mathbf{g}, \mathbf{a} can be chosen arbitrarily as long as the resulting \mathbf{f} contains sufficiently many equations. Obviously, $E(\mathbf{f} | \mathbf{Z}) = \mathbf{0}$ hence $E(\mathbf{f} | \beta^\top \mathbf{Z} \gamma) = \mathbf{0}$, and $\mathbf{f} - E(\mathbf{f} | Y, \beta^\top \mathbf{Z} \gamma) = \{\mathbf{g} - E(\mathbf{g} | \beta^\top \mathbf{Z} \gamma)\} \{\mathbf{a} - E(\mathbf{a} | \beta^\top \mathbf{Z} \gamma)\}$. Thus, we can construct estimating equation based on the sample version of

$$\begin{aligned} & E([\mathbf{g}(Y, \beta^\top \mathbf{Z} \gamma) - E\{\mathbf{g}(Y, \beta^\top \mathbf{Z} \gamma) | \beta^\top \mathbf{Z} \gamma\}][\mathbf{a}(\mathbf{Z}) - E\{\mathbf{a}(\mathbf{Z}) | \beta^\top \mathbf{Z} \gamma\}]) \\ &= \mathbf{0}, \end{aligned} \tag{3.8}$$

and it provides a class of estimators for β, γ .

We now perform a set of analysis somewhat in the spirit of Ma and Zhu (2012) to illustrate that by different choice of \mathbf{g} and \mathbf{a} , (3.8) leads to the classical

dimension reduction estimators.

3.1 The relation with the sliced inverse regression

As a first choice of \mathbf{g} and \mathbf{a} , let $\mathbf{V} = \text{vec}(\mathbf{Z})$, and select $\mathbf{g}(Y, \beta^T \mathbf{Z} \gamma) = E(\mathbf{V} | Y)$, $\mathbf{a}(\mathbf{Z}) = \mathbf{V}^T$. This provides an estimator with the flavor of the sliced inverse regression (SIR, Li (1991)) in the classical dimension reduction framework. Specifically, under this choice of \mathbf{g} , \mathbf{a} , (3.8) has the form

$$E \left([E(\mathbf{V} | Y) - E\{E(\mathbf{V} | \mathbf{Y}) | \beta^T \mathbf{Z} \gamma\}] \{ \mathbf{V}^T - E(\mathbf{V}^T | \beta^T \mathbf{Z} \gamma) \} \right) = \mathbf{0}. \quad (3.9)$$

The above estimating equation set contains $J^2 d_\gamma^2$ equations while we only have $J + d_\gamma - 2$ free parameters. We can use GMM to reduce the number of equations in practice. We can also construct \mathbf{g} or \mathbf{a} or both using only a subset of \mathbf{V} .

3.2 The relation with the sliced average variance estimator

As a second choice of \mathbf{g} , \mathbf{a} , we select $\mathbf{g}_1(Y, \beta^T \mathbf{Z} \gamma) = \mathbf{I}_{J d_\gamma} - \text{cov}(\mathbf{V} | Y)$, $\mathbf{g}_2(Y, \beta^T \mathbf{Z} \gamma) = \mathbf{g}_1 E(\mathbf{V} | Y)$, $\mathbf{a}_1(\mathbf{Z}) = -\mathbf{V} \{ \mathbf{V} - E(\mathbf{V} | \beta^T \mathbf{Z} \gamma) \}^T$, $\mathbf{a}_2(\mathbf{Z}) = \mathbf{V}^T$. We then construct a classical sliced average variance estimator (SAVE, Cook and Weisberg (1991)) flavored estimator based on

$$\begin{aligned} & E \left[\{ \mathbf{g}_1 - E(\mathbf{g}_1 | \beta^T \mathbf{Z} \gamma) \} \{ \mathbf{a}_1 - E(\mathbf{a}_1 | \beta^T \mathbf{Z} \gamma) \} \right] \\ & + E \left[\{ \mathbf{g}_2 - E(\mathbf{g}_2 | \beta^T \mathbf{Z} \gamma) \} \{ \mathbf{a}_2 - E(\mathbf{a}_2 | \beta^T \mathbf{Z} \gamma) \} \right] = \mathbf{0}. \end{aligned} \quad (3.10)$$

3.3 The relation with the directional regression

The third choice of \mathbf{g} , \mathbf{a} that we would like to present is $\mathbf{g}_1(Y, \beta^T \mathbf{Z} \gamma) = \mathbf{I}_{Jd_\gamma} - E(\mathbf{V}\mathbf{V}^T | Y)$, $\mathbf{g}_2(Y, \beta^T \mathbf{Z} \gamma) = E\{E(\mathbf{V} | Y)E(\mathbf{V}^T | Y)\}E(\mathbf{V} | Y)$, $\mathbf{g}_3(Y, \beta^T \mathbf{Z} \gamma) = E\{E(\mathbf{V}^T | Y)E(\mathbf{V} | Y)\}E(\mathbf{V} | Y)$, $\mathbf{a}_1(\mathbf{Z}) = -\mathbf{V}\{\mathbf{V} - E(\mathbf{V} | \beta^T \mathbf{Z} \gamma)\}^T$, $\mathbf{a}_2(\mathbf{Z}) = \mathbf{a}_3(\mathbf{Z}) = \mathbf{V}^T$. This leads to a classic directional regression (DR, Li and Wang (2007)) flavored estimator from

$$\sum_{i=1}^3 E [\{\mathbf{g}_i - E(\mathbf{g}_i | \beta^T \mathbf{Z} \gamma)\}\{\mathbf{a}_i - E(\mathbf{a}_i | \beta^T \mathbf{Z} \gamma)\}] = \mathbf{0}. \quad (3.11)$$

The three estimators given in (3.9), (3.10) and (3.11) are respectively of the flavours of SIR, SAVE and DR, because if we had worked in the classical sufficient dimension reduction context, and if further equipped with the additional linearity condition and constant variance condition, the choices of \mathbf{g} and \mathbf{a} that led to the three estimating equations above would have further led to SIR, SAVE and DR (Ma and Zhu, 2012). Further, the choices of \mathbf{g} , \mathbf{a} in (3.9), (3.10) and (3.11) only depend on the moments of \mathbf{Z} instead of the conditional density as the one used in \mathbf{S}_{eff} defined in (2.3). Hence, these estimators can serve as alternatives to the proposed efficient estimators when the conditional density is difficult to obtain.

4. Simulation Studies

We carry out three simulation studies under the following settings in order to assess the finite sample performance of our estimation method. In each simulation, we generate 1000 data sets with the sample size $n = 500$.

Simulation 1

(1) $J = 9$, $\beta = (1, 1.2, 1.5, 0.5, -0.5, -1.5, -1.2, -1, 1)^T$ $\alpha_0(t) = \sin(\pi t) + 1$, $t \in [0, 1]$;

(2) $X_{ji}(t)$, $j = 1 \dots, 4$ follows $U(-5, 5)$, where $U[a, b]$ denotes a random variable from the uniform distribution in the range of $[a, b]$;

(3) Y_i follows a normal distribution with mean $\int_0^1 W_i(t)\alpha_0(t)dt$ and variance 1, where $W_i(t) = \beta^T \mathbf{X}_i(t)$.

Simulation 2 investigates the ability of our methods in handling the nonlinear mean and variance.

(1) Y_i follows a normal distribution with mean $\sin\{2 \int_0^1 W_i(t)\alpha_0(t)dt\} + \log[1 + \{\int_0^1 W_i(t)\alpha_0(t)dt\}^2] - 3$ and variance $0.5[1 + \{\beta^T \int_0^1 \mathbf{X}_i(t)\alpha_0(t)dt\}^2]^{1/5}$.

Simulation 3 resembles the air pollution data structure in Section 5.

(1) $J = 4$, $\beta = (-0.2, -1, -1.5, 1)^T$ and $\alpha_0(t) = 1 - 26.76t + 145.3t^2 - 227.27t^3 + 107.99t^4$;

(2) $X_{i1}(t) = 0.66 - 4.84t + 5.12t^2 + U[-4, 6]$, $X_{i2}(t) = 0.43 - 2.95t +$

$3.11t^2 + U[-5, 5]$, $X_{i3}(t) = -1.61 + 10.40t - 10.85t^2 + U[-4, 4]$, $X_{i4}(t) = 0.58 - 3.59t + 3.52t^2 + U[-4, 8]$;

(3) Y_i is from a normal distribution with mean $0.075 + 0.53(\int_0^1 W_i(t)\alpha_0(t)dt + 1.23)$ and variance 0.05.

We applied the proposed method to estimate both β and $\alpha(t)$, where $\alpha(t)$ is approximated with cubic B-spline basis functions using three equally spaced internal knots. For comparison, we implemented three estimators: the oracle, the efficient, and the locally efficient estimators. In the oracle estimator, we specified $f(Y, \beta^T \mathbf{Z}\gamma)$ using the normal pdf form, and used the true $g(Y, \beta^T \mathbf{Z}\gamma)$ in the estimation. In the efficient estimator, $E(\cdot | \beta^T \mathbf{Z}\gamma)$, $f(Y, \beta^T \mathbf{Z}\gamma)$, and $f'_2(Y, \beta^T \mathbf{Z}\gamma)$ are estimated via nonparametric method. In the local estimator, we specified an incorrect model of $f(Y, \beta^T \mathbf{Z}\gamma)$, hence a misspecified $g^*(Y, \beta^T \mathbf{Z}\gamma)$ function was used, and estimated $E(\cdot | \beta^T \mathbf{Z}\gamma)$ nonparametrically. Note that the form of $f(Y, \beta^T \mathbf{Z}\gamma)$ is generally unknown, hence the oracle estimator is unrealistic and is only included here as a benchmark for a comparison purpose.

The numerical performances of the estimation of β in Simulation 1, 2 and 3 are summarized in Table 1, 2 and 3, respectively. Based on the asymptotic results in Theorem 2, the average of the estimated standard error is obtained and the coverage of the 95% confidence interval is also provided. As expected, both the efficient and the locally efficient estimators have very small bias, the esti-

mated variances are close to their empirical ones, and the 95% coverage are also reasonably close to the nominal level. The variances of the efficient estimators are smaller than those of the locally efficient estimators. In fact, the performance of the efficient estimators is very close to the oracle estimators. We illustrate the performance of the estimation of $\alpha_0(t)$ in Figure 1, where we show the mean estimated curves and the pointwise 90% confidence bands. The performance shown in Figure 1 is rather typical for spline approximations.

In functional data analysis, a simple stacking approach is often used to study the effect of the functional covariates (Ramsay and Silverman, 2005) in a less structured model

$$E\{Y|\mathbf{X}(t)\} = \int_0^1 \mathbf{X}(t)^T \boldsymbol{\eta}(t) dt, \quad (4.12)$$

where $\boldsymbol{\eta}(t) = \{\eta_1(t), \dots, \eta_p(t)\}^T$. The stacking approach is a special case of the proposed functional single index model. We thus implemented the stacking approach and compared the two estimators in Figure 2. It is easy to see that our estimator performs better than the stacking approach, with narrower confidence bands. This pattern also applies to simulations 2 and 3, and we provide the corresponding plots in Figures S1 and S2 of the supplementary document.

5. Application

We apply the proposed method to study the effect of various air pollutants on the rate of death caused by CVD, where we adopt the model in (1.1) without specifying any special link function.

In the NMMAP data (Peng and Welty, 2004), all four pollutants (CO, NO₂, SO₂, and O₃) were recorded on a daily basis in 108 U.S. cities. The measurements unit is parts per billion (ppb) by volume and spans the range from 1987 to 2000. 400 observations with a relatively small portion of missing values are used for the analysis and each observation has 365 daily median measurements of four air pollutants, where we imputed a few missing days in some observations by linear interpolation. We also standardized each pollutant across the whole year so that the 365 observations yield a sample mean 0 and sample variance 1. The time interval is normalized to $[0, 1]$. Figure S3 of the supplementary document displays the mean trajectories for four pollutants.

We fit model (1.1) to estimate the air pollution index directly related to the following year's CVD death rate. Throughout the implementation, we set the kernel bandwidth h to be $n^{-1/5}\text{range}(\beta^T \mathbf{z}_i \gamma)$ and $b = n^{-1/7}\text{range}(y_i)$, where the unknown parameters β and γ are updated during each iteration. The functional parameter $\alpha(t)$ is estimated using a linear combination of cubic B-splines with three equally spaced internal knots in $[0, 1]$, where the optimal number of

internal knots is determined through a ten-fold cross-validation. We calculated the confidence band for $\alpha(t)$ using the asymptotic results in Theorem 1.

Figure 3 shows the time-varying effect $\hat{\alpha}(t)$ of the estimated air pollution index to the CVD death rate. The time-varying effect of the air pollution index is significantly positive in the spring, summer, and fall seasons but insignificant in the winter. The air pollution index has the largest positive effect on the CVD death rate in the summer.

Figure 4 displays the air pollution index for three major cities: Boston, New York and Chicago together with their CVD annual death rates. With the largest air pollution index in the summertime, New York has the largest CVD death rate. On the other hand, Boston has the lowest air pollution index in the summer and hence the CVD death rate is the smallest in Boston, although Boston has the highest air pollution index in the winter.

Table 4 displays the estimated coefficients for all four air pollutants β . The standard errors and the p -values are obtained based on the asymptotic normality of $\hat{\beta}$ shown in Theorem 2. All estimated coefficients $\hat{\beta}$ are statistically significant, which indicates that CO, NO₂, O₃ and SO₂ are all significant risk factors for the air pollution index related to the CVD death rate. It reaffirms that all of the pollutants have a significant effect on the CVD death rate. The estimated coefficients for CO, NO₂, and SO₂ are negative, which is caused by the corre-

lation of these three air pollutants with O_3 . The correlation coefficients among these four air pollutants are provided in Section S5 of the supplementary files. We also study the time-varying effect of each individual pollutant on the CVD death rate by fitting a simple functional linear regression $E(Y) = \int_0^1 \eta(t)X(t)dt$ to the air pollution data, where the response variable Y is the annual CVD death rate, and the functional covariate $X(t)$ is the daily concentration of the air pollutant CO, NO_2 , SO_2 , and O_3 , respectively. Figure S4 in the supplementary file displays the estimated functional coefficient $\hat{\eta}(t)$ with the 95% pointwise confidence interval. It shows that no significant effect of each individual pollutant is found on the CVD death rate. This is another motivation for us to estimate a comprehensive air pollution index to measure the contributions of air pollutants simultaneously.

For comparison, we implemented the stacking approach to estimate the functional linear model (4.12). Figure 5 compares the estimated $\hat{\eta}_k(t)$ for the stacking functional linear model (4.12) and the estimated $\hat{\beta}_k \hat{\alpha}(t)$ for our functional index model (1.1), where $k = 1, \dots, 4$. While there is slight disagreement between the two sets of estimations from the two models, it is clear that the unstructured model has very large variability and can hardly deliver any statistically significant results.

We further assessed the prediction performance of our proposed method in

comparison with three other methods, including the stacking functional linear model (4.12), the functional additive model (Müller and Yao, 2008), and a single index model where each covariate is simply the yearly average of each pollutant. The evaluation is conducted through a ten-fold cross-validation. Table 5 displays the mean squared prediction errors (MSPE) of our proposed method and the three comparison methods. It shows that our proposed functional single index model has the smallest MSPE among all four methods. For instance, MSPE is decreased by 31% when using our proposed functional single index in comparison with the stacking functional linear model (4.12).

6. Discussion

We proposed a functional single index model to study the relation between the pollutants and CVD death rate. The model contains a single index which summarizes the pollution severity level and a time-varying coefficient which captures the seasonality of the pollution effects. Furthermore, the model is robust against the misspecification of the conditional density function $f_{Y|\mathbf{X}(t)}(\cdot)$. When replacing the function $\alpha(\cdot)$ by its B-spline approximation, the model reduces to a dimension folding model, and our estimator yields a new estimator as a by-product. This new estimator requires much more relaxed conditions on the covariates while at the same time performs much better than all existing meth-

ods. Finally, the model and method can be used in the high dimensional settings thanks to the fact that the numbers of covariate functions and spline basis are added. In contrast, the traditional functional single index described in (4.12) would result in multiplication of these two numbers.

In our analysis, we assume the functional covariate $X_i(t)$ is known to simplify the problem. However, in practice, the measurements for the functional covariate $X_i(t)$ may contain errors. To take into account the errors, model (1.1) should be further augmented. The resulting model falls into the measurement error framework and deserves careful investigation in future work.

Supplementary Materials

The supplementary document online includes the comprehensive proofs of all theoretic results. The computing codes for our simulation studies and application can be downloaded from <https://github.com///sbaek306/FSIM>.

References

- Atkinson, K. E. (1989). *An Introduction to Numerical Analysis (2nd ed.)*. New York: John Wiley & Sons.
- Bickel, P. J., C. A. Klaassen, P. J. Bickel, Y. Ritov, J. Klaassen, J. A. Wellner, and Y. Ritov (1993). *Efficient and adaptive estimation for semiparametric models*. New York: Springer.
- Chen, D., P. Hall, H.-G. Müller, et al. (2011). Single and multiple index functional regression models with nonparametric link. *The Annals of Statistics* 39(3), 1720–1747.

- Cook, D. (1998). *Regression Graphics: Ideas for Studying Regressions through Graphics*. New York: Wiley.
- Cook, R. and S. Weisberg (1991). Discussion of “sliced inverse regression for dimension reduction”. *Journal of the American Statistical Association* 86, 28–33.
- Cox, L. A. T. and D. A. Popken (2015). Has reducing fine particulate matter and ozone caused reduced mortality rates in the united states? *Annals of epidemiology* 25(3), 162–173.
- Ferré, L. and A. Yao (2003). Functional sliced inverse regression analysis. *Statistics* 37(6), 475–488.
- Ferré, L. and A. Yao (2005). Smoothed functional inverse regression. *Statistica Sinica* 15(3), 665–683.
- Ferré, L. and A. Yao (2007). Reply to the paper ‘a note on smoothed functional inverse regression’ by liliana forzani and r. dennis cook. *Statistica Sinica* 17(4), 1534–1544.
- Li, B., M. K. Kim, and N. Altman (2010). On dimension folding of matrix or array valued statistical objects. *Annals of Statistics* 38, 1094–1121.
- Li, B. and S. Wang (2007). On directional regression for dimension reduction. *Journal of the American Statistical Association* 102, 997–1008.
- Li, K. C. (1991). Sliced inverse regression for dimension reduction (with discussion). *Journal of the American Statistical Association* 86, 316–342.
- Ma, S. (2016). Estimation and inference in functional single-index models. *Annals of the Institute of Statistical Mathematics* 68(1), 181–208.
- Ma, Y. and L. Zhu (2012). A semiparametric approach to dimension reduction. *Journal of the American*

Statistical Association 107, 168–179.

Ma, Y. and L. Zhu (2013). Efficient estimation in sufficient dimension reduction. *Annals of Statistics* 41, 250–268.

Mack, Y. and B. W. Silverman (1982). Weak and strong uniform consistency of kernel regression estimates. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* 61(3), 405–415.

Müller, H. and F. Yao (2008). Functional additive models. *Journal of the American Statistical Association* 103(4), 1683.

Peng, R. and L. Welty (2004). The nmmapsdata package. *R News* 4, 10–14.

Qu, S., J.-L. Wang, X. Wang, et al. (2016). Optimal estimation for the functional cox model. *The Annals of Statistics* 44(4), 1708–1738.

Ramsay, J. and B. W. Silverman (2005). *Functional data analysis* (Second ed.). New York: Springer-Verlag.

Samet, J., F. Dominici, S. Zeger, J. Schwartz, and D. Dockery (2000). The national morbidity, mortality, and air pollution study. part i: Methods and methodologic issues. *Research report (Health Effects Institute)* (94 Pt 1), 5–14.

Tsiatis, A. A. (2004). *Semiparametric Theory and Missing Data*. New York: Springer.

Turner, M. C., M. Jerrett, C. A. Pope III, D. Krewski, S. M. Gapstur, W. R. Diver, B. S. Beckerman, J. D. Marshall, J. Su, D. L. Crouse, et al. (2016). Long-term ozone exposure and mortality in a large prospective study. *American journal of respiratory and critical care medicine* 193(10), 1134–1142.

F. Jiang

Department of Statistics and Actuarial Science,

University of Hong Kong, Hong Kong, China. E-mail: feijiang@hku.hk

Seungchul Baek

Department of Statistics, University of South Carolina,

Columbia, SC 29208, USA. E-mail: sbaek@stat.sc.edu

Jiguo Cao

Department of Statistics and Actuarial Science,

Simon Fraser University, Burnaby, BC V5A 1S6, Canada. E-mail: jiguo_cao@sfu.ca

Yanyuan Ma

Department of Statistics, Penn State University,

University Park, PA 16802, USA. E-mail: yzm63@psu.edu




Figure 1: The mean and point wise 90% confidence bands of the estimated $\hat{\alpha}(t)$ for the functional single index model (1.1) in Simulations 1 (left), 2 (middle) and 3 (right). The true $\alpha_0(t)$ is plotted in the solid curve.

Table 1: The average (AVE), the sample standard deviations (STD), the average of the estimated standard deviation (\widehat{STD}), the square root of the mean squared error (MSE) and the coverage of the estimated 95% confidence interval (CI) from the oracle (Ora), efficient (Eff) and Locally efficient (Loc) estimates of β, γ in Simulation 1.

		β_1	β_2	β_3	β_4	β_5	β_6	β_7	β_8
		1	1.2	1.5	0.5	-0.5	-1.5	-1.2	-1
Ora	AVE	1.0089	1.2070	1.5134	0.5067	-0.5033	-1.5081	-1.2141	-1.0062
	STD	0.1294	0.1364	0.1588	0.1020	0.1007	0.1553	0.1412	0.1304
	\widehat{STD}	0.1244	0.1365	0.1577	0.0994	0.0996	0.1570	0.1373	0.1242
	MSE	0.0168	0.0186	0.0254	0.0104	0.0101	0.0242	0.0201	0.0170
	CI	0.9470	0.9520	0.9540	0.9420	0.9470	0.9530	0.9420	0.9330
Eff	AVE	1.0264	1.2279	1.5398	0.5160	-0.5122	-1.5344	-1.2347	-1.0240
	STD	0.1368	0.1460	0.1719	0.1065	0.1056	0.1669	0.1522	0.1389
	\widehat{STD}	0.1349	0.1495	0.1744	0.1051	0.1052	0.1734	0.1503	0.1349
	MSE	0.0194	0.0221	0.0311	0.0116	0.0113	0.0290	0.0243	0.0198
	CI	0.9640	0.9650	0.9660	0.9500	0.9510	0.9630	0.9580	0.9520
Loc	AVE	1.0335	1.2427	1.5533	0.5194	-0.5139	-1.5479	-1.2444	-1.0341
	STD	0.1535	0.1674	0.1962	0.1229	0.1192	0.1903	0.1728	0.1559
	\widehat{STD}	0.1544	0.1700	0.1997	0.1208	0.1207	0.1969	0.1702	0.1538
	MSE	0.0247	0.0298	0.0413	0.0155	0.0144	0.0385	0.0318	0.0254
	CI	0.9550	0.9540	0.9580	0.9520	0.9570	0.9640	0.9550	0.9600

Table 2: The average (AVE), the sample standard deviations (STD), the average of the estimated standard deviation (\widehat{STD}), the square root of the mean squared error (MSE) and the coverage of the estimated 95% confidence interval (CI) from the oracle (Ora), efficient (Eff) and Locally efficient (Loc) estimates of β, γ in Simulation 2.

		β_1	β_2	β_3	β_4	β_5	β_6	β_7	β_8
		1	1.2	1.5	0.5	-0.5	-1.5	-1.2	-1
Ora	AVE	1.0037	1.2019	1.5031	0.5015	-0.5030	-1.5015	-1.2001	-1.0014
	STD	0.0611	0.0677	0.0752	0.0503	0.0493	0.0750	0.0682	0.0568
	\widehat{STD}	0.0603	0.0662	0.0764	0.0484	0.0485	0.0763	0.0662	0.0604
	MSE	0.0037	0.0046	0.0057	0.0025	0.0024	0.0056	0.0046	0.0032
	CI	0.9500	0.9430	0.9430	0.9370	0.9480	0.9550	0.9310	0.9580
Eff	AVE	1.0027	1.2018	1.5042	0.5024	-0.5037	-1.5005	-1.2004	-1.0011
	STD	0.0752	0.0835	0.0930	0.0571	0.0564	0.0966	0.0801	0.0718
	\widehat{STD}	0.0734	0.0805	0.0931	0.0579	0.0578	0.0937	0.0805	0.0730
	MSE	0.0057	0.0070	0.0087	0.0033	0.0032	0.0093	0.0064	0.0051
	CI	0.9460	0.9460	0.9470	0.9470	0.9500	0.9410	0.9470	0.9500
Loc	AVE	0.9944	1.1907	1.4877	0.4963	-0.4973	-1.4868	-1.1889	-0.9913
	STD	0.1494	0.1720	0.2098	0.0830	0.0842	0.2126	0.1736	0.1426
	\widehat{STD}	0.1571	0.1855	0.2282	0.0893	0.0895	0.2296	0.1854	0.1568
	MSE	0.0223	0.0296	0.0441	0.0069	0.0071	0.0453	0.0302	0.0204
	CI	0.9440	0.9400	0.9440	0.9530	0.9470	0.9460	0.9430	0.9460

Table 3: The average (AVE), the sample standard deviations (STD), the average of the estimated standard deviation (\widehat{STD}), the square root of the mean squared error (MSE) and the coverage of the estimated 95% confidence interval (CI) from the oracle, efficient, and locally efficient estimates of β, γ in Simulation 3.

		β_1	β_2	β_3
TRUE		-0.2	-1.0	-1.5
Oracle	AVE	-0.2009	-1.0015	-1.5005
	STD	0.0497	0.0650	0.0842
	\widehat{STD}	0.0493	0.0634	0.0860
	MSE	0.0025	0.0042	0.0071
	CI	0.9520	0.9480	0.9520
Efficient	AVE	-0.2017	-1.0057	-1.5058
	STD	0.0502	0.0662	0.0851
	\widehat{STD}	0.0497	0.0642	0.0871
	MSE	0.0025	0.0044	0.0071
	CI	0.9480	0.9440	0.9540
Locally Efficient	AVE	-0.2020	-0.9893	-1.4849
Efficient	STD	0.0769	0.1002	0.1246
	\widehat{STD}	0.0790	0.1069	0.1508
	MSE	0.0059	0.0101	0.0157
	CI	0.9630	0.9420	0.9530

Table 4: The estimated coefficients for the air pollutants CO, NO₂, SO₂ and standard errors for the functional single index model (1.1) in the air pollution data using the efficient method. The coefficient for O₃ is fixed to be 1 for identifiability, as introduced in Section 2.1.

	$\hat{\beta}_1$ (CO)	$\hat{\beta}_2$ (NO ₂)	$\hat{\beta}_3$ (SO ₂)	β_4 (O ₃)
Coefficients	-0.286	-0.971	-1.833	1.000
Standard Errors	0.080	0.006	0.002	-
<i>p</i> -values	3e-4	<5e-5	<5e-5	-

Table 5: The mean squared prediction errors of the four methods for the CVD death rate.

Methods	Mean Squared Prediction Errors
Functional single index Model (1)	2.14×10^{-6}
Stacking functional linear model (12)	3.11×10^{-6}
Functional additive model	2.56×10^{-6}
Single index model	2.44×10^{-6}




Figure 2: The estimated $\beta_k\alpha(t)$, $k = 1, \dots, 9$, and their point-wise 90% confidence bands for the proposed functional single index model (1.1), in comparison with the estimated $\hat{\eta}_k(t)$ for the simple stacking functional linear model (4.12) in Simulation 1.




Figure 3: The estimated $\hat{\alpha}(t)$ for the functional single index model (1.1) from the air pollution data. It captures the time-varying effect of the air pollution index on the annual CVD death rate. The pointwise 90% confidence band of the estimated $\hat{\alpha}(t)$ is also provided.




Figure 4: The pollution indices for Boston, New York and Chicago. The CVD death rates are shown in the legend.




Figure 5: Comparison of the estimated $\hat{\beta}_k \hat{\alpha}(t)$ in the proposed functional single index model (1.1) with their 90% confidence bands, and the estimated $\hat{\eta}_k(t)$ with their 90% confidence bands in the simple stacking functional linear model (4.12), for $k = 1, \dots, 4$. The top left panel is $\hat{\beta}_1 \hat{\alpha}(t)$ and $\hat{\eta}_1(t)$, the top right panel is $\hat{\beta}_2 \hat{\alpha}(t)$ and $\hat{\eta}_2(t)$, the bottom left panel is $\hat{\beta}_3 \hat{\alpha}(t)$ and $\hat{\eta}_3(t)$, and the bottom right panel is $\hat{\beta}_4 \hat{\alpha}(t)$ and $\hat{\eta}_4(t)$.