

NONPARAMETRIC REGRESSION ANALYSIS OF MULTIVARIATE LONGITUDINAL DATA

Dongdong Xiang¹, Peihua Qiu² and Xiaolong Pu¹

¹*East China Normal University* and ²*University of Minnesota*

Abstract: Multivariate longitudinal data are common in medical, industrial, and social science research. However, statistical analysis of such data in the current literature is restricted to linear or parametric modeling, which may well be inappropriate in applications. On the other hand, all existing nonparametric methods for analyzing longitudinal data are for univariate cases only. When longitudinal data are multivariate, nonparametric modeling becomes challenging, as one needs to properly handle the association among the observed data across different time points and across different components of the multivariate response. Motivated by data from the National Heart Lung and Blood Institute, this paper proposes a nonparametric modeling approach for analyzing multivariate longitudinal data. Our method is based on multivariate local polynomial smoothing. Both theoretical and numerical results show that it is useful in various settings.

Key words and phrases: Cluster data, local polynomial regression, longitudinal data, multivariate regression.

1. Introduction

Some nonparametric methods have been proposed in the literature for the analysis of longitudinal data. Most of them restrict their attention to the analysis of a single outcome variable measured repeatedly over time. However, experiments in medical, industrial, and social science research are often complex, characterized by several outcomes measured repeatedly over time. This paper focuses on statistical modeling of multivariate longitudinal data that are obtained from such experiments.

The example that motivates our research is the SHARe Framingham Heart Study of the National Heart Lung and Blood Institute (cf., Cupples et al. (2007)), in which 1,826 participants were followed seven times each at different ages. Multiple medical indices that are important risk factors of stroke, including systolic blood pressure (mmHg), diastolic blood pressure (mmHg), total cholesterol level (mg/100ml), and glucose level (mg/100ml), were measured at each time for each participant, and it was of interest to the medical researchers to know how these indices change over time. Similar studies have been reported in

the literature. See, for instance, Godleski et al. (2000), Roy and Lin (2000), and Fieuws and Verbeke (2006).

There is some existing research about the statistical analysis of multivariate longitudinal data. However, almost all of it assumes that the mean response follows a parametric model (cf., Gray and Brookmeyer (2000); O'Brien and Fitzmaurice (2004)) or that the error term has a given parametric distribution (cf., Coull and Staudenmayer (2004); Fieuws and Verbeke (2006); Roy and Lin (2000)). When the model assumptions are valid, these methods should be effective. But, in practice, it is difficult to obtain sufficient prior information to properly specify parametric models. There is some existing research on nonparametric or semiparametric modeling of longitudinal data, see, for instance, Liang and Zeger (1986), Lin and Carroll (2000, 2001), Wang (2003), Fitzmaurice, Laird and Ware (2004), Weiss (2005), Chen and Jin (2005), and Li (2011). The existing nonparametric or semiparametric methods are for analyzing univariate longitudinal data; we have not found existing research on nonparametric modeling for multivariate longitudinal data.

In this paper, we develop a nonparametric modeling approach for analyzing multivariate longitudinal data. In our approach, possible correlation among different components of the response is properly accommodated, along with possible correlation across different time points. The method is based on local polynomial kernel smoothing, and is described in Section 2. In Section 3, some of its theoretical properties are discussed. In section 4, the results of a simulation study are presented. Furthermore and we apply our method to the data of the SHARe Framingham Heart Study. Some concluding remarks are given in Section 5. Technical details are provided in the Appendix.

2. Proposed Method

Let $\mathbf{y}_{ij} = (y_{ij1}, y_{ij2}, \dots, y_{ijq})^T$ be the q -dimensional response observed at the j th time point t_{ij} from the i th subject, $j = 1, \dots, J$ and $i = 1, \dots, n$. we assume the multivariate nonparametric regression model

$$\mathbf{y}_{ij} = \mathbf{m}(t_{ij}) + \boldsymbol{\varepsilon}_{ij}, \quad j = 1, \dots, J, \quad i = 1, \dots, n, \quad (2.1)$$

where $\mathbf{m}(t_{ij}) = (m_1(t_{ij}), m_2(t_{ij}), \dots, m_q(t_{ij}))^T$ is the mean of \mathbf{y}_{ij} , and $\boldsymbol{\varepsilon}_{ij} = (\varepsilon_{ij1}, \dots, \varepsilon_{ijq})^T$ is the q -dimensional random error. Let

$$Y_i = (\mathbf{y}_{i1}, \dots, \mathbf{y}_{iJ})^T, \quad \boldsymbol{\varepsilon}_i = (\boldsymbol{\varepsilon}_{i1}, \dots, \boldsymbol{\varepsilon}_{iJ})^T,$$

$\text{vec}(Y_i)$ be created by connecting all columns of Y_i one after another, and $\text{vec}(\boldsymbol{\varepsilon}_i)$ be created from the columns of $\boldsymbol{\varepsilon}_i$ in the same way. Thus, Y_i and $\boldsymbol{\varepsilon}_i$ are $J \times q$

matrices, and $\text{vec}(Y_i)$ and $\text{vec}(\varepsilon_i)$ are Jq -dimensional vectors. In (2.1), we assume that, for $i = 1, \dots, n$,

$$E(\text{vec}(\varepsilon_i)|t_{i1}, \dots, t_{iJ}) = 0, \quad (2.2)$$

$$\text{Cov}(\text{vec}(\varepsilon_i)|t_{i1}, \dots, t_{iJ}) = \text{Cov}(\text{vec}(Y_i)|t_{i1}, \dots, t_{iJ}) =: V_i, \quad (2.3)$$

where V_i is the conditional covariance matrix of $\text{vec}(Y_i)$ containing $q \times q$ sub-matrices, each sub-matrix a $J \times J$ matrix. The diagonal sub-matrices give the correlation among different components of the response at individual time points for the i th subject, and the off-diagonal sub-matrices give the correlation among response vectors at different time points. The model at (2.1) is quite general in that it accommodates correlation among the observed data across different time points and across different components of the multivariate response vector.

For estimation, we employ the local polynomial kernel smoothing approach that has been used in the literature for handling cases with univariate longitudinal data (e.g., Lin and Carroll (2001), Wang (2003), Chen and Jin (2005)). With multivariate longitudinal data, it is much more complicated to use this approach while allowing correlations among different components of the response. To this end, let us first define some notation. We use $\text{diag}\{a_{jl}, j = 1, \dots, J, l = 1, \dots, q\}$ to denote a diagonal matrix with the $[j + (l - 1)J]$ th diagonal element a_{jl} . The inverse of a matrix means the Moore-Penrose generalized inverse, and t denotes an arbitrary but fixed interior point of the domain of t_{ij} . The kernel function is denoted by $K(\cdot)$, chosen to be a symmetric density function with support $[-1, 1]$. Typical choices of $K(\cdot)$ are the Epanechnikov kernel $K(u) = 0.75(1 - u^2)I(|u| \leq 1)$ and the uniform kernel $K(u) = 0.5I(|u| \leq 1)$, where $I(\cdot)$ is the indicator function. Let $K_h(u) = K(u/h)/h$, where h is a bandwidth. In our setting, we need a q -dimensional bandwidth vector \mathbf{H} to allow different degrees of smoothing in different components. Let $\mathbf{H} = (h_1, \dots, h_q)^T$,

$$K_{i\mathbf{H}} = \text{diag}\{K_{h_l}(t_{ij} - t), j = 1, \dots, J, l = 1, \dots, q\},$$

$$W_i = \left(K_{i\mathbf{H}}^{-1/2} \widehat{V}_i K_{i\mathbf{H}}^{-1/2}\right)^{-1} = K_{i\mathbf{H}}^{1/2} \left(\tilde{I}_i \widehat{V}_i \tilde{I}_i\right)^{-1} K_{i\mathbf{H}}^{1/2},$$

where \widehat{V}_i is an estimator of V_i , and

$$\begin{aligned} \tilde{I}_i &= \text{diag}\{I(K_{h_l}(t_{ij} - t) > 0), j = 1, \dots, J, l = 1, \dots, q\} \\ &= \text{diag}\{I(|t_{ij} - t| \leq h_l), j = 1, \dots, J, l = 1, \dots, q\}. \end{aligned}$$

For a positive integer p , consider the p th order local polynomial kernel smoothing procedure

$$\min_{\text{vec}(\beta) \in R^{q(p+1)}} \sum_{i=1}^n [\text{vec}(Y_i) - (I_q \otimes X_i) \text{vec}(\beta)]^T W_i [\text{vec}(Y_i) - (I_q \otimes X_i) \text{vec}(\beta)], \quad (2.4)$$

where \otimes denotes the Kronecker product, and

$$X_i = \begin{pmatrix} 1 & (t_{i1} - t) & \dots & (t_{i1} - t)^p \\ \vdots & \vdots & \ddots & \vdots \\ 1 & (t_{iJ} - t) & \dots & (t_{iJ} - t)^p \end{pmatrix}_{J \times (p+1)}, \quad \beta = \begin{pmatrix} \beta_0^{(1)} & \dots & \beta_0^{(q)} \\ \vdots & \ddots & \vdots \\ \beta_p^{(1)} & \dots & \beta_p^{(q)} \end{pmatrix}_{(p+1) \times q}.$$

In (2.4), the possible correlation among different response components has been accommodated by using $W_i = \left(K_{i\mathbf{H}}^{-1/2} \widehat{V}_i K_{i\mathbf{H}}^{-1/2} \right)^{-1}$. When we know that the q response components are independent of each other, the procedure (2.4) is equivalent to applying the univariate method of Chen and Jin (2005) to each component of the response vector.

The solution to (2.4) is

$$\widehat{\text{vec}}(\beta) = \left[\sum_{i=1}^n (I_q \otimes X_i)^T W_i (I_q \otimes X_i) \right]^{-1} \left[\sum_{i=1}^n (I_q \otimes X_i)^T W_i \text{vec}(Y_i) \right]. \quad (2.5)$$

Then, the p th order local polynomial kernel estimators of $\mathbf{m}^{(k)}(t) = (m_1^{(k)}(t), \dots, m_q^{(k)}(t))^T$ for $k = 0, \dots, p$, are

$$\widehat{\mathbf{m}}^{(k)}(t) = k! \widehat{\text{vec}}(\beta)^T (I_q \otimes \mathbf{e}_{k+1}), \quad (2.6)$$

where \mathbf{e}_{k+1} is a $(p+1)$ -dimensional vector that is 1 at the $(k+1)$ th position and 0 otherwise. When $k = 0$, (2.6) becomes

$$\widehat{\mathbf{m}}^{(0)}(t) = \widehat{\text{vec}}(\beta)^T (I_q \otimes \mathbf{e}_1),$$

the p th order local polynomial kernel estimator of $\mathbf{m}(t)$.

In (2.4), we need to provide a reasonable estimator \widehat{V}_i of the covariance matrix V_i . In practice, if there are replicated observations at each time point for each subject, then the V_i can be estimated by their sample covariance matrices. Otherwise, some assumptions on the V_i are necessary. If it is reasonable to assume that V_i are the same for all i , then the common covariance matrix can be estimated as follows. First, we use the local linear kernel smoothing procedure to estimate individual components of $\mathbf{m}(\cdot)$ separately, using the Epanechnikov kernel function and the bandwidths determined by the conventional cross-validation (CV) procedure. The estimators are denoted as $\widetilde{\mathbf{m}}(\cdot) = (\widetilde{m}_1(\cdot), \dots, \widetilde{m}_q(\cdot))$. Then, we compute the residuals

$$\widetilde{\varepsilon}_{ijl} = y_{ijl} - \widetilde{m}_l(t_{ij}), \quad i = 1, \dots, n, j = 1, \dots, J, l = 1, \dots, q.$$

The $([(l - 1)J + j], [(s - 1)J + k])$ th element of V_i can then be estimated by

$$\widehat{\text{Cov}}(\varepsilon_{ijl}, \varepsilon_{iks}) = \begin{cases} \frac{\sum_{v=1}^n \tilde{\varepsilon}_{vjl} \tilde{\varepsilon}_{vks} K\left(\frac{t_{vj} - t_{ij}}{g_l}\right) K\left(\frac{t_{vk} - t_{ik}}{g_s}\right)}{\sum_{v=1}^n K\left(\frac{t_{vj} - t_{ij}}{g_l}\right) K\left(\frac{t_{vk} - t_{ik}}{g_s}\right)}, & j \neq k \text{ or } l \neq s; \\ \frac{\sum_{v=1}^n \tilde{\varepsilon}_{vjl}^2 K\left(\frac{t_{vj} - t_{ij}}{g_l}\right)}{\sum_{v=1}^n K\left(\frac{t_{vj} - t_{ij}}{g_l}\right)}, & j = k, l = s. \end{cases} \tag{2.7}$$

where g_l is the bandwidth for the response component l , $j, k = 1, \dots, J$, and $l, s = 1, \dots, q$. In (2.7), we can still use the Epanechnikov kernel function, and the bandwidths $(g_1, \dots, g_J)^T$ can be chosen as follows. With

$$y_{ijl}^* = \tilde{\varepsilon}_{ijl}^2, \quad i = 1, \dots, n, j = 1, \dots, J, l = 1, \dots, q,$$

the mean of $\mathbf{y}_{ij}^* = (y_{ij1}^*, y_{ij2}^*, \dots, y_{ijq}^*)^T$ is a good approximation of the variance of \mathbf{y}_{ij} , denoted by $\sigma^2(t_{ij})$. Then, we can use the CV procedure to choose the bandwidths for the local linear kernel smoothing of the new data when the Epanechnikov kernel function is used. The resulting bandwidths can be used as the chosen values of $(g_1, \dots, g_J)^T$. To specify V_i , one might also use a time series model for specifying the correlation of the observed data across different time points, as mentioned by Chen and Jin (2005) for univariate cases.

In certain applications, it is possible that some response components are missing at some time points. To handle such cases, our proposed method should be modified as follows. Let δ_{ijl} be a binary variable taking the value of 0 when the observation of the l th component of $\mathbf{y}(t_{ij})$ is missing and 1 otherwise. Take

$$\Delta_i = \text{diag}\{\delta_{ijl}, j = 1, \dots, J, l = 1, \dots, q\}.$$

Then, $\widehat{\text{Cov}}(\varepsilon_{ijl}, \varepsilon_{iks})$ in (2.7) is changed to

$$\widehat{\text{Cov}}'(\varepsilon_{ijl}, \varepsilon_{iks}) = \begin{cases} \frac{\sum_{v=1}^n \tilde{\varepsilon}_{vjl} \tilde{\varepsilon}_{vks} \delta_{vjl} \delta_{vks} K\left(\frac{t_{vj} - t_{ij}}{g_l}\right) K\left(\frac{t_{vk} - t_{ik}}{g_s}\right)}{\sum_{v=1}^n \delta_{vjl} \delta_{vks} K\left(\frac{t_{vj} - t_{ij}}{g_l}\right) K\left(\frac{t_{vk} - t_{ik}}{g_s}\right)}, & j \neq k \text{ or } l \neq s; \\ \frac{\sum_{v=1}^n \tilde{\varepsilon}_{vjl}^2 \delta_{vjl} K\left(\frac{t_{vj} - t_{ij}}{g_l}\right)}{\sum_{v=1}^n \delta_{vjl} K\left(\frac{t_{vj} - t_{ij}}{g_l}\right)}, & j = k, l = s. \end{cases} \tag{2.8}$$

The resulting estimator of V_i is denoted as \widehat{V}_i' and (2.5) becomes

$$\widehat{\text{vec}}(\beta)' = \left[\sum_{i=1}^n (I_q \otimes X_i)^T \Delta_i W_i' \Delta_i (I_q \otimes X_i) \right]^{-1} \left[\sum_{i=1}^n (I_q \otimes X_i)^T \Delta_i W_i' \Delta_i \text{vec}(Y_i) \right], \tag{2.9}$$

where

$$W_i' = \left(K_{i\mathbf{H}}^{-1/2} \Delta_i \widehat{V}_i' \Delta_i K_{i\mathbf{H}}^{-1/2} \right)^{-1} = K_{i\mathbf{H}}^{1/2} \left(\tilde{I}_i \Delta_i \widehat{V}_i' \Delta_i \tilde{I}_i \right)^{-1} K_{i\mathbf{H}}^{1/2}.$$

Finally, the p th order local polynomial kernel estimators of $\mathbf{m}^{(k)}(t)$, for $k = 0, \dots, p$, can still be computed by (2.6) after $\widehat{\text{vec}(\beta)}$ is replaced by $\widehat{\text{vec}(\beta)'}^{\prime}$ in (2.9). The resulting estimators are denoted by $\widehat{\mathbf{m}}^{(k)'}(t)$.

3. Asymptotic Properties

In this section we study the theoretical properties of the proposed method. These properties require some regularity conditions on the local distribution of the design points, here described along with the necessary notation.

Let Ω_v , for $1 \leq v \leq 2^J - 1$, be the $2^J - 1$ distinct non-empty subsets of $\{1, \dots, J\}$, and $B(t, \delta)$ be the interval $[t - \delta, t + \delta]$. Assume that the design points $(t_{i1}, \dots, t_{iJ})^T$, for $i = 1, \dots, n$, are independent and identically distributed, and that their partial density at any given point t in the design space exists. Partial density at t , according to Chen and Jin (2005), exists if there is a constant $\delta_0 > 0$ such that, for all $u \in B(t, \delta_0)$ and all $v = 1, \dots, 2^J - 1$,

$$\begin{aligned} & \Pr\left\{t_{1j} \in B(u, \delta), \text{ and elements in } \{t_{1j}, j \in \Omega_v\} \text{ are equal, and } t_{1j_1} \neq t_{1j} \right. \\ & \quad \left. \text{if } j_1 \notin \Omega_v \text{ and } j \in \Omega_v\right\} \\ &= \int_{-\delta}^{\delta} f_v(z + u) dz \\ &= \Pr\{t_{1j} \in B(u, \delta) \text{ for all } j \in \Omega_v, \text{ and } t_{1j} \notin B(u, \delta) \text{ for all } j \notin \Omega_v\} + o(\delta) \end{aligned}$$

for all $0 < \delta < 2\delta_0$. Here $f_v(\cdot)$, for $1 \leq v \leq 2^J - 1$, are nonnegative continuous functions on $B(t_0, 2\delta_0)$ such that $\sum_{v=1}^{2^J-1} f_v(z) > 0$ for all $z \in B(t, 2\delta_0)$. The condition ensures that the chance that two design points take values in a small neighborhood of t is negligible unless they belong to the same Ω_v .

Let $\mathcal{S}_v(0) = \{t_{1j} = t \text{ for all } j \in \Omega_v, \text{ and } t_{1j} \neq t \text{ for all } j \notin \Omega_v\}$, and take

$$\xi_v^{(sk)} = E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\}, \quad \text{for } s, k = 1, \dots, q,$$

where $\tilde{I}_{v0} = I_q \otimes \text{diag}\{I(1 \in \Omega_v), \dots, I(J \in \Omega_v)\}$ is a $qJ \times qJ$ nonrandom matrix, $\tilde{\mathbf{e}}_k$ is a q -dimensional vector with 1 at the k th position and 0 otherwise, and $\mathbf{1}_0$ is a J -dimensional vector with all components equal to 1. We further take

$$\begin{aligned} V_0(t) &= \text{Cov}(\text{vec}(\varepsilon_1) | t_{11} = t, \dots, t_{1J} = t), \\ \tilde{\xi}_v^{(sk)}(t) &= E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1} V_0(t) (\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\}, \\ \bar{\xi}_{v, l_1 l_2}^{(sk)}(t) &= E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1} (E_{l_1} \otimes I_J) V_0(t) (E_{l_2} \otimes I_J) (\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1} \\ & \quad \times (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\}, \end{aligned}$$

where E_l is a $q \times q$ matrix with 1 at the l th diagonal position and 0 otherwise, $l = 1, \dots, q$. Set $h_{\max} = \max\{h_1, \dots, h_q\}$, and let $h_l = c_l h_{\max}$, where $0 < c_l \leq 1$

are constants, $l = 1, \dots, q$. Let

$$\begin{aligned} \mu_j(h_s, h_k) &= (h_s h_k)^{-1/2} \int z^j K^{1/2}\left(\frac{h_{\max} z}{h_s}\right) K^{1/2}\left(\frac{h_{\max} z}{h_k}\right) dz, \\ \nu_j(h_s, h_k, h_{l_1}, h_{l_2}) &= (h_s h_k h_{l_1} h_{l_2})^{-1/2} \int z^j K^{1/2}\left(\frac{h_{\max} z}{h_s}\right) \\ &\quad \times K^{1/2}\left(\frac{h_{\max} z}{h_k}\right) K^{1/2}\left(\frac{h_{\max} z}{h_{l_1}}\right) K^{1/2}\left(\frac{h_{\max} z}{h_{l_2}}\right) dz, \\ \nu_{m+l,v}^{(sk)}(t) &= \sum_{l_1, l_2=1}^q \xi_{v, l_1 l_2}^{(sk)}(t) \nu_{m+l}(h_s, h_k, h_{l_1}, h_{l_2}). \end{aligned}$$

Then it can be checked that $\mu_j(h_s, h_k) = O(h_{\max}^{-1})$ and $\nu_j(h_s, h_k, h_{l_1}, h_{l_2}) = O(h_{\max}^{-2})$ for any $s, k, l_1, l_2 \in \{1, \dots, q\}$. Let S and \bar{S} be $[q(p+1)] \times [q(p+1)]$ matrices with the $[(p+1)(s-1) + m + 1, (p+1)(k-1) + l + 1]$ th elements equal to $\sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(sk)} \mu_{m+l}(h_s, h_k)$ and $\sum_{v=1}^{2^J-1} f_v(t) \nu_{m+l,v}^{(sk)}(t)$, respectively, for $s, k, m, l \in \{1, \dots, q\}$.

Proposition 1. Denote \mathcal{F}_n as the σ -algebra generated by (t_{i1}, \dots, t_{iJ}) , $i = 1, \dots, n$. Let the design points $(t_{i1}, \dots, t_{iJ})^T$, $i = 1, \dots, n$, be independent and identity distributed and assume their partial density exists at any given point t in the design space. Suppose the V_i at (2.3) are continuous functions of (t_{i1}, \dots, t_{iJ}) and the components of the $(p+1)$ th derivative $\mathbf{m}^{(p+1)}(t)$ of $\mathbf{m}(t)$ are continuous functions of t , $i = 1, \dots, n$. If it is further assumed that $h_l = c_l h_{\max}$, where $0 < c_l \leq 1$ are constants, for $l = 1, \dots, q$, $h_{\max} = o(1)$, and $1/(nh_{\max}) = o(1)$, then the following hold.

(i) The conditional covariance of $\widehat{\mathbf{m}}^{(k)}(t)$ is

$$\text{Cov}\{\widehat{\mathbf{m}}^{(k)}(t) | \mathcal{F}_n\} = \frac{k!^2}{nh_{\max}^{1+2k}} [(I_q \otimes \mathbf{e}_{k+1}^T) S^{-1} \bar{S} S^{-1} (I_q \otimes \mathbf{e}_{k+1})] + o_P\left(\frac{1}{nh_{\max}^{1+2k}}\right). \tag{3.1}$$

(ii) The conditional bias of $\widehat{\mathbf{m}}^{(k)}(t)$ is

$$\text{Bias}\{\widehat{\mathbf{m}}^{(k)}(t) | \mathcal{F}_n\} = \frac{k!}{(p+1)!} h_{\max}^{p+1-k} [(I_q \otimes \mathbf{e}_{k+1}^T) S^{-1} \mathbf{D}] + o_P(h_{\max}^{p+1-k}), \tag{3.2}$$

where $\mathbf{D} = (d_{10}, \dots, d_{1p}, \dots, d_{q0}, \dots, d_{qp})^T$ and

$$d_{sk} = \sum_{v=1}^{2^J-1} \sum_{l=1}^q f_v(t) m_l^{(p+1)}(t) \xi_v^{(sl)} \mu_{k+p+1}(h_s, h_l), \text{ for } s=1, \dots, q, k=0, \dots, p.$$

The convergence rates of the conditional covariance and the conditional bias of $\widehat{\mathbf{m}}^{(k)}(t)$ here are the same as those in univariate cases (cf., Chen and Jin

(2005)). Our results are derived in a quite general setting while in some special cases, they have simpler expressions. For instance, in cases when the components of $\mathbf{m}(\cdot)$ have similar smoothness, we can use a bandwidth vector with $h_1 \sim \dots \sim h_q \sim h_{\max}$. In such cases, $\mu_j(h_s, h_k) \approx (1/h_{\max}) \int u^j K(u) du =: (1/h_{\max})\mu_j$, and $\nu_j(h_s, h_k, h_{l_1}, h_{l_2}) \approx (1/h_{\max}^2) \int u^j K^2(u) du =: (1/h_{\max}^2)\nu_j$, where “ \approx ” means that some higher order terms have been omitted in the related expressions. Then, take $\mathbf{c}_p = (\mu_{p+1}, \dots, \mu_{2p+1})^T$, $S_1 = (\mu_{i+j})_{0 \leq i, j \leq p}$, $\bar{S}_1 = (\nu_{i+j})_{0 \leq i, j \leq p}$, and let

$$\mathbf{C} = \text{diag} \left\{ \sum_{v=1}^{2^J-1} \sum_{l=1}^q f_v(t) m_l^{(p+1)}(t) \xi_v^{(1l)}, \dots, \sum_{v=1}^{2^J-1} \sum_{l=1}^q f_v(t) m_l^{(p+1)}(t) \xi_v^{(ql)} \right\},$$

$$N = \left(\sum_{v=1}^{2^J-1} f_v(t) \bar{\xi}_v^{(sk)}(t) \right)_{q \times q} \quad \text{and} \quad M = \left(\sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(sk)} \right)_{q \times q},$$

where $\bar{\xi}_v^{(sk)}(t) = \sum_{l_1, l_2=1}^q \bar{\xi}_{v, l_1 l_2}^{(sk)}(t)$. In such cases, the results in Proposition 1 can be simplified.

Corollary 1. *Under conditions in Proposition 1, if $h_1 \sim \dots \sim h_q$. the following hold.*

(i) *The conditional covariance of $\widehat{\mathbf{m}}^{(k)}(t)$ is*

$$\text{Cov}\{\widehat{\mathbf{m}}^{(k)}(t) | \mathcal{F}_n\} = \frac{k!^2}{nh_{\max}^{1+2k}} \mathbf{e}_{k+1}^T S_1^{-1} \bar{S}_1 S_1^{-1} \mathbf{e}_{k+1} M^{-1} N M^{-1} + o_P \left(\frac{1}{nh_{\max}^{1+2k}} \right). \tag{3.3}$$

(ii) *The conditional bias of $\widehat{\mathbf{m}}^{(k)}(t)$ is*

$$\text{Bias}\{\widehat{\mathbf{m}}^{(k)}(t) | \mathcal{F}_n\} = \frac{k! h_{\max}^{p+1-k}}{(p+1)!} \mathbf{e}_{k+1}^T S_1^{-1} \mathbf{c}_p M^{-1} \mathbf{C} + o_P(h_{\max}^{p+1-k}). \tag{3.4}$$

When $p - k$ is even, the first term on the right side of the above expression is 0.

Compared to (3.1) and (3.2), the leading terms of (3.3) and (3.4) are much simpler. For practical purpose, we can use $h_1 = \dots = h_q = h$ for simplicity. Another special case that deserves attention has different response components independent of each other. Then the matrices \widehat{V}_i are nearly block diagonal and our method is similar to the one that handles individual response components separately.

Corollary 2. *Under conditions in Proposition 1, suppose that different response components are independent of each other. Then, the following hold.*

(i) The conditional covariance of $\widehat{\mathbf{m}}^{(k)}(t)$ is

$$\begin{aligned} \text{Cov}\{\widehat{\mathbf{m}}^{(k)}(t)|\mathcal{F}_n\} &= \frac{k!^2}{n} \mathbf{e}_{k+1}^T S_1^{-1} \bar{S}_1 S_1^{-1} \mathbf{e}_{k+1} \\ &\quad \times \text{diag} \left\{ \frac{\sum_{v=1}^{2^J-1} f_v(t) \tilde{\xi}_v^{(ll)}(t)}{\{\sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(ll)}\}^2} h_l^{1+2k}, l = 1, \dots, q \right\} \\ &\quad + o_P \left(\frac{1}{nh_{\max}^{1+2k}} \right). \end{aligned} \tag{3.5}$$

(ii) The conditional bias of $\widehat{\mathbf{m}}^{(k)}(t)$ is

$$\begin{aligned} \text{Bias}\{\widehat{\mathbf{m}}^{(k)}(t)|\mathcal{F}_n\} &= \frac{k!}{(p+1)!} \mathbf{e}_{k+1}^T S_1^{-1} \mathbf{c}_p \\ &\quad \times \text{diag} \left\{ m_l^{(p+1)}(t) h_l^{p+1-k}, l = 1, \dots, q \right\} \\ &\quad + o_P(h_{\max}^{p+1-k}). \end{aligned} \tag{3.6}$$

In cases when $p - k$ is even, the first term on the right side of the above expression is 0.

We next discuss the properties of our proposed method when there are missing observations. First, we introduce some notation. Let $\xi_v^{(sk)}$ denote $\xi_v^{(sk)}$ after $(\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1}$ is replaced by $(\tilde{I}_{v0} \Delta_1 \widehat{V}_1 \Delta_1 \tilde{I}_{v0})^{-1}$ in its definition, and $\bar{\xi}_{v,l_1 l_2}^{(sk)}(t)$ denote $\bar{\xi}_{v,l_1 l_2}^{(sk)}(t)$ after $(\tilde{I}_{v0} \widehat{V}_1 \tilde{I}_{v0})^{-1}$ and $\widehat{V}_0(t)$ are replaced by $(\tilde{I}_{v0} \Delta_1 \widehat{V}_1 \Delta_1 \tilde{I}_{v0})^{-1}$ and $\Delta_1 \widehat{V}_0(t) \Delta_1$, respectively. Furthermore, let S' and \bar{S}' be $[q(p+1)] \times [q(p+1)]$ matrices with their $[(p+1)(s-1) + m + 1, (p+1)(k-1) + l + 1]$ th elements $\sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(sk)} \mu_{m+l}(h_s, h_k)$ and $\sum_{v=1}^{2^J-1} f_v(t) \nu_{m+l,v}^{(sk)}(t)$, respectively, for $s, k, m, l \in \{1, \dots, q\}$, where

$$\nu_{m+l,v}^{(sk)}(t) = \sum_{l_1, l_2=1}^q \bar{\xi}_{v, l_1 l_2}^{(sk)}(t) \nu_{m+l}^{\prime}(h_s, h_k, h_{l_1}, h_{l_2}).$$

Corollary 3. Under the assumptions in Proposition 1 and that $P(\delta_{ijl} = 0) = p_l$, $i = 1, \dots, n$, $j = 1, \dots, J$, and $l = 1, \dots, q$, where $p_l \in [0, 1)$ does not depend on i and j . Then, the following hold.

(i) The conditional covariance of $\widehat{\mathbf{m}}^{(k)'}(t)$ is

$$\text{Cov}\{\widehat{\mathbf{m}}^{(k)'}(t)|\mathcal{F}_n\} \tag{3.7}$$

$$= \frac{k!^2}{nh_{\max}^{1+2k}} [(P^{-1} \otimes \mathbf{e}_{k+1}^T) S'^{-1} \bar{S}' S'^{-1} (P^{-1} \otimes \mathbf{e}_{k+1})] + o_P \left(\frac{1}{nh_{\max}^{1+2k}} \right), \tag{3.8}$$

where $P = \text{diag}\{p_1, \dots, p_q\}$.

(ii) The conditional bias of $\widehat{\mathbf{m}}^{(k)'}(t)$ is

$$\text{Bias}\{\widehat{\mathbf{m}}^{(k)'}(t)|\mathcal{F}_n\} = \frac{k!}{(p+1)!} h_{\max}^{p+1-k} [(P^{-1} \otimes \mathbf{e}_{k+1}^T) S'^{-1} \mathbf{D}'] + o_P(h_{\max}^{p+1-k}),$$

where $\mathbf{D}' = (d'_{10}, \dots, d'_{1p}, \dots, d'_{q0}, \dots, d'_{qp})^T$ and

$$d'_{sk} = \sum_{v=1}^{2^J-1} \sum_{l=1}^q f_v(t) m_l^{(p+1)}(t) \xi_v^{l(s)} \mu_{k+p+1}(h_s, h_l), \text{ for } s=1, \dots, q, k=0, \dots, p.$$

4. Numerical Study

In this section, we investigate the numerical performance of the proposed method using simulation and a data example. We also discuss estimation of the covariance matrices V_i defined at (2.3) and the selection of the bandwidth vector used in the local smoothing estimators.

We first considered observed data without missing values. The simulated data were generated from the model (2.1) with $J = 3$, $q = 3$, and

$$m_1(t) = 2 \times \exp\{\sin(10t)\}, \quad m_2(t) = 1 - \exp\{-t\}, \quad m_3(x) = 1 - \exp\{-t\} + 2 \sin(10t).$$

The error term $\text{vec}(\varepsilon_i)$ was normal with mean $\mathbf{0}$. Its correlation matrix was specified as follows: for $j, k, l, s = 1, 2, 3$,

$$\text{corr}(\varepsilon_{1jl}, \varepsilon_{1ks}) = \begin{cases} 1, & \text{if } j = k, l = s, \\ \rho_1, & \text{if } j \neq k, l = s, \\ \rho_2, & \text{if } j = k, l \neq s, \\ \rho_1 \rho_2, & \text{if } j \neq k, l \neq s, \end{cases}$$

$$\text{var}(\varepsilon_{111}) = \frac{\text{var}(\varepsilon_{112})}{2} = \frac{\text{var}(\varepsilon_{113})}{3} = 0.25,$$

$$\text{var}(\varepsilon_{121}) = \frac{\text{var}(\varepsilon_{122})}{2} = \frac{\text{var}(\varepsilon_{123})}{3} = 0.64,$$

$$\text{var}(\varepsilon_{131}) = \frac{\text{var}(\varepsilon_{132})}{2} = \frac{\text{var}(\varepsilon_{133})}{3} = 0.36.$$

The design points $\{t_{ij}, j = 1, 2, 3, i = 1, \dots, n\}$ were uniform $U[-2, 2]$, and independent of the errors.

In this, ρ_1 specifies the association of individual response components over different time points, and ρ_2 specifies the association among different response components at a given time point. We considered cases I: $\rho_1 = 0.8$ and $\rho_2 = 0$; II: $\rho_1 = 0$ and $\rho_2 = 0.8$; III: $\rho_1 = 0.8$ and $\rho_2 = 0.8$.

With multivariate longitudinal data, besides the proposed method denoted MULTIVARIATE here, there are a number of alternative approaches. One is

to apply the univariate method of Chen and Jin (2005) to each dimension of the multivariate longitudinal data to obtain estimators of the individual components of $\mathbf{m}(\cdot)$, denoted as INDIVIDUAL. Another is a simplified version of MULTIVARIATE using the same bandwidth h in all dimensions, denoted as SIMPLIFIED. For each method, we computed the values of the estimator $\widehat{\mathbf{m}}(t)$ at 101 grid points $\{t_j = -1.8 + 0.036 \times j, j = 0, \dots, 100\}$, and three performance measures were computed:

$$\text{Bias}_l = \frac{1}{101} \sum_{j=0}^{100} |m_l(t_j) - \widehat{m}_l(t_j)|,$$

$$\text{SD}_l = \text{sample standard deviation of } \{m_l(t_j) - \widehat{m}_l(t_j), j = 0, 1, \dots, 100\},$$

$$\text{MISE}_l = \frac{4}{101} \sum_{j=0}^{100} (m_l(t_j) - \widehat{m}_l(t_j))^2,$$

where $l = 1, 2, 3$ is the index of the response components. To remove some randomness, the values of these measures that are given are averages computed from 100 replicated simulations.

In all methods considered, the Epanechnikov kernel function and local linear smoothing were used (cf., (2.4)). For a fair comparison, we first used the true covariance matrices V_i , instead of their estimates, in all methods. The optimal bandwidths of each method were then searched by minimizing the MISE value. The searched optimal bandwidths and the corresponding values of the performance measures in the three cases considered are presented in Tables 1 and 2, respectively, for sample sizes $n = 100$ and $n = 200$. For each measure, its values corresponding to the three response components are presented separately, together with their summation SUM.

From Table 1, it can be seen that, in case I the methods MULTIVARIATE and INDIVIDUAL perform exactly the same. As a matter of fact, it can be checked that the two methods are equivalent in such cases. Compared to SIMPLIFIED, their MISE values are smaller across all three response components. This means that when the curvature of the three components of $\mathbf{m}(\cdot)$ are quite different, the method SIMPLIFIED may not be appropriate to use. It also suggests that the proposed method MULTIVARIATE is appropriate even in cases when the response components are independent. In case II, we see that the method MULTIVARIATE performs better than both methods INDIVIDUAL and SIMPLIFIED in terms of the SUMs of the three performance measures, although it is slightly worse than the method INDIVIDUAL for estimating $m_2(\cdot)$. In case III, the method MULTIVARIATE also performs better than both methods INDIVIDUAL and SIMPLIFIED in terms of the SUMs of the three performance measures. Similar conclusions can be made from results in Table 2.

Table 1. Averaged performance measures Bias, SD, and MISE, based on 100 replicated simulations, of the methods MULTIVARIATE, INDIVIDUAL, and SIMPLIFIED when $n = 100$. The numbers in \mathbf{H} are the searched optimal bandwidths.

| Case | Components | MULTIVARIATE | | | INDIVIDUAL | | | SIMPLIFIED | | |
|------|------------|-------------------------------------|-------|-------|-------------------------------------|-------|-------|-------------------------------------|-------|-------|
| | | Bias | SD | MISE | Bias | SD | MISE | Bias | SD | MISE |
| I | | $\mathbf{H} = (0.08, 0.65, 0.11)^T$ | | | $\mathbf{H} = (0.08, 0.65, 0.11)^T$ | | | $\mathbf{H} = (0.11, 0.11, 0.11)^T$ | | |
| | 1 | 0.103 | 0.224 | 0.246 | 0.103 | 0.224 | 0.246 | 0.159 | 0.199 | 0.286 |
| | 2 | 0.036 | 0.114 | 0.057 | 0.036 | 0.114 | 0.057 | 0.020 | 0.269 | 0.267 |
| | 3 | 0.140 | 0.317 | 0.455 | 0.140 | 0.317 | 0.455 | 0.117 | 0.334 | 0.470 |
| | SUM | 0.279 | 0.655 | 0.758 | 0.279 | 0.655 | 0.758 | 0.295 | 0.802 | 1.023 |
| II | | $\mathbf{H} = (0.08, 0.45, 0.11)^T$ | | | $\mathbf{H} = (0.09, 0.5, 0.11)^T$ | | | $\mathbf{H} = (0.1, 0.1, 0.1)^T$ | | |
| | 1 | 0.102 | 0.186 | 0.201 | 0.139 | 0.211 | 0.280 | 0.169 | 0.196 | 0.304 |
| | 2 | 0.067 | 0.128 | 0.080 | 0.043 | 0.112 | 0.056 | 0.024 | 0.264 | 0.259 |
| | 3 | 0.119 | 0.253 | 0.297 | 0.151 | 0.307 | 0.449 | 0.127 | 0.325 | 0.459 |
| | SUM | 0.288 | 0.567 | 0.578 | 0.333 | 0.630 | 0.785 | 0.320 | 0.785 | 1.022 |
| III | | $\mathbf{H} = (0.07, 0.5, 0.11)^T$ | | | $\mathbf{H} = (0.09, 0.65, 0.12)^T$ | | | $\mathbf{H} = (0.1, 0.1, 0.1)^T$ | | |
| | 1 | 0.074 | 0.208 | 0.211 | 0.129 | 0.217 | 0.287 | 0.188 | 0.193 | 0.334 |
| | 2 | 0.067 | 0.139 | 0.091 | 0.044 | 0.112 | 0.058 | 0.018 | 0.251 | 0.232 |
| | 3 | 0.105 | 0.268 | 0.312 | 0.159 | 0.298 | 0.440 | 0.133 | 0.314 | 0.443 |
| | SUM | 0.245 | 0.615 | 0.615 | 0.332 | 0.627 | 0.785 | 0.339 | 0.758 | 1.009 |

Table 2. Averaged performance measures Bias, SD, and MISE, based on 100 replicated simulations, of the methods MULTIVARIATE, INDIVIDUAL, and SIMPLIFIED when $n = 200$. The numbers in \mathbf{H} are the searched optimal bandwidths.

| Case | Components | MULTIVARIATE | | | INDIVIDUAL | | | SIMPLIFIED | | |
|------|------------|-------------------------------------|-------|-------|-------------------------------------|-------|-------|-------------------------------------|-------|-------|
| | | Bias | SD | MISE | Bias | SD | MISE | Bias | SD | MISE |
| I | | $\mathbf{H} = (0.06, 0.5, 0.09)^T$ | | | $\mathbf{H} = (0.06, 0.5, 0.09)^T$ | | | $\mathbf{H} = (0.09, 0.09, 0.09)^T$ | | |
| | 1 | 0.064 | 0.169 | 0.127 | 0.064 | 0.169 | 0.127 | 0.137 | 0.140 | 0.171 |
| | 2 | 0.031 | 0.083 | 0.031 | 0.031 | 0.083 | 0.031 | 0.015 | 0.184 | 0.125 |
| | 3 | 0.091 | 0.227 | 0.226 | 0.091 | 0.227 | 0.226 | 0.091 | 0.227 | 0.226 |
| | SUM | 0.185 | 0.479 | 0.383 | 0.185 | 0.479 | 0.383 | 0.242 | 0.551 | 0.522 |
| II | | $\mathbf{H} = (0.06, 0.4, 0.08)^T$ | | | $\mathbf{H} = (0.06, 0.45, 0.09)^T$ | | | $\mathbf{H} = (0.08, 0.08, 0.08)^T$ | | |
| | 1 | 0.059 | 0.137 | 0.091 | 0.066 | 0.170 | 0.132 | 0.113 | 0.146 | 0.151 |
| | 2 | 0.041 | 0.091 | 0.038 | 0.035 | 0.079 | 0.030 | 0.017 | 0.198 | 0.144 |
| | 3 | 0.070 | 0.200 | 0.169 | 0.106 | 0.228 | 0.242 | 0.085 | 0.244 | 0.251 |
| | SUM | 0.170 | 0.428 | 0.298 | 0.207 | 0.477 | 0.404 | 0.214 | 0.587 | 0.546 |
| III | | $\mathbf{H} = (0.06, 0.45, 0.10)^T$ | | | $\mathbf{H} = (0.06, 0.55, 0.1)^T$ | | | $\mathbf{H} = (0.09, 0.09, 0.09)^T$ | | |
| | 1 | 0.059 | 0.136 | 0.087 | 0.061 | 0.168 | 0.123 | 0.134 | 0.137 | 0.164 |
| | 2 | 0.054 | 0.101 | 0.050 | 0.031 | 0.082 | 0.030 | 0.019 | 0.186 | 0.128 |
| | 3 | 0.093 | 0.191 | 0.171 | 0.120 | 0.221 | 0.244 | 0.099 | 0.233 | 0.243 |
| | SUM | 0.206 | 0.427 | 0.308 | 0.212 | 0.470 | 0.397 | 0.252 | 0.555 | 0.535 |

Table 3. Averaged performance measures Bias, SD, and MISE, based on 100 replicated simulations, of the methods MULTIVARIATE, INDIVIDUAL, and SIMPLIFIED when $n = 200$ and V_i are estimated. The numbers in \mathbf{H} are the searched optimal bandwidths.

| Case | Components | MULTIVARIATE | | | INDIVIDUAL | | | SIMPLIFIED | | |
|------|------------|-------------------------------------|-------|-------|------------------------------------|-------|-------|-------------------------------------|-------|-------|
| | | Bias | SD | MISE | Bias | SD | MISE | Bias | SD | MISE |
| I | | $\mathbf{H} = (0.06, 0.5, 0.09)^T$ | | | $\mathbf{H} = (0.06, 0.5, 0.09)^T$ | | | $\mathbf{H} = (0.08, 0.08, 0.08)^T$ | | |
| | 1 | 0.065 | 0.170 | 0.129 | 0.064 | 0.170 | 0.128 | 0.112 | 0.147 | 0.147 |
| | 2 | 0.032 | 0.083 | 0.032 | 0.032 | 0.082 | 0.032 | 0.016 | 0.197 | 0.143 |
| | 3 | 0.092 | 0.227 | 0.227 | 0.092 | 0.227 | 0.226 | 0.072 | 0.243 | 0.239 |
| | SUM | 0.189 | 0.480 | 0.387 | 0.189 | 0.479 | 0.386 | 0.200 | 0.587 | 0.529 |
| II | | $\mathbf{H} = (0.06, 0.4, 0.09)^T$ | | | $\mathbf{H} = (0.06, 0.4, 0.09)^T$ | | | $\mathbf{H} = (0.08, 0.08, 0.08)^T$ | | |
| | 1 | 0.066 | 0.138 | 0.096 | 0.066 | 0.171 | 0.133 | 0.114 | 0.147 | 0.153 |
| | 2 | 0.041 | 0.091 | 0.038 | 0.029 | 0.084 | 0.030 | 0.017 | 0.199 | 0.146 |
| | 3 | 0.091 | 0.193 | 0.172 | 0.107 | 0.229 | 0.244 | 0.086 | 0.246 | 0.256 |
| | SUM | 0.197 | 0.421 | 0.306 | 0.202 | 0.483 | 0.407 | 0.217 | 0.591 | 0.555 |
| III | | $\mathbf{H} = (0.06, 0.45, 0.09)^T$ | | | $\mathbf{H} = (0.06, 0.55, 0.1)^T$ | | | $\mathbf{H} = (0.09, 0.09, 0.09)^T$ | | |
| | 1 | 0.063 | 0.138 | 0.091 | 0.061 | 0.168 | 0.124 | 0.138 | 0.138 | 0.171 |
| | 2 | 0.040 | 0.095 | 0.040 | 0.033 | 0.081 | 0.031 | 0.019 | 0.187 | 0.130 |
| | 3 | 0.084 | 0.200 | 0.178 | 0.122 | 0.222 | 0.247 | 0.101 | 0.235 | 0.248 |
| | SUM | 0.187 | 0.432 | 0.309 | 0.216 | 0.470 | 0.402 | 0.258 | 0.560 | 0.549 |

Generally, the covariance matrices V_i are unknown and need to be estimated from observed data. We investigated the performance of the three methods when V_i , $i = 1, \dots, n$, are assumed the same and are estimated by the procedure (2.7). The estimated V_i were used in the three methods in place of the true matrices V_i . The corresponding results of the three methods in cases when $n = 200$, and when the bandwidths are chosen to be optimal by minimizing the MISE values, are presented in Table 3. From the table, we can see that similar conclusions can be made here to those from Tables 1 and 2, regarding the relative performance of the three methods. By comparing the results of MULTIVARIATE in Tables 2 and 3, we see that they are almost the same, which suggests that the procedure (2.7) for specifying V_i is quite efficient. Corresponding results in the case when $n = 100$ are similar and thus omitted here.

In practice, the optimal bandwidths are also unknown. To implement our method, we propose using cross-validation (CV) to determine the bandwidths, as follows. Let

$$CV_l(h_l) = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^J (y_l(t_{ij}) - \hat{m}_{l,-i}(t_{ij}))^2, \quad \text{for } l = 1, 2, 3,$$

$$CV(\mathbf{H}) = CV_1(h_1) + CV_2(h_2) + CV_3(h_3),$$

Table 4. Averaged performance measures Bias, SD, and MISE, based on 100 replicated simulations, of the method MULTIVARIATE-CV when $n = 100$ or 200.

| n | Components | Case I | | | Case II | | | Case III | | |
|-----|------------|--------|-------|-------|---------|-------|-------|----------|-------|-------|
| | | Bias | SD | MISE | Bias | SD | MISE | Bias | SD | MISE |
| 100 | 1 | 0.118 | 0.236 | 0.297 | 0.136 | 0.192 | 0.258 | 0.137 | 0.197 | 0.252 |
| | 2 | 0.062 | 0.117 | 0.075 | 0.083 | 0.127 | 0.093 | 0.084 | 0.123 | 0.094 |
| | 3 | 0.157 | 0.326 | 0.497 | 0.151 | 0.274 | 0.389 | 0.144 | 0.283 | 0.376 |
| | SUM | 0.336 | 0.679 | 0.870 | 0.371 | 0.593 | 0.740 | 0.365 | 0.604 | 0.722 |
| 200 | 1 | 0.079 | 0.165 | 0.138 | 0.090 | 0.130 | 0.118 | 0.101 | 0.125 | 0.117 |
| | 2 | 0.048 | 0.081 | 0.041 | 0.065 | 0.076 | 0.046 | 0.060 | 0.078 | 0.041 |
| | 3 | 0.109 | 0.211 | 0.227 | 0.117 | 0.179 | 0.187 | 0.121 | 0.192 | 0.211 |
| | SUM | 0.236 | 0.457 | 0.407 | 0.272 | 0.385 | 0.351 | 0.282 | 0.396 | 0.369 |

where $\widehat{m}_{l,-i}(\cdot)$ is the “leave-one-subject-out” estimator of $m_l(\cdot)$ obtained when the observations of the i th subject are not used. Then the three bandwidths can be determined by minimizing $CV(\mathbf{H})$ over R_+^3 . However, this minimization process is time-consuming. To simplify the computation, we suggest using a two-step CV procedure instead, noticing in Tables 1 and 2 that the optimal bandwidths of MULTIVARIATE and INDIVIDUAL are actually quite close to each other. In the first step, we determine the individual bandwidths $\{h_l, l = 1, 2, 3\}$ separately by applying CV to the method INDIVIDUAL. The selected bandwidths from this step are denoted as $\{h_{l,0}, l = 1, 2, 3\}$. Then, in the second step, we determine the three bandwidths by minimizing $CV(\mathbf{H})$ in a small neighborhood of $(h_{1,0}, h_{2,0}, h_{3,0})^T$. In our simulation study, we used the neighborhood $\{(h_1, h_2, h_3) | h_1 = h_{10} + 0.01\delta_1, h_2 = h_{20} + 0.05\delta_2, h_3 = h_{30} + 0.01\delta_3, \delta_1, \delta_2, \delta_3 = 0, \pm 1, \pm 2\}$. The method MULTIVARIATE with the bandwidths chosen by the CV procedure and the covariance matrix estimated by (2.7) is denoted as MULTIVARIATE-CV. Its results corresponding to the cases considered in Tables 1 and 2 are presented in Table 4. By comparing tables, we can see that MULTIVARIATE-CV performs a little worse than MULTIVARIATE, but still favorably compared to the methods INDIVIDUAL and SIMPLIFIED, in these cases when the response components are correlated, even if INDIVIDUAL and SIMPLIFIED use their optimal bandwidths.

Next, we considered an example in which missing observations were present in the observed data. The setup of this example was the same as that of Table 4, except that $n = 200$, with probabilities of missing observations for the components taken as $p_1 = p_2 = p_3 = \pi$, and $\pi = 0.05, 0.1$, or 0.2 . The results are presented in Table 5, computed by (2.6) after $\widehat{\text{vec}}(\beta)$ is replaced by $\widehat{\text{vec}}(\beta)$ in (2.9). From the table, it can be seen that the MISE value increases when π increases, intuitively reasonable, and that our method performs reasonably well in such cases.

Table 5. Averaged performance measures Bias, SD, and MISE, based on 100 replicated simulations, of MULTIVARIATE-CV when $n = 200$, the probabilities of missing observations for the three components are $p_1 = p_2 = p_3 = \pi = 0.05, 0.1$, or 0.2 .

| π | Components | Case I | | | Case II | | | Case III | | |
|-------|------------|--------|-------|-------|---------|-------|-------|----------|-------|-------|
| | | Bias | SD | MISE | Bias | SD | MISE | Bias | SD | MISE |
| 0.05 | 1 | 0.061 | 0.188 | 0.148 | 0.059 | 0.152 | 0.105 | 0.052 | 0.151 | 0.098 |
| | 2 | 0.014 | 0.123 | 0.056 | 0.034 | 0.124 | 0.061 | 0.036 | 0.131 | 0.069 |
| | 3 | 0.092 | 0.241 | 0.251 | 0.081 | 0.213 | 0.195 | 0.070 | 0.222 | 0.203 |
| | SUM | 0.167 | 0.552 | 0.455 | 0.174 | 0.489 | 0.361 | 0.158 | 0.504 | 0.370 |
| 0.1 | 1 | 0.062 | 0.193 | 0.158 | 0.058 | 0.158 | 0.114 | 0.067 | 0.151 | 0.118 |
| | 2 | 0.015 | 0.122 | 0.056 | 0.034 | 0.123 | 0.061 | 0.045 | 0.121 | 0.069 |
| | 3 | 0.094 | 0.248 | 0.265 | 0.078 | 0.220 | 0.204 | 0.079 | 0.214 | 0.213 |
| | SUM | 0.171 | 0.563 | 0.479 | 0.170 | 0.501 | 0.379 | 0.191 | 0.486 | 0.400 |
| 0.2 | 1 | 0.066 | 0.207 | 0.182 | 0.063 | 0.169 | 0.132 | 0.076 | 0.158 | 0.128 |
| | 2 | 0.014 | 0.131 | 0.063 | 0.034 | 0.138 | 0.082 | 0.049 | 0.138 | 0.091 |
| | 3 | 0.096 | 0.268 | 0.305 | 0.092 | 0.237 | 0.247 | 0.115 | 0.224 | 0.311 |
| | SUM | 0.176 | 0.606 | 0.550 | 0.189 | 0.544 | 0.461 | 0.240 | 0.520 | 0.530 |

Finally, we apply our proposed method to the SHARe Framingham Heart Study data that is described in Section 1. The raw data can be downloaded from the web page http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000007.v4.p2. After deleting patients with outlier values, a total of $n = 1028$ non-stroke patients with ages from 14 to 85 were included in our analysis. In this example, the response is 4-dimensional, $q = 4$, and each patient was followed seven times, $J = 7$. In our proposed method, we used $p = 1$ and the covariance matrices V_i were determined by the procedure (2.7). The bandwidth vector \mathbf{H} was chosen using two-step CV, and the chosen bandwidth vector was $\mathbf{H} = (9, 5, 4, 6)^T$. The four estimated components of $\mathbf{m}(\cdot)$ are shown in the four plots of Figure 1 by the solid curves. After obtaining the estimator $\widehat{\mathbf{m}}(\cdot)$, we estimated the variance functions of the components of the multivariate response $\mathbf{y}(t)$ by first computing the residuals

$$\widehat{\varepsilon}_{ijl} = y_{ijl} - \widehat{m}_l(t_{ij}), \quad i = 1, \dots, n, j = 1, \dots, J, l = 1, \dots, q.$$

The estimators of the variance functions were obtained by applying the method described in Section 2. Using the estimated variance functions, pointwise 95% confidence bands of the components of $\mathbf{m}(\cdot)$ were constructed and are shown in Figure 1 (a)–(d) by the dashed curves, along with the observed longitudinal data of the first 20 patients, shown by little circles connected by thin lines. From the plots, it can be seen that our estimators describe the observed data reasonably well.




Figure 1. Estimated mean components, the pointwise 95% confidence bands of the true mean response components, and the observed longitudinal data of the first 20 patients in the dataset of the SHARe Framingham Heart Study. (a) systolic blood pressure, (b) diastolic blood pressure, (c) cholesterol level, and (d) glucose level.

5. Concluding Remarks

We have proposed a local smoothing method for analyzing multivariate longitudinal data. That can accommodate not only the correlation among observations across different time points, but the correlation among different response components. Numerical results presented in the paper suggests that our method can perform well in applications. Although the explanatory variable t is univariate here, it is possible to generalize the method to handle multiple explanatory variables, using methods similar to these in Ruppert and Wand (1994).

There are several issues that have not been addressed. First, our numerical results show that the cross-validation procedure for choosing the bandwidths

works reasonably well. But, as pointed out by Hall and Robinson (2009), the bandwidths chosen by this approach usually have a large variability. They proposed two computationally procedures to overcome this limitation. Thus, research is needed to produce an efficient and computationally simple procedure for choosing bandwidths. Second, our method may not be suitable for high-dimensional multivariate longitudinal data because of the complexity in computing estimators of V_i and in choosing bandwidths, and research might develop appropriate methods for handling such cases. Third, in Corollary 3, it is assumed that the probabilities of missing observations of the response components are unchanged over time. In applications, this assumption may not be valid. If the probabilities of missing observations depend on observation times, variable bandwidths might be more appropriate to our method. At places with more missing observations, bandwidths should be chosen larger and this not trivial topic is left for our future research.

Acknowledgement

We thank the Co-Editor, Professor Naisyin Wang, an AE, and a referee for many helpful comments and suggestions, that greatly improved the quality of the paper. This research is supported in part by an NSF grant and by China Scholarship Council through its joint Ph.D program.

Appendix

Proof of Proposition 1. By the definitions of S , \bar{S} , and D , we can show that

$$S = O(h_{\max}^{-1}), \quad \bar{S} = O(h_{\max}^{-2}), \quad D = O(h_{\max}^{-1}). \quad (\text{A.1})$$

By the continuity of V_i and a direct algebraic manipulation, we get

$$\text{Cov}(\widehat{\text{vec}}(\beta)|\mathcal{F}_n) = A_n^{-1}B_nA_n^{-1}\{1 + o_P(1)\}, \quad (\text{A.2})$$

where $A_n = \sum_{i=1}^n (I_q \otimes X_i)^T W_i (I_q \otimes X_i)$ and

$$B_n = \sum_{i=1}^n (I_q \otimes X_i^T) K_{i\mathbf{H}}^{1/2} (\tilde{I}_i \hat{V}_i \tilde{I}_i)^{-1} K_{i\mathbf{H}}^{1/2} V_0(t) K_{i\mathbf{H}}^{1/2} (\tilde{I}_i \hat{V}_i \tilde{I}_i)^{-1} K_{i\mathbf{H}}^{1/2} (I_q \otimes X_i).$$

Set $\tilde{H} = \text{diag}\{1, h_{\max}, \dots, h_{\max}^p\}$ and $c_{ijsm} = (t_{ij} - t)^m K_{h_s}^{1/2}(t_{ij} - t)$, for $i = 1, \dots, n$, $j = 1, \dots, J$, $s = 1, \dots, q$ and $m = 0, \dots, p$. For every fixed $v = 1, \dots, 2^J - 1$, let

$$\mathcal{S}_v(h_{\max}) = \{t_{1j} \in B(t, h_{\max}) \text{ for all } j \in \Omega_v, \text{ and } t_{1,j} \notin B(t, h_{\max}) \text{ for all } j \notin \Omega_v\}.$$

Then, the existence condition of the partial density of $\{t_{ij}\}$ ensures that $\Pr\{t_{1j}$ are all equal for all $j \in \Omega_v | \mathcal{S}_v(h_{\max})\} = 1 + o(1)$ on $B(t, h_{\max})$, as $h_{\max} \rightarrow 0$. Let $a_{m+1,l+1}^{(sk)}$ denote the $((s-1)(p+1) + m + 1, (k-1)(p+1) + l + 1)$ th element of A_n , $j_v \in \Omega_v$, and $\mathbf{C}_{ism} = (0, \dots, 0, c_{i1sm}, \dots, c_{iJsm}, 0, \dots, 0)_{1 \times qJ}^T$. Then,

$$\begin{aligned} E(a_{m+1,l+1}^{(sk)}) &= \sum_{i=1}^n E\{\mathbf{C}_{ism}^T (\tilde{I}_i \widehat{V}_i \tilde{I}_i)^{-1} \mathbf{C}_{ikl}\} \\ &= n \sum_{v=1}^{2^J-1} E\{\mathbf{C}_{1sm}^T (\tilde{I}_1 \widehat{V}_1 \tilde{I}_1)^{-1} \mathbf{C}_{1kl} | \mathcal{S}_v(h_{\max})\} \Pr\{\mathcal{S}_v(h_{\max})\} \\ &= n \sum_{v=1}^{2^J-1} E[(t_{1j_v} - t)^{m+l} K_{h_s}^{1/2} (t_{1j_v} - t) K_{h_k}^{1/2} (t_{1j_v} - t) I\{\mathcal{S}_v(h_{\max})\}] \\ &\quad \times E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v_0} \widehat{V}_1 \tilde{I}_{v_0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\} \{1 + o(1)\} \\ &= n \sum_{v=1}^{2^J-1} \frac{1}{\sqrt{h_s h_k}} \int_{t-h_{\max}}^{t+h_{\max}} (u-t)^{m+l} K^{1/2} \left(\frac{u-t}{h_s}\right) K^{1/2} \left(\frac{u-t}{h_k}\right) f_v(u) du \\ &\quad \times E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v_0} \widehat{V}_1 \tilde{I}_{v_0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\} \{1 + o(1)\} \\ &= n \sum_{v=1}^{2^J-1} \frac{h_{\max}^{m+l+1}}{\sqrt{h_s h_k}} \int_{-1}^1 z^{m+l} K^{1/2} \left(\frac{h_{\max}}{h_s} z\right) K^{1/2} \left(\frac{h_{\max}}{h_k} z\right) f_v(t) dz \\ &\quad \times E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v_0} \widehat{V}_1 \tilde{I}_{v_0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\} \{1 + o(1)\} \\ &= nh_{\max}^{m+l+1} \mu_{m+l}(h_s, h_k) \sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(sk)} \{1 + o(1)\}. \end{aligned}$$

Similarly, we can show that $\{\text{var}(a_{m+1,l+1}^{(sk)})\}^{1/2} = o(nh_{\max}^{m+l})$. By combining these results, we have

$$\begin{aligned} a_{m+1,l+1}^{(sk)} &= E(a_{m+1,l+1}^{(sk)}) + O_p[\{\text{var}(a_{m+1,l+1}^{(sk)})\}^{1/2}] \\ &= nh_{\max}^{m+l+1} \mu_{m+l}(h_s, h_k) \sum_{v=1}^{2^J-1} f_v(t) \xi_v^{(sk)} \{1 + o(1)\}. \end{aligned}$$

Therefore,

$$A_n = nh_{\max} [(I_q \otimes \tilde{H}) S (I_q \otimes \tilde{H})] \{1 + o_P(1)\}. \tag{A.3}$$

Let $b_{m+1,l+1}^{(sk)}$ denote the $((s-1)(p+1) + m + 1, (k-1)(p+1) + l + 1)$ th element of B_n . Then, we have

$$E(b_{m+1,l+1}^{(sk)}) = \sum_{i=1}^n E\{\mathbf{C}_{ism}^T (\tilde{I}_i \widehat{V}_i \tilde{I}_i)^{-1} K_{i\mathbf{H}}^{1/2} V_0(t) K_{i\mathbf{H}}^{1/2} (\tilde{I}_i \widehat{V}_i \tilde{I}_i)^{-1} \mathbf{C}_{ikl}\}$$

$$\begin{aligned}
 &= n \sum_{v=1}^{2^J-1} E\{C_{1sm}^T(\tilde{I}_1 \widehat{V}_1 \tilde{I}_1)^{-1} K_{1H}^{1/2} V_0(t) K_{1H}^{1/2} (\tilde{I}_1 \widehat{V}_1 \tilde{I}_1)^{-1} C_{1kl} | \mathcal{S}_v(h_{\max})\} \Pr\{\mathcal{S}_v(h_{\max})\} \\
 &= n \sum_{v=1}^{2^J-1} \left\{ \sum_{l_1, l_2=1}^q E[(t_{1j_v} - t)^{m+l} K_{h_s}^{1/2}(t_{1j_v} - t) K_{h_k}^{1/2}(t_{1j_v} - t) \right. \\
 &\quad \times K_{h_{l_1}}^{1/2}(t_{1j_v} - t) K_{h_{l_2}}^{1/2}(t_{1j_v} - t) I\{\mathcal{S}_v(h_{\max})\}] E\{(\tilde{\mathbf{e}}_s \otimes \mathbf{1}_0)^T (\tilde{I}_{v_0} \widehat{V}_1 \tilde{I}_{v_0})^{-1} \\
 &\quad \times (E_{l_1} \otimes I_J) V_0(t) (E_{l_2} \otimes I_J) (\tilde{I}_{v_0} \widehat{V}_1 \tilde{I}_{v_0})^{-1} (\tilde{\mathbf{e}}_k \otimes \mathbf{1}_0) | \mathcal{S}_v(0)\} \{1 + o(1)\} \\
 &= n \sum_{v=1}^{2^J-1} f_v(t) h_{\max}^{m+l+1} \left\{ \sum_{l_1, l_2=1}^q \bar{\xi}_{v, l_1 l_2}^{(sk)}(t) \frac{1}{\sqrt{h_s h_k h_{l_1} h_{l_2}}} \right. \\
 &\quad \times \int_{-1}^1 z^{l+m} K^{1/2}\left(\frac{h_{\max}}{h_s} z\right) K^{1/2}\left(\frac{h_{\max}}{h_k} z\right) K^{1/2}\left(\frac{h_{\max}}{h_{l_1}} z\right) K^{1/2}\left(\frac{h_{\max}}{h_{l_2}} z\right) dz \} \{1 + o(1)\} \\
 &= n \sum_{v=1}^{2^J-1} f_v(t) h_{\max}^{m+l+1} \left\{ \sum_{l_1, l_2=1}^q \bar{\xi}_{v, l_1 l_2}^{(sk)}(t) \nu_{m+l}(h_s, h_k, h_{l_1}, h_{l_2}) \right\} \{1 + o(1)\} \\
 &= n h_{\max}^{m+l+1} \sum_{v=1}^{2^J-1} f_v(t) \nu_{m+l, v}^{(sk)}(t) \{1 + o(1)\}.
 \end{aligned}$$

As with (A.3), we have

$$B_n = n h_{\max} [(I_q \otimes \tilde{H}) \bar{S} (I_q \otimes \tilde{H})] \{1 + o_P(1)\}. \tag{A.4}$$

By combining (A.1)–(A.4), we have

$$\text{Cov}\{\widehat{\mathbf{m}}^{(k)}(t) | \mathcal{F}_n\} = \frac{k!^2}{n h_{\max}^{1+2k}} [(I_q \otimes \mathbf{e}_{k+1}^T) S^{-1} \bar{S} S^{-1} (I_q \otimes \mathbf{e}_{k+1})] + o_P\left(\frac{1}{n h_{\max}^{1+2k}}\right).$$

Similar to the asymptotic expansion of B_n in (A.4), we can show that

$$\begin{aligned}
 \text{Bias}\{\widehat{\mathbf{m}}^{(k)}(t)\} &= k! (I_q \otimes \mathbf{e}_{k+1}^T) [E(\widehat{\text{vec}}(\beta) | \mathcal{F}_n) - \text{vec}(\beta)] \\
 &= k! (I_q \otimes \mathbf{e}_{k+1}^T) A_n^{-1} \sum_{i=1}^n (I_q \otimes X_i^T) W_i E[\text{vec}(Y_i) - (I_q \otimes X_i) \text{vec}(\beta)] \\
 &= \frac{k!}{(p+1)!} (I_q \otimes \mathbf{e}_{k+1}^T) A_n^{-1} \sum_{i=1}^n (I_q \otimes X_i^T) W_i \begin{pmatrix} m_1^{(p+1)}(t) \\ \vdots \\ m_q^{(p+1)}(t) \end{pmatrix}
 \end{aligned}$$

$$\begin{aligned}
& \otimes \begin{pmatrix} (t_{i1} - t)^{p+1} \\ \vdots \\ (t_{iJ} - t)^{p+1} \end{pmatrix} \{1 + o(1)\} \\
&= \frac{nk!h_{\max}^{p+2}}{(p+1)!} (I_q \otimes \mathbf{e}_{k+1}^T) \{nh_{\max}^{-1} [(I_q \otimes \tilde{H})S(I_q \otimes \tilde{H})] \{1 + o_P(1)\}\}^{-1} \\
& \quad \times [(I_q \otimes \tilde{H})\mathbf{D}] \{1 + o_P(1)\} \\
&= \frac{k!}{(p+1)!} h_{\max}^{p+1-k} [(I_q \otimes \mathbf{e}_{k+1}^T S^{-1} \mathbf{D}) + o_P(h_{\max}^{p+1-k})].
\end{aligned}$$

The last equation holds because $\mathbf{D} = O(h_{\max}^{-1})$, as specified in (A.1). By now, we have proved (3.1) and (3.2).

Proof of Corollary 1. Similar to the proof of Proposition 1, we can show that

$$A_n = n[M \otimes \tilde{H}S_1\tilde{H}] \{1 + o_P(1)\} \quad (\text{A.5})$$

$$B_n = nh^{-1}[N \otimes \tilde{H}\tilde{S}_1\tilde{H}] \{1 + o_P(1)\} \quad (\text{A.6})$$

$$\sum_{i=1}^n (I_q \otimes X_i^T) W_i E[\text{vec}(Y_i) - (I_q \otimes X_i)\text{vec}(\beta)] = [\mathbf{C} \otimes \tilde{H}c_p] \{1 + o_P(1)\}. \quad (\text{A.7})$$

The results (3.3) and (3.4) can be obtained from combining (A.5)–(A.7).

Proof of Corollary 2. In cases when response components are independent, the covariance matrices V_i are block diagonal. By combining this result with those in (3.1) and (3.2) in Proposition 1, the conclusions (3.5) and (3.6) are straightforward.

Proof of Corollary 3. The proof of Corollary 3 is similar to the one of Proposition 1. Thus, it is omitted here.

References

- Chen, K. and Jin, Z. (2005). Local polynomial regression analysis of clustered data. *Biometrika* **92**, 59-74.
- Coull, B. A. and Staudenmayer, J. (2004). Self-modeling regression for multivariate curve data. *Statist. Sinica* **14**, 695-711.
- Cupples, L. A. et al. (2007). The Framingham Heart Study 100K SNP genome-wide association study resource: overview of 17 phenotype working group reports. *BMC Medical Genetics* **8** (Suppl 1), S1.
- Fieuwis, S. and Verbeke, G. (2006). Pairwise fitting of mixed models for the joint modeling of multivariate longitudinal profiles. *Biometrics* **62**, 424-431.
- Fitzmaurice, G. M., Laird, N. M. and Ware, J. H. (2004). Applied longitudinal analysis. *John Wiley & Sons, New York*

- Gray, S. M. and Brookmeyer, R. (2000). Multidimensional longitudinal data: estimating treatment effect from continuous, discrete or time-to-event response variable. *J. Amer. Statist. Assoc.* **95**, 396-406.
- Godleski, J. J. Verrier, R. L. Koutrakis, P. Catalano, P. et al. (2000). Mechanisms of morbidity and mortality from exposure to ambient air particles. *Research Report, Health Effects Institute* **91**, 5-103.
- Hall, P. and Robinson, A. P. (2009). Reducing variability of cross-validation for smoothing-parameter choice. *Biometrika* **96**, 175-186.
- Li, Y. (2011). Efficient semiparametric regression for longitudinal data with nonparametric covariance estimation. *Biometrika* **98**, 355-370.
- Liang, K. Y. and Zeger, S. L. (1986). Longitudinal data analysis using generalized linear models. *Biometrika* **73**, 13-22.
- Lin, X. and Carroll, R. (2000). Nonparametric function estimation for clustered data when the predictor is measured without/with error. *J. Am. Statist. Assoc.* **95**, 520-534.
- Lin, X. and Carroll, R. (2001). Semiparametric regression for clustered data using generalized estimating equations. *J. Am. Statist. Assoc.* **96**, 1045-1056.
- O'Brien, L. M. and Fitzmaurice, G. M. (2004). Analysis of longitudinal multiple-source binary data using generalized estimating equations. *Appl. Statist.* **53**, 177-193.
- Ruppert, D. and Wand, M. P. (1994). Multivariate locally weighted least squares regression. *Ann. Statist.* **22**, 1346-1370.
- Roy, J. and Lin, X. (2000). Latent variable model for longitudinal data with multiple continuous outcomes. *Biometrics* **56**, 1047-1054.
- Wang, N. (2003). Marginal nonparametric kernel regression accounting for within-subject correlation. *Biometrika* **90**, 43-52.
- Weiss, R. E. (2005). *Modeling Longitudinal Data*. Springer, USA.

School of Finance and Statistics, East China Normal University, Shanghai 200241, China.

E-mail: terryxdd@gmail.com

School of Statistics, University of Minnesota, 389 Ford Hall, 224 Church Street SE, Minneapolis, MN 55455, USA.

E-mail: qiuxx008@umn.edu

School of Finance and Statistics, East China Normal University, Shanghai 200241, China.

E-mail: xlpu@stat.ecnu.edu.cn

(Received December 2011; accepted July 2012)