

Detailed Asymptotic Analysis of k -recursive Sequences

Daniel Krenn



June 29, 2023



This presentation is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 3.0 Unported License.

(Un-)bordered Factors

(Un-)bordered

- word w **bordered**:
 - exists non-empty word $v \neq w$
 - v is prefix and suffix of w
- otherwise **unbordered**

(Un-)bordered Factors

(Un-)bordered

- word w **bordered**:
 - exists non-empty word $v \neq w$
 - v is prefix and suffix of w
- otherwise **unbordered**

bordered factor	border	length
00	0	2
11	1	2
010	0	3
101	1	3
1010	10	4
0110100110	0110	10

unbordered factor	length
ε	0
0	1
1	1
01	2
10	2
011	3
110	3
100	3
001	3

(Un-)bordered Factors

(Un-)bordered

- word w **bordered**:
 - exists non-empty word $v \neq w$
 - v is prefix and suffix of w
- otherwise **unbordered**

bordered factor	border	length
$t[5..6] = 00$	0	2
$t[1..2] = 11$	1	2
$t[3..5] = 010$	0	3
$t[2..4] = 101$	1	3
$t[2..5] = 1010$	10	4
$t[0..9] = 0110100110$	0110	10

unbordered factor	length
ε	0
$t[0..0] = 0$	1
$t[1..1] = 1$	1
$t[0..1] = 01$	2
$t[2..3] = 10$	2
$t[0..2] = 011$	3
$t[1..3] = 110$	3
$t[4..6] = 100$	3
$t[5..7] = 001$	3

Thue–Morse Sequence

$t = 01101001\ 10010110\ 10010110\ 01101001\ \dots$

Number of Unbordered Factors

Theorem (Goč–Henshall–Shallit 2013)

*exists unbordered factor
of length n
in Thue–Morse sequence*

\iff

$$(n)_2 \notin 1(01^*0)^*10^*1$$

Number of Unbordered Factors

Theorem (Goč–Henshall–Shallit 2013)

*exists unbordered factor
of length n
in Thue–Morse sequence* $\iff (n)_2 \notin 1(01^*0)^*10^*1$

- number $f(n)$ of unbordered factors of length n in the Thue–Morse sequence

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(n)$	1	2	2	4	2	4	6	0	4	4	4	4	12	0	4	4

Number of Unbordered Factors

Theorem (Goč–Henshall–Shallit 2013)

*exists unbordered factor
of length n
in Thue–Morse sequence*

$$\iff (n)_2 \notin 1(01^*0)^*10^*1$$

- number $f(n)$ of unbordered factors of length n in the Thue–Morse sequence

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$f(n)$	1	2	2	4	2	4	6	0	4	4	4	4	12	0	4	4

Theorem (Goč–Mousavi–Shallit 2013)

- inequality $f(n) \leq n$ holds for all $n \geq 4$*
- $f(n) = n$ infinitely often*
- $\limsup_{n \geq 1} \frac{f(n)}{n} = 1$*

Recurrence Relations

- number $f(n)$ of unbordered factors of length n in Thue–Morse sequence
- recurrence relations

$$\begin{aligned}f(4n) &= 2f(2n) && (n \geq 2) \\f(4n + 1) &= f(2n + 1) && (n \geq 0) \\f(8n + 2) &= f(2n + 1) + f(4n + 3) && (n \geq 1) \\f(8n + 3) &= -f(2n + 1) + f(4n + 2) && (n \geq 2) \\f(8n + 6) &= -f(2n + 1) + f(4n + 2) + f(4n + 3) && (n \geq 2) \\f(8n + 7) &= 2f(2n + 1) + f(4n + 3) && (n \geq 3)\end{aligned}$$



Theorem (Goč–Mousavi–Shallit 2013)

$f(n)$ satisfies recurrence relations above

Recurrence Relations

- number $f(n)$ of unbordered factors of length n in Thue–Morse sequence
- recurrence relations

$$\begin{array}{ll}
 f(8n) = 2f(4n) & (n \geq 1) \\
 f(8n + 1) = f(4n + 1) & (n \geq 0) \\
 f(8n + 2) = f(4n + 1) + f(4n + 3) & (n \geq 1) \\
 f(8n + 3) = -f(4n + 1) + f(4n + 2) & (n \geq 2) \\
 f(8n + 4) = 2f(4n + 2) & (n \geq 1) \\
 f(8n + 5) = f(4n + 3) & (n \geq 0) \\
 f(8n + 6) = -f(4n + 1) + f(4n + 2) + f(4n + 3) & (n \geq 2) \\
 f(8n + 7) = 2f(4n + 1) + f(4n + 3) & (n \geq 3)
 \end{array}$$



Recurrence Relations

- number $f(n)$ of unbordered factors of length n in Thue–Morse sequence
- recurrence relations

$$f(8n) = 2f(4n) \quad (n \geq 1)$$

$$f(8n + 1) = f(4n + 1) \quad (n \geq 0)$$

$$f(8n + 2) = f(4n + 1) + f(4n + 3) \quad (n \geq 1)$$

$$f(8n + 3) = -f(4n + 1) + f(4n + 2) \quad (n \geq 2)$$

$$f(8n + 4) = 2f(4n + 2) \quad (n \geq 1)$$

$$f(8n + 5) = f(4n + 3) \quad (n \geq 0)$$

$$f(8n + 6) = -f(4n + 1) + f(4n + 2) + f(4n + 3) \quad (n \geq 2)$$

$$f(8n + 7) = 2f(4n + 1) + f(4n + 3) \quad (n \geq 3)$$

- $f(n)$ is a 2-recursive sequence



k -recursive Sequence

- integer $k \geq 2$

k -recursive Sequence $x(n)$

there exist

- integers $M > m \geq 0$, $\ell \leq u$,
 $n_0 \geq \max\{-\ell/k^m, 0\}$
- constants $c_{s,j} \in \mathbb{C}$

such that

$$x(k^M n + s) = \sum_{\ell \leq j \leq u} c_{s,j} x(k^m n + j)$$

holds for all $n \geq n_0$ and $0 \leq s < k^M$

k -recursive Sequence

- integer $k \geq 2$

k -recursive Sequence $x(n)$

there exist

- integers $M > m \geq 0$, $\ell \leq u$,
 $n_0 \geq \max\{-\ell/k^m, 0\}$
- constants $c_{s,j} \in \mathbb{C}$

such that

$$x(k^M n + s) = \sum_{\ell \leq j \leq u} c_{s,j} x(k^m n + j)$$

holds for all $n \geq n_0$ and $0 \leq s < k^M$

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2n) = 2h(n)$, $h(2n+1) = 2h(n)$ for $n \geq 1$, $h(1) = 1$
 - $k = 2$, $M = 1$, $m = 0$, $\ell = 0$, $u = 1$, $n_0 = 1$

k -recursive Sequence

- integer $k \geq 2$

k -recursive Sequence $x(n)$

there exist

- integers $M > m \geq 0$, $\ell \leq u$,
 $n_0 \geq \max\{-\ell/k^m, 0\}$
- constants $c_{s,j} \in \mathbb{C}$

such that

$$x(k^M n + s) = \sum_{\ell \leq j \leq u} c_{s,j} x(k^m n + j)$$

holds for all $n \geq n_0$ and $0 \leq s < k^M$

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2n) = 2h(n)$, $h(2n+1) = 2h(n)$ for $n \geq 1$, $h(1) = 1$
 - $k = 2$, $M = 1$, $m = 0$, $\ell = 0$, $u = 1$, $n_0 = 1$
- binary sum of digits
 - $s(2n) = s(n)$, $s(2n+1) = s(n) + 1$
 - no direct fit because of constant sequence
 - deal with **inhomogeneities** by increasing the exponents

k -recursive Sequence

- integer $k \geq 2$

k -recursive Sequence $x(n)$

there exist

- integers $M > m \geq 0$, $\ell \leq u$,
 $n_0 \geq \max\{-\ell/k^m, 0\}$
- constants $c_{s,j} \in \mathbb{C}$

such that

$$x(k^M n + s) = \sum_{\ell \leq j \leq u} c_{s,j} x(k^m n + j)$$

holds for all $n \geq n_0$ and $0 \leq s < k^M$

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2n) = 2h(n)$, $h(2n+1) = 2h(n)$ for $n \geq 1$, $h(1) = 1$
 - $k = 2$, $M = 1$, $m = 0$, $\ell = 0$, $u = 1$, $n_0 = 1$
- binary sum of digits
 - $s(2n) = s(n)$, $s(2n+1) = s(n) + 1$
 - no direct fit because of constant sequence
 - deal with **inhomogeneities** by increasing the exponents
- number of comparisons for MergeSort

k -recursive Sequence

- integer $k \geq 2$

k -recursive Sequence $x(n)$

there exist

such that

- integers $M > m \geq 0$, $\ell \leq u$,
 $n_0 \geq \max\{-\ell/k^m, 0\}$
- constants $c_{s,j} \in \mathbb{C}$

$$x(k^M n + s) = \sum_{\ell \leq j \leq u} c_{s,j} x(k^m n + j)$$

holds for all $n \geq n_0$ and $0 \leq s < k^M$

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2n) = 2h(n)$, $h(2n+1) = 2h(n)$ for $n \geq 1$, $h(1) = 1$
 - $k = 2$, $M = 1$, $m = 0$, $\ell = 0$, $u = 1$, $n_0 = 1$
- binary sum of digits
 - $s(2n) = s(n)$, $s(2n+1) = s(n) + 1$
 - no direct fit because of constant sequence
 - deal with **inhomogeneities** by increasing the exponents
- number of comparisons for MergeSort
- number of **unbordered factors** of length n in Thue–Morse sequence

k -linear Representation

binary sum of digits $s(n)$:

- recurrence relations

even numbers: $s(2n) = s(n)$

odd numbers: $s(2n + 1) = s(n) + 1$

k -linear Representation

binary sum of digits $s(n)$:

- recurrence relations

even numbers: $s(2n) = s(n)$

odd numbers: $s(2n + 1) = s(n) + 1$

- vector-valued sequence

set $v(n) = (s(n), 1)^T$

even $v(2n) = \begin{pmatrix} s(n) \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} v(n)$

odd $v(2n + 1) = \begin{pmatrix} s(n) + 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} v(n)$

k -linear Representation

binary sum of digits $s(n)$:

- recurrence relations

even numbers: $s(2n) = s(n)$

odd numbers: $s(2n + 1) = s(n) + 1$

- vector-valued sequence

set $v(n) = (s(n), 1)^T$

even $v(2n) = \begin{pmatrix} s(n) \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} v(n)$

odd $v(2n + 1) = \begin{pmatrix} s(n) + 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} v(n)$

- iterate \rightsquigarrow product of matrices

k -linear Representation

binary sum of digits $s(n)$:

- recurrence relations

even numbers: $s(2n) = s(n)$

odd numbers: $s(2n+1) = s(n) + 1$

- vector-valued sequence

set $v(n) = (s(n), 1)^T$

even $v(2n) = \begin{pmatrix} s(n) \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} v(n)$

odd $v(2n+1) = \begin{pmatrix} s(n)+1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} v(n)$

- iterate \rightsquigarrow product of matrices

k -regular Sequence $f(n)$

- square matrices M_0, \dots, M_{k-1}
- vectors u and w
- k -linear representation

$$f(n) = u^T M_{n_0} M_{n_1} \dots M_{n_{\ell-1}} w$$

with standard k -ary expansion

$$n = (n_{\ell-1} \dots n_1 n_0)_k$$

Some k -regular Sequences

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2^j n + r) = 2^j h(n)$ for $n \geq 1, j \geq 0, 0 \leq r < 2^j$
- binary sum of digits

Some k -regular Sequences

- $h(n)$... largest power of 2 less than or equal to n
 - $h(2^j n + r) = 2^j h(n)$ for $n \geq 1, j \geq 0, 0 \leq r < 2^j$
- binary sum of digits
- k -recursive sequences:

Theorem (Heuberger–K–Lipnik 2022)

- k -recursive sequence $x(n)$

Then

- $x(n)$ is k -regular sequence
- k -linear representation of $x(n)$
 - vector-valued sequence $v(n)$ in block form
 - block matrices M_0, \dots, M_{k-1}
 - computed by coefficients of k -recursive sequence
 - explicit formulæ for the rows available

Unbordered Factors: Coefficient Matrices

- number $f(n)$ of unbordered factors of length n in Thue–Morse sequence



$$f(8n) = 2f(4n)$$

$$f(8n + 1) = f(4n + 1)$$

$$f(8n + 2) = f(4n + 1) + f(4n + 3)$$

$$f(8n + 3) = -f(4n + 1) + f(4n + 2)$$

$$f(8n + 4) = 2f(4n + 2)$$

$$f(8n + 5) = f(4n + 3)$$

$$f(8n + 6) = -f(4n + 1) + f(4n + 2) + f(4n + 3)$$

$$f(8n + 7) = 2f(4n + 1) + f(4n + 3)$$

Unbordered Factors: Coefficient Matrices

- number $f(n)$ of unbordered factors of length n in Thue–Morse sequence
- coefficient matrices B_0, B_1 :



$$f(8n) = 2f(4n)$$

$$f(8n+1) = f(4n+1)$$

$$f(8n+2) = f(4n+1) + f(4n+3)$$

$$f(8n+3) = -f(4n+1) + f(4n+2)$$

$$f(8n+4) = 2f(4n+2)$$

$$f(8n+5) = f(4n+3)$$

$$f(8n+6) = -f(4n+1) + f(4n+2) + f(4n+3)$$

$$f(8n+7) = 2f(4n+1) + f(4n+3)$$

 \implies

$$B_0 = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & 1 & 0 \end{pmatrix}$$

$$B_1 = \begin{pmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 1 \\ 0 & 2 & 0 & 1 \end{pmatrix}$$

Unbordered Factors: Linear Representation

- coefficient matrices B_0, B_1
- 2-linear representation of $f(n)$:

$$v = \begin{pmatrix} f \\ f \circ (n \mapsto 2n) \\ f \circ (n \mapsto 2n+1) \\ f \circ (n \mapsto 4n) \\ f \circ (n \mapsto 4n+1) \\ f \circ (n \mapsto 4n+2) \\ f \circ (n \mapsto 4n+3) \end{pmatrix}$$

$$M_0 = \begin{pmatrix} J_{00} & J_{01} \\ 0 & B_0 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} J_{10} & J_{11} \\ 0 & B_1 \end{pmatrix}$$

- J_{r0}, J_{r1} entries 0, 1



Unbordered Factors: Linear Representation

- coefficient matrices B_0, B_1
- 2-linear representation of $f(n)$:

$$v = \begin{pmatrix} f \\ f \circ (n \mapsto 2n) \\ f \circ (n \mapsto 2n+1) \\ f \circ (n \mapsto 4n) \\ f \circ (n \mapsto 4n+1) \\ f \circ (n \mapsto 4n+2) \\ f \circ (n \mapsto 4n+3) \end{pmatrix}$$

$$M_0 = \begin{pmatrix} J_{00} & J_{01} \\ 0 & B_0 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} J_{10} & J_{11} \\ 0 & B_1 \end{pmatrix}$$



- J_{r0}, J_{r1} entries 0, 1
- initial value compensation \rightsquigarrow 2-linear representation of $f(n)$:

$$\tilde{M}_0 = \begin{pmatrix} M_0 & W_0 \\ 0 & J_0 \end{pmatrix} \quad \text{and} \quad \tilde{M}_1 = \begin{pmatrix} M_1 & W_1 \\ 0 & J_1 \end{pmatrix}$$

- J_r entries 0, 1
- W_r entries from initial values

Unbordered Factors: Linear Representation

- coefficient matrices B_0, B_1
- 2-linear representation of $f(n)$:

$$v = \begin{pmatrix} f \\ f \circ (n \mapsto 2n) \\ f \circ (n \mapsto 2n+1) \\ f \circ (n \mapsto 4n) \\ f \circ (n \mapsto 4n+1) \\ f \circ (n \mapsto 4n+2) \\ f \circ (n \mapsto 4n+3) \end{pmatrix}$$

$$M_0 = \begin{pmatrix} J_{00} & J_{01} \\ 0 & B_0 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} J_{10} & J_{11} \\ 0 & B_1 \end{pmatrix}$$



- J_{r0}, J_{r1} entries 0, 1
- initial value compensation \rightsquigarrow 2-linear representation of $f(n)$:

$$\tilde{M}_0 = \begin{pmatrix} M_0 & W_0 \\ 0 & J_0 \end{pmatrix} \quad \text{and} \quad \tilde{M}_1 = \begin{pmatrix} M_1 & W_1 \\ 0 & J_1 \end{pmatrix}$$

- J_r entries 0, 1
- W_r entries from initial values
- minimization algorithm: dimension 10 \rightsquigarrow dimension 8

Asymptotics of Partial Sums

- k -regular sequence $f(n)$
- partial sums $F(N) = \sum_{n < N} f(n)$

Asymptotics of Partial Sums

- k -regular sequence $f(n)$
- partial sums $F(N) = \sum_{n < N} f(n)$

Theorem (Heuberger–K–Prodinger 2018, Heuberger–K 2020)

$$F(N) = \sum_{\substack{\lambda \in \sigma(M_0 + \dots + M_{k-1}) \\ |\lambda| > \rho}} N^{\log_k \lambda} \sum_{0 \leq \ell < m(\lambda)} (\log_k N)^\ell \Phi_{\lambda \ell}(\{\log_k N\}) + O(N^{\log_k R} (\log N)^{\widehat{m}})$$

- 1-periodic (Hölder) continuous functions $\Phi_{\lambda \ell}$
- functional equation

$$\left(1 - \frac{1}{k^s} (M_0 + \dots + M_{k-1})\right) \mathcal{V}(s) = \sum_{n=1}^{k-1} \frac{v(n)}{n^s} + \frac{1}{k^s} \sum_{r=0}^{k-1} M_r \sum_{\ell \geq 1} \binom{-s}{\ell} \left(\frac{r}{k}\right)^\ell \mathcal{V}(s+\ell)$$

- meromorphic continuation on the half plane $\Re s > \log_k R$
- Fourier series $\Phi_{\lambda \ell}(u) = \sum_{h \in \mathbb{Z}} \varphi_{\lambda \ell h} \exp(2\ell \pi i u)$

$$\varphi_{\lambda \ell h} = \frac{(\log k)^\ell}{\ell!} \operatorname{Res} \left(\frac{(f(0) + \mathcal{F}(s)) \left(s - \log_k \lambda - \frac{2h\pi i}{\log k}\right)^\ell}{s}, s = \log_k \lambda + \frac{2h\pi i}{\log k} \right)$$

Unbordered Factors: towards Asymptotics



Asymptotics of k -recursive Sequences

Only properties of coefficient matrices needed from k -linear representation!

- coefficient matrices

$$B_0 = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & 1 & 0 \end{pmatrix}$$

$$B_1 = \begin{pmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 1 \\ 0 & 2 & 0 & 1 \end{pmatrix}$$

Unbordered Factors: towards Asymptotics



Asymptotics of k -recursive Sequences

Only properties of coefficient matrices needed from k -linear representation!

- coefficient matrices

$$B_0 = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & 1 & 0 \end{pmatrix}$$

$$B_1 = \begin{pmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 1 \\ 0 & 2 & 0 & 1 \end{pmatrix}$$

- spectrum

$$\sigma(B_0 + B_1) = \{1 - \sqrt{3}, 1, 2, 1 + \sqrt{3}\}$$

Unbordered Factors: towards Asymptotics



Asymptotics of k -recursive Sequences

Only properties of coefficient matrices needed from k -linear representation!

- coefficient matrices

$$B_0 = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & -1 & 1 & 0 \end{pmatrix}$$

$$B_1 = \begin{pmatrix} 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & -1 & 1 & 1 \\ 0 & 2 & 0 & 1 \end{pmatrix}$$

- spectrum

$$\sigma(B_0 + B_1) = \{1 - \sqrt{3}, 1, 2, 1 + \sqrt{3}\}$$

- joint spectral radius of $\{B_0, B_1\}$ is 2
- has simple growth property


Unbordered Factors: Asymptotics

- number $f(n)$ of unbordered factors of length n in the Thue–Morse sequence

Theorem (Heuberger–K–Lipnik 2022)

$$F(N) = \sum_{0 \leq n < N} f(n) = N^\kappa \cdot \Phi_F(\{\log_2 N\}) + O(N \log N) \quad \text{as } N \rightarrow \infty$$

- • $\kappa = \log_2(1 + \sqrt{3}) = 1.44998431347650\dots$
- • 1-periodic continuous function Φ_F , Hölder continuous with any exponent smaller than $\kappa - 1$
- • explicit functional equation for Dirichlet series + analyticity properties, poles
- • efficiently computable Fourier coefficients of Φ_F



Stern's Diatomic Sequence



Stern's Diatomic Sequence

$$d(2n) = d(n)$$

$$d(2n+1) = d(n) + d(n+1)$$

for all $n \geq 0$ and

$$d(0) = 0, d(1) = 1$$

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$d(n)$	0	1	1	2	1	3	2	3	1	4	3	5	2	5	3	4

Stern's Diatomic Sequence



Stern's Diatomic Sequence

$$d(2n) = d(n)$$

$$d(2n+1) = d(n) + d(n+1)$$

for all $n \geq 0$ and

$$d(0) = 0, d(1) = 1$$

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$d(n)$	0	1	1	2	1	3	2	3	1	4	3	5	2	5	3	4

- number of different hyperbinary representations
(Northshield 2010)
- number of integers $r \in \mathbb{N}_0$ such that Stirling partition numbers $\left\{ \begin{smallmatrix} n \\ 2r \end{smallmatrix} \right\}$ are even and non-zero
(Carlitz 1964)
- number of different representations as a sum of distinct Fibonacci numbers F_{2k}
(Bicknell-Johnson 2003)
- number of different alternating bit sets
(Finch 2003)
- relation to the Towers of Hanoi
(Hinz-Klavžar-Milutinović-Parisse-Petr 2005)


Stern's Diatomic Sequence: Asymptotics

- Stern's diatomic sequence $d(n)$
- 2-recursive, 2-regular

Asymptotics

$$D(N) = \sum_{0 \leq n < N} d(n) = N^\kappa \cdot \Phi_D(\{\log_2 N\}) + O(N^{\log_2 \varphi}) \quad \text{as } N \rightarrow \infty$$

- • $\kappa = \log_2 3 = 1.5849625007211\dots$
- • $\varphi = \frac{1+\sqrt{5}}{2} = 1.6180339887498\dots$, $\log_2 \varphi = 0.69424191363061\dots$
- • 1-periodic continuous function Φ_D ,
Hölder continuous with any exponent smaller than $\kappa - \log_2 \varphi$
- • explicit functional equation for Dirichlet series
+ analyticity properties, poles
- • efficiently computable Fourier coefficients of Φ_D



Generalized Pascal's Triangle

Binomial Coefficients of Words

binomial coefficient $\binom{u}{v}$ equals
number of different occurrences of v
as a scattered subword of u



Generalized Pascal's Triangle

Binomial Coefficients of Words

binomial coefficient $\binom{u}{v}$ equals
 number of different occurrences of v
 as a scattered subword of u



n	$u = (n)_2$	k	$v = (k)_2$								$z(n)$	
			0	1	2	3	4	5	6	7		8
	ε		ε	1	10	11	100	101	110	111	1000	
0	ε		1	0	0	0	0	0	0	0	0	1
1	1		1	1	0	0	0	0	0	0	0	2
2	10		1	1	1	0	0	0	0	0	0	3
3	11		1	2	0	1	0	0	0	0	0	3
4	100		1	1	2	0	1	0	0	0	0	4
5	101		1	2	1	1	0	1	0	0	0	5
6	110		1	2	2	1	0	0	1	0	0	5
7	111		1	3	0	3	0	0	0	1	0	4
8	1000		1	1	3	0	3	0	0	0	1	5

Generalized Pascal's Triangle

Binomial Coefficients of Words


binomial coefficient $\binom{u}{v}$ equals
 number of different occurrences of v
 as a scattered subword of u

"classical"

binomial coefficient
 $\binom{n}{k} = \binom{1^n}{1^k}$

n	k $v = \binom{k}{2}$ $u = \binom{n}{2}$	0	1	2	3	4	5	6	7	8	$z(n)$
		ε	1	10	11	100	101	110	111	1000	
0	ε	1	0	0	0	0	0	0	0	0	1
1	1	1	1	0	0	0	0	0	0	0	2
2	10	1	1	1	0	0	0	0	0	0	3
3	11	1	2	0	1	0	0	0	0	0	3
4	100	1	1	2	0	1	0	0	0	0	4
5	101	1	2	1	1	0	1	0	0	0	5
6	110	1	2	2	1	0	0	1	0	0	5
7	111	1	3	0	3	0	0	0	1	0	4
8	1000	1	1	3	0	3	0	0	0	1	5

Non-zeros in Generalized Pascal's Triangle



Stern's Diatomic Sequence & Generalized Pascal's Triangle

- Stern's diatomic sequence $d(n)$
- number $z(n)$ of non-zero elements in n th row of generalized Pascal's triangle $\binom{n}{k}_2$

Theorem (Leroy–Rigo–Stipulanti 2017)

$$z(n) = d(2n + 1) \quad \text{for all } n \geq 0$$



Stern's Diatomic Sequence & Generalized Pascal's Triangle

- Stern's diatomic sequence $d(n)$
- number $z(n)$ of non-zero elements in n th row of generalized Pascal's triangle $\binom{n}{k}_2$



Theorem (Leroy–Rigo–Stipulanti 2017)

$$z(n) = d(2n + 1) \quad \text{for all } n \geq 0$$

Theorem (Leroy–Rigo–Stipulanti 2017)

recurrence relations

$$z(2n + 1) = 3z(n) - z(2n)$$

$$z(4n) = -z(n) + 2z(2n)$$

$$z(4n + 2) = 4z(n) - z(2n)$$

for all $n \geq 0$



Generalized Pascal's Triangle: Asymptotic Analysis

- reformulate as 2-recursive sequence:

$$z(4n) = \frac{5}{3}z(2n) - \frac{1}{3}z(2n+1)$$

$$z(4n+1) = \frac{4}{3}z(2n) + \frac{1}{3}z(2n+1)$$

$$z(4n+2) = \frac{1}{3}z(2n) + \frac{4}{3}z(2n+1)$$

$$z(4n+3) = -\frac{1}{3}z(2n) + \frac{5}{3}z(2n+1)$$



Generalized Pascal's Triangle: Asymptotic Analysis

- reformulate as 2-recursive sequence & read off coefficient matrices:

$$\begin{aligned}z(4n) &= \frac{5}{3}z(2n) - \frac{1}{3}z(2n+1) \\z(4n+1) &= \frac{4}{3}z(2n) + \frac{1}{3}z(2n+1) \\z(4n+2) &= \frac{1}{3}z(2n) + \frac{4}{3}z(2n+1) \\z(4n+3) &= -\frac{1}{3}z(2n) + \frac{5}{3}z(2n+1)\end{aligned} \quad \Rightarrow \quad \begin{aligned}B_0 &= \frac{1}{3} \begin{pmatrix} 5 & -1 \\ 4 & 1 \end{pmatrix} \\B_1 &= \frac{1}{3} \begin{pmatrix} 1 & 4 \\ -1 & 5 \end{pmatrix}\end{aligned}$$

- 2-linear representation of dimension 3



Generalized Pascal's Triangle: Asymptotic Analysis

- reformulate as 2-recursive sequence & read off coefficient matrices:

$$\begin{aligned}z(4n) &= \frac{5}{3}z(2n) - \frac{1}{3}z(2n+1) \\z(4n+1) &= \frac{4}{3}z(2n) + \frac{1}{3}z(2n+1) \\z(4n+2) &= \frac{1}{3}z(2n) + \frac{4}{3}z(2n+1) \\z(4n+3) &= -\frac{1}{3}z(2n) + \frac{5}{3}z(2n+1)\end{aligned} \quad \Rightarrow \quad \begin{aligned}B_0 &= \frac{1}{3} \begin{pmatrix} 5 & -1 \\ 4 & 1 \end{pmatrix} \\B_1 &= \frac{1}{3} \begin{pmatrix} 1 & 4 \\ -1 & 5 \end{pmatrix}\end{aligned}$$

- 2-linear representation of dimension 3
- towards asymptotics: spectrum & joint spectral radius
- investigating eigenstructure \rightsquigarrow no error term



Generalized Pascal's Triangle: Asymptotic Analysis

- reformulate as 2-recursive sequence & read off coefficient matrices:

$$\begin{aligned}
 z(4n) &= \frac{5}{3}z(2n) - \frac{1}{3}z(2n+1) \\
 z(4n+1) &= \frac{4}{3}z(2n) + \frac{1}{3}z(2n+1) \\
 z(4n+2) &= \frac{1}{3}z(2n) + \frac{4}{3}z(2n+1) \\
 z(4n+3) &= -\frac{1}{3}z(2n) + \frac{5}{3}z(2n+1)
 \end{aligned}
 \implies
 \begin{aligned}
 B_0 &= \frac{1}{3} \begin{pmatrix} 5 & -1 \\ 4 & 1 \end{pmatrix} \\
 B_1 &= \frac{1}{3} \begin{pmatrix} 1 & 4 \\ -1 & 5 \end{pmatrix}
 \end{aligned}$$

- 2-linear representation of dimension 3
- towards asymptotics: spectrum & joint spectral radius
- investigating eigenstructure \rightsquigarrow no error term
- reconsider connection to Stern's diatomic sequence
- (compute Fourier coefficients of fluctuation)




Generalized Pascal's Triangle: Asymptotics

- number $z(n)$ of non-zero elements in n th row of generalized Pascal's triangle $\binom{(n)_2}{(k)_2}$

Asymptotics

$$Z(N) = \sum_{0 \leq n < N} z(n) = N^\kappa \cdot \Phi_Z(\{\log_2 N\}) \quad \text{for } N \geq 1$$

- $\kappa = \log_2 3$
- $\Phi_Z = 2\Phi_D$ (with Φ_D from Stern's diatomic sequence)




Generalized Pascal's Triangle: Asymptotics

- number $z(n)$ of non-zero elements in n th row of generalized Pascal's triangle $\binom{n}{k}_2$

Asymptotics

$$Z(N) = \sum_{0 \leq n < N} z(n) = N^\kappa \cdot \Phi_Z(\{\log_2 N\}) \quad \text{for } N \geq 1$$

- $\kappa = \log_2 3$
- $\Phi_Z = 2\Phi_D$ (with Φ_D from Stern's diatomic sequence)



Corollary (Heuberger–K–Lipnik 2022)

- Stern's diatomic sequence $d(n)$

$$\sum_{0 \leq n < N} d(n) + \frac{1}{2}d(N) = N^\kappa \cdot \Phi_D(\{\log_2 N\})$$