

Statistica Sinica Preprint No: SS-12-322R2

Title	Testing for change points due to a covariate threshold in regression
Manuscript ID	SS-12-322R2
URL	http://www.stat.sinica.edu.tw/statistica/
DOI	10.5705/ss.2012.322
Complete List of Authors	Liwen Zhang Huixia Judy Wang and Zhongyi Zhu
Corresponding Author	Zhongyi Zhu
E-mail	zhuzy@fudan.edu.cn
Notice: Accepted version subject to English editing.	

TESTING FOR CHANGE POINTS DUE TO A COVARIATE THRESHOLD IN QUANTILE REGRESSION

Liwen Zhang¹, Huixia Judy Wang², and Zhongyi Zhu³

Fudan University^{1,3}, North Carolina State University²

Abstract: We develop a new procedure for testing change points due to a covariate threshold in regression quantiles. The proposed test is based on the CUSUM of the subgradient of the quantile objective function and requires fitting the model only under the null hypothesis. The critical values can be obtained by simulating the Gaussian process that characterizes the limiting distribution of the test statistic. The proposed method can be used to detect change points at a single quantile level or across multiple quantiles, and can accommodate both homoscedastic and heteroscedastic errors. Simulation study suggests that the proposed method has higher computational efficiency and comparable power with the existing likelihood-ratio-based method in the finite samples. The performance of the proposed method is further illustrated by the analysis of a blood pressure and body mass index data set.

Key words and phrases: Change point, Covariate threshold, Hypothesis testing, Quantile regression, Threshold regression model.

1. Introduction

In regression models, the regression functions are often assumed to take the same parametric form across the entire domain of interest. However, in some applications, regression functions may have different forms in different regions of the conditioning variables. For instance, blood pressure (BP) and body mass index (BMI) showed different relationships for those below and above a BMI threshold (Kaufman et al., 1997; Kerry et al., 2005; Tesfaye et al., 2007); in acute HIV infection, CD4 counts increased rapidly in the first 2-4 weeks but gradually decreased afterwards (Ghosh and Vaidya, 2007); consumer price index showed a significantly positive correlation with the contribution of renewables to energy supply in the higher-economic growth regime, but no significant correlation in the lower-economic growth regime (Chang et al., 2009). The questions in these applications can be addressed by threshold regression models, which specify different regression functions in subsamples segmented by a continuous predictor, referred to as the *threshold variable* thereafter. In other words, in threshold

regression models, the covariate effects are piecewise functions of the threshold variable with jumps occurring at the unknown change points. Therefore, threshold regression models can be viewed as special cases of varying coefficient models (Hastie and Tibshirani, 1993). Threshold regression models have wide applications in biostatistics, epidemiology, economics and finance; see Huang (2009) and Pennell et al. (2010) for additional examples. In the threshold regression literature, most existing work focus on mean regression models (Hansen, 1996). In this paper, we focus on quantile regression.

Quantile regression, pioneered by Koenker and Bassett (1978), is a valuable alternative regression technique to mean regression for modeling the stochastic relationships between random variables. Using quantile regression, we can explore the stochastic relationships in a comprehensive way by studying different tails of the conditional distribution of the response variable. In addition, quantile regression offers an automatic approach for capturing the heteroscedasticity in the population. We refer to Koenker (2005) for a more comprehensive review of quantile regression.

A number of researchers have studied the detection and estimation of change points in quantile regression for time series data where the model structure changes after an unknown time point. Su and Xiao (2008) proposed a sup-Wald test for detecting the structural change of conditional distribution based on sequential quantile regression estimators. Qu (2008) developed two test procedures for detecting change points, one based on a subgradient statistic and the other based on a Wald-type statistic. Oka and Qu (2011) studied the estimation of multiple structural changes in conditional quantile functions. The situations discussed in these papers differ from threshold regression models, where the change point is due to an unknown threshold in the continuous threshold covariate.

There exist very limited work for quantile regression with threshold effects. Caner (2002) studied the least absolute deviation estimators in threshold linear regression models. Kato (2009) developed the extended convexity arguments and applied the theory to establish the asymptotic properties of a proposed Wald-type test for median threshold regression model. The proposed test relies on the identically distributed error assumption and it requires estimating the unknown error density function even for homoscedastic errors. Lee et al. (2011) proposed a sup-likelihood-ratio-based method for testing the existence of threshold effects in regression models, which includes quantile regression as a special case. Li et al. (2011) discussed estimation for a bent line quantile regres-

sion, where a threshold covariate has different effects in two segments separated at an unknown point and the other covariates have constant effects across the entire domain. Therefore, bent line regression can be viewed as a special case of the threshold regression model considered in Lee et al. (2011). Galvao et al. (2011) studied the estimation for threshold quantile autoregressive models and established the asymptotic property for the threshold and regression parameter estimators. Galvao et al. (2013) developed a uniform test of linearity against threshold effects in quantile regression model for stationary time series processes. Cai and Stander (2008) and Cai (2010) discussed forecasting for quantile threshold autoregressive time series models with known threshold values.

In this article, we propose a new procedure for testing the presence of change points due to covariate thresholding in quantile regression. We focus on the general threshold regression models as in Lee et al. (2011), which allow a set of covariates to have heterogeneous effects across the domain of the threshold covariate. Our proposed test statistic is based on a CUSUM process of subgradient obtained by fitting the quantile regression model under the null hypothesis of no threshold effects. We study the asymptotic properties of the proposed test statistics under both null and local alternative models, and develop a convenient procedure for calculating the critical values by simulating the Gaussian process. The proposed test can accommodate both homoscedastic and heteroscedastic errors, and it can be used to test for the presence of change points occurring at either a specified single quantile level or at multiple quantiles. Our numerical studies show that the proposed test tends to have similar power with the likelihood-ratio test of Lee et al. (2011) in finite samples when sizes are controlled at the nominal levels, but the former is less sensitive to the choice of bandwidth. In addition, since it requires only fitting the null model, the proposed method is computationally more efficient than the likelihood-ratio test, which involves grid search over the domain of the threshold covariate for fitting the alternative model.

The rest of the paper is organized as follows. In Section 2, we present the proposed testing procedure at one quantile level together with the asymptotic properties of the proposed test statistic and the calculation of critical values. In Section 3, we discuss the extension of testing change points at multiple quantiles. We assess the finite sample performance of the proposed method through a simulation study in Section 4 and the analysis of a blood pressure and body mass index data set in Section 5. All technical proofs are given in the Supplementary Material.

2. Testing for threshold effects at a single quantile level

2.1. Model setup and testing procedure

At a given quantile level $\tau \in (0, 1)$, we consider the following threshold quantile regression model,

$$Q_Y(\tau|\mathbf{X}, \mathbf{Z}, U) = \mathbf{X}^T \boldsymbol{\beta}_0(\tau) + \mathbf{Z}^T \boldsymbol{\alpha}_0(\tau) I(U > u_0), \quad (2.1)$$

where $Q_Y(\tau|\cdot)$ is the τ th conditional quantile of the response variable Y , \mathbf{X} is a p -dimensional design vector with the first element as 1 corresponding to the intercept, U is the univariate threshold variable that might be an element of \mathbf{X} , \mathbf{Z} is a q -dimensional design vector, $\boldsymbol{\beta}_0(\tau)$ and $\boldsymbol{\alpha}_0(\tau)$ are the unknown regression coefficients, and u_0 is the unknown threshold parameter lying in the region $(0, 1)$. Without loss of generality, we assume that U and \mathbf{Z} are subsets of \mathbf{X} , and $U \in [0, 1]$. Suppose $\{(Y_i, \mathbf{X}_i); i = 1, \dots, n\}$ is a random sample of (Y, \mathbf{X}) .

We are interested in testing the existence of a threshold effect at the quantile level τ . That is, we consider the null and alternative hypotheses

$$H_0 : \boldsymbol{\alpha}_0(\tau) = \mathbf{0} \text{ for any } u_0 \in (0, 1) \text{ v.s. } H_1 : \boldsymbol{\alpha}_0(\tau) \neq \mathbf{0} \text{ for some } u_0 \in (0, 1).$$

Under the null hypothesis, the threshold parameter $\boldsymbol{\alpha}_0(\tau)$ is not identifiable, while $\boldsymbol{\beta}_0(\tau)$ can be estimated by

$$\hat{\boldsymbol{\beta}}(\tau) = \arg \min_{\boldsymbol{\beta}(\tau) \in R^p} \sum_{i=1}^n \rho_\tau\{Y_i - \mathbf{X}_i^T \boldsymbol{\beta}(\tau)\},$$

where $\rho_\tau(r) = r\{\tau - I(r < 0)\}$ is the quantile check function. The building block of our test statistic is

$$\mathbf{R}_n\{u, \tau, \hat{\boldsymbol{\beta}}(\tau)\} = n^{-1/2} \sum_{i=1}^n \psi_\tau\{Y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}}(\tau)\} \mathbf{Z}_i I(U_i \leq u), \quad (2.2)$$

where $\psi_\tau(r) = \tau - I(r < 0)$. To test for the presence of an unknown change point, we search over all possible candidates $u \in (0, 1)$ and define the test statistic as

$$T_n(\tau) = \sup_{u \in (0, 1)} \|\mathbf{R}_n\{u, \tau, \hat{\boldsymbol{\beta}}(\tau)\}\|, \quad (2.3)$$

where $\|\cdot\|$ stands for the L_2 norm.

Note that $\mathbf{R}_n\{u, \tau, \hat{\beta}(\tau)\}$ is the negative subgradient of the quantile objective function corresponding to the subsample with U_i below the threshold u . Therefore, our proposed test statistic can be viewed as a CUSUM statistic based on signs of quantile residuals; see Bai (1996) for a discussion of CUSUM test statistic in least squares regression. Under the null hypothesis, $\hat{\beta}(\tau)$ is a root-n consistent estimator of $\beta_0(\tau)$ and $\mathbf{R}_n\{u, \tau, \hat{\beta}(\tau)\}$ converges to a Gaussian process with mean zero. On the other hand, if there exists a threshold effect, the significant difference of $\hat{\beta}(\tau)$ and $\beta_0(\tau)$ will make the estimated residuals consistently fall below or above zero for a subsample and thus force the statistic $T_n(\tau)$ to take a large value. Our proposed test statistic depends on the score-type statistic, and thus requires fitting the model only under H_0 . This is in contrast with the likelihood ratio approach in Lee et al. (2011), which requires fitting alternative models under all possible partitions of the domain $(0, 1)$ of the threshold variable.

2.2. Asymptotic properties

We study the asymptotic properties of the proposed test statistic under both the null and local alternative models. We first fix some notation. Let $f(\cdot|\mathbf{X})$ and $F(\cdot|\mathbf{X})$ denote the conditional density and distribution functions of Y given \mathbf{X} , respectively. To simplify the notation, we write $f(\cdot|\mathbf{X}_i)$ and $F(\cdot|\mathbf{X}_i)$ as $f_i(\cdot)$ and $F_i(\cdot)$. Also denote $f_U(u)$ and $F_U(u)$ as the density and distribution functions of U , $\mathbf{Z}_u = \mathbf{Z}I(U \leq u)$, $\mathbf{S}_z(u) = E[\mathbf{Z}_u\mathbf{X}^T f\{F^{-1}(\tau|\mathbf{X})|\mathbf{X}\}]$, and $\mathbf{S} = \mathbf{S}_x(1) = E[\mathbf{X}\mathbf{X}^T f\{F^{-1}(\tau|\mathbf{X})|\mathbf{X}\}]$. We make the following assumptions.

- A1 The density $f_i(\cdot)$ is continuous, uniformly bounded away from zero and infinity and has a bounded first derivative in the neighborhood of $F_i^{-1}(\tau)$ for $i = 1, \dots, n$.
- A2 The density function of U , $f_U(u)$, is continuous.
- A3 (a) The design vector satisfies $\max_{1 \leq i \leq n} \|\mathbf{X}_i\| = O_p(n^{1/10}/\log n)$; (b) The matrix \mathbf{S} is positive definite; (c) $E\|\mathbf{X}\|^6$ is bounded.

Assumption A1 imposes some conditions on the smoothness and boundedness of the conditional density functions and their derivatives, which are standard in quantile regression literature. Assumption A2 specifies the condition on the threshold effect covariate. Assumption A3(a) imposes some conditions on the moments of the covariates, which are needed to establish the asymptotic representation of $\hat{\beta}(\tau)$ and the uniform

convergence of the subgradient process under the null and alternative hypotheses. Assumption A3(b) is needed to obtain the asymptotic representation of $\hat{\beta}(\tau)$. Assumption A3(c) is due to Bai (1996) and it ensures the stochastic equicontinuity of the subgradient process $\mathbf{R}_n\{\mathbf{u}, \tau, \beta_0(\tau)\}$.

Theorem 1 *Suppose that Assumptions A1-A3 hold. Under the null hypothesis H_0 , we have*

$$T_n(\tau) \Rightarrow \sup_{u \in (0,1)} \|\mathbf{R}(u)\|, \text{ as } n \rightarrow \infty, \quad (2.4)$$

where “ \Rightarrow ” denotes the weak convergence, $\mathbf{R}(u)$ is the Gaussian process with mean 0 and covariance function $\mathbf{W}(u, u') = \tau(1 - \tau)\{E(\mathbf{Z}_{u'}\mathbf{Z}_u^T) - E(\mathbf{Z}_u\mathbf{X}^T)\mathbf{S}^{-1}\mathbf{S}_z^T(u') - \mathbf{S}_z(u)\mathbf{S}^{-1}E(\mathbf{X}\mathbf{Z}_{u'}^T) + \mathbf{S}_z(u)\mathbf{S}^{-1}E(\mathbf{X}\mathbf{X}^T)\mathbf{S}^{-1}\mathbf{S}_z^T(u')\}$.

We next establish the asymptotic properties of the proposed test statistic under the local alternative model

$$Q_{Y_i}(\tau|\mathbf{X}_i) = \mathbf{X}_i^T \beta_0(\tau) + n^{-1/2} \mathbf{Z}_i^T \alpha_0(\tau) I(U_i > u_0), i = 1, \dots, n, \quad (2.5)$$

where $u_0 \in (0, 1)$ is the change point and $\alpha_0(\tau) \neq 0$ is a q -dimensional vector.

Theorem 2 *Suppose that Assumptions A1-A3 hold. Under the local alternative model (2.5), we have*

$$T_n(\tau) \Rightarrow \sup_{u \in (0,1)} \|\mathbf{R}(u) + \mathbf{q}\{u, \alpha_0(\tau)\}\|, \text{ as } n \rightarrow \infty, \quad (2.6)$$

where $\mathbf{R}(u)$ is the same Gaussian process as defined in Theorem 1, and $\mathbf{q}\{u, \alpha_0(\tau)\} = -\mathbf{S}_z(u)\mathbf{S}^{-1}\mathbf{Q}\{\alpha_0(\tau)\} + \mathbf{P}\{u, \alpha_0(\tau)\}$, where $\mathbf{Q}(\alpha) = E[\mathbf{X}\mathbf{Z}^T \alpha I(U > u_0) f\{F^{-1}(\tau|\mathbf{X})|\mathbf{X}\}]$ and $\mathbf{P}(u, \alpha) = E[\mathbf{Z}\mathbf{Z}^T \alpha I(u_0 < U \leq u) f\{F^{-1}(\tau|\mathbf{X})|\mathbf{X}\}]$.

Note that $\mathbf{q}\{u, \alpha_0(\tau)\} = 0$ for all u under H_0 , and it is a nonzero function of u under the local alternative. Therefore, the proposed test statistic can be used to distinguish the alternative with a change point due to a threshold effect from the null hypothesis of no change point. The following Corollary 1 shows that the power of the test $T_n(\tau)$ approaches to one when the order of the threshold effect under the alternative model is arbitrarily close to $n^{-1/2}$.

Corollary 1 *Suppose that Assumptions A1-A3 hold. Under the local alternative model $Q_{Y_i}(\tau|\mathbf{X}_i) = \mathbf{X}_i^T \boldsymbol{\beta}_0(\tau) + n^{-1/2} a_n \mathbf{Z}_i^T \boldsymbol{\alpha}_0(\tau) I(U_i > u_0)$ with $u_0 \in (0, 1)$, if $a_n \rightarrow \infty$, then we have $\lim_{n \rightarrow \infty} P\{\|T_n(\tau)\| \geq t\} = 1$ for all $t > 0$.*

2.3. Calculation of critical values

Theorem 1 suggests that the asymptotic null distribution of $T_n(\tau)$ is nonstandard, and it depends on the unknown density function for heteroscedastic errors. Therefore, critical values for $T_n(\tau)$ can not be tabulated for general cases. We overcome this difficulty by simulating the asymptotic representation of $\mathbf{R}_n\{u, \tau, \hat{\boldsymbol{\beta}}(\tau)\}$.

Let $\{e_i; i = 1, \dots, n\}$ be a random sample with τ th quantile zero and $\{\omega_i; i = 1, \dots, n\}$ be a random sample independent of e_i with zero mean, unit variance and a finite third moment. Define

$$\mathbf{R}_n^*(u) = n^{-1/2} \sum_{i=1}^n \omega_i \psi_\tau(e_i) \{I(U_i \leq u) \mathbf{Z}_i - \mathbf{S}_{z,n}(u) \mathbf{S}_n^{-1} \mathbf{X}_i\}, \quad (2.7)$$

where $\mathbf{S}_{z,n}(u) = n^{-1} \sum_{i=1}^n \mathbf{Z}_i \mathbf{X}_i^T I(U_i \leq u) K_{h_n}\{Y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}}(\tau)\}$, $\mathbf{S}_n = n^{-1} \sum_{i=1}^n \mathbf{X}_i \mathbf{X}_i^T K_{h_n}\{Y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}}(\tau)\}$, $K_{h_n}(\cdot) = h_n^{-1} K(\cdot/h_n)$, $K(\cdot)$ is a kernel function, and h_n is a positive bandwidth. Throughout our numerical studies, we follow the suggestion in Lee et al. (2011) and use Silverman's rule of thumb (Silverman 1986), $h_n = 1.06 \hat{\sigma} n^{-1/5}$, where $\hat{\sigma}$ is the sample standard deviation of $\{Y_i - \mathbf{X}_i^T \hat{\boldsymbol{\beta}}(\tau); i = 1, \dots, n\}$. The sensitivity analysis in Section 3 shows that the performance of the proposed test procedure is stable with $h_n = c \hat{\sigma} n^{-1/5}$ for $c \in [0.1, 3]$.

Before presenting the simulation procedure, we first study the asymptotic property of $\mathbf{R}_n^*(u)$. We assume the following assumptions on the kernel function and the bandwidth.

- A4 The function $K(\cdot)$ is a symmetric kernel function with compact support, and it satisfies $\int K(u) du = 1$ and has a bounded first derivative.
- A5 The positive bandwidth h_n satisfies $h_n \rightarrow 0$ and $h_n n^{1/5} \sqrt{\log n} \rightarrow \infty$ as $n \rightarrow \infty$.

Theorem 3 *Suppose that Assumptions A1-A5 hold. Under H_0 , $\mathbf{R}_n^*(u)$ converges to the Gaussian process $\mathbf{R}(u)$ defined in Theorem 1 as $n \rightarrow \infty$.*

Theorem 3 suggests that we can calculate the critical values by the quantiles of $\max_u \|\mathbf{R}_n^*(u)\|$. The detailed computing procedure is as follows:

Step 1. Generate $\{e_i; i = 1, \dots, n\}$ as a random sample with the τ th quantile zero and $\{\omega_i, i = 1, \dots, n\}$ as a random sample (independent of e_i) with mean 0, variance 1 and a finite third moment. In our implementation, we generate e_i independently from $N(0, 1) - \Phi^{-1}(\tau)$ with Φ being the cumulative distribution function of $N(0, 1)$, and ω_i from the two-point mass distribution with equal probability at 1 and -1.

Step 2. Simulate the process

$$\mathbf{R}_n^*(u) = n^{-1/2} \sum_{i=1}^n \omega_i \psi_\tau(e_i) \{I(U_i \leq u) \mathbf{Z}_i - \mathbf{S}_{z,n}(u) \mathbf{S}_n^{-1} \mathbf{X}_i\},$$

and obtain the test statistic $T_n^*(\tau) = \sup_{u \in (0,1)} \|\mathbf{R}_n^*(u)\|$.

Step 3. Repeat Steps 1–2 J times to get $T_{n1}^*(\tau), \dots, T_{nJ}^*(\tau)$. Calculate the critical value for a level α test by the $(1 - \alpha)$ th sample quantile of $\{T_{nj}^*(\tau); j = 1, \dots, J\}$.

Remark 1 For model (2.5), let $\epsilon = Y - \mathbf{X}^T \beta_0(\tau) - n^{-1/2} \mathbf{Z}^T \alpha_0(\tau) I(U > u_0)$ denote the error term whose τ th quantile is zero conditional on \mathbf{X} . Let $f_\epsilon(\cdot | \mathbf{X})$ be the conditional density function of ϵ given \mathbf{X} , then we have $f\{F^{-1}(\tau | \mathbf{X}) | \mathbf{X}\} = f_\epsilon(0 | \mathbf{X})$. Hence for model (2.5) with i.i.d. (independent and identically distributed) errors, the density functions in $\mathbf{S}(u)$ and \mathbf{S}^{-1} are cancelled out and thus do not need to be estimated. Therefore, the process $\mathbf{R}_n^*(u)$ can be simplified as

$$\mathbf{R}_n^*(u) = n^{-1/2} \sum_{i=1}^n \omega_i \psi_\tau(e_i) \{I(U_i \leq u) \mathbf{Z}_i - \mathbf{S}_{z,1n}(u) \mathbf{S}_{1n}^{-1} \mathbf{X}_i\}, \quad (2.8)$$

where $\mathbf{S}_{z,1n}(u) = n^{-1} \sum_{i=1}^n \mathbf{Z}_i \mathbf{X}_i^T I(U_i \leq u)$ and $\mathbf{S}_{1n} = n^{-1} \sum_{i=1}^n \mathbf{X}_i \mathbf{X}_i^T$.

2.4. Comparison to the sup-likelihood-ratio-type test of Lee et al. (2011)

Lee et al. (2011) developed a general sup-likelihood-ratio-type (LRT) test for testing threshold effects in regression models including quantile regression. We carry out a systematic comparison of our proposed sup-score-type (SS) test with the LRT test.

The LRT test statistic in Lee et al. (2011) is defined as

$$LRT_n = \sup_{u \in (0,1)} n \{Q_n(u) - \tilde{Q}_n\}, \quad (2.9)$$

where $\tilde{Q}_n = \min_{\beta(\tau)} n^{-1} \sum_{i=1}^n \rho_\tau\{Y_i - \mathbf{X}_i^T \beta(\tau)\}$ and $Q_n(u) = \min_{\beta(\tau), \alpha(\tau)} n^{-1} \sum_{i=1}^n \rho_\tau\{Y_i - \mathbf{X}_i^T \beta(\tau) - \mathbf{Z}_i^T \alpha(\tau) I(U_i > u)\}$ are the minimum values of the quantile objective

functions under the null and the alternative hypothesis with the change point $u_0 = u$, respectively.

Note that the proposed SS method is based on the score-type test statistic obtained by only fitting the null model, while the LRT method is based on the pseudo-likelihood-ratio-type test statistic and requires fitting both the null and alternative models.

The following Proposition 1 gives the asymptotic distributions of LRT_n under the null and local alternative models, where (i) comes directly from (3.6) of Lee et al. (2011), and the proof of (ii) is provided in the Supplementary Material.

Proposition 1 (i) Under Assumptions A1–A3 and H_0 , $LRT_n \Rightarrow \frac{1}{2} \{ \sup_{u \in (0,1)} \mathcal{G}(u)^T \mathbf{V}(u)^{-1} \mathcal{G}(u) - \mathcal{G}_1^T \mathbf{V}_1^{-1} \mathcal{G}_1 \}$ as $n \rightarrow \infty$, where $\mathcal{G}(u)$ is a mean-zero Gaussian process with covariance kernel $\widetilde{\mathbf{W}}(u, u') = \tau(1 - \tau) E(\widetilde{\mathbf{X}}_u \widetilde{\mathbf{X}}_{u'}^T)$ with $\widetilde{\mathbf{X}}_u = (\mathbf{X}^T, \mathbf{Z}^T I(U > u))^T$ being the vector of covariates under the alternative hypothesis, $\mathbf{V}(u) = E[\widetilde{\mathbf{X}}_u \widetilde{\mathbf{X}}_u^T f\{\mathbf{X}^T \boldsymbol{\beta}_0(\tau) | \mathbf{X}\}]$, \mathcal{G}_1 and \mathbf{V}_1 denote the first p elements of \mathcal{G} and the first $p \times p$ block of $\mathbf{V}(u)$. (ii) Under Assumptions A1–A5 and the local alternative model (2.5), $LRT_n \Rightarrow \frac{1}{2} (\sup_{u \in (0,1)} [\mathcal{G}(u) + \widetilde{\mathbf{q}}_L\{u, \boldsymbol{\alpha}_0(\tau)\}]^T \mathbf{V}(u)^{-1} [\mathcal{G}(u) + \widetilde{\mathbf{q}}_L\{u, \boldsymbol{\alpha}_0(\tau)\}] - (\mathcal{G}_1 + \widetilde{\mathbf{q}}_1)^T \mathbf{V}_1^{-1} (\mathcal{G}_1 + \widetilde{\mathbf{q}}_1))$ as $n \rightarrow \infty$, where $\widetilde{\mathbf{q}}_L\{u, \boldsymbol{\alpha}_0(\tau)\} = (E[\mathbf{XZ}^T \boldsymbol{\alpha}_0(\tau) I(U > u_0)] f\{\mathbf{X}^T \boldsymbol{\beta}_0(\tau) | \mathbf{X}\})^T$, $E[\mathbf{ZZ}^T \boldsymbol{\alpha}_0(\tau) I\{U > \max(u, u_0)\}] f\{\mathbf{X}^T \boldsymbol{\beta}_0(\tau) | \mathbf{X}\})^T$ and $\widetilde{\mathbf{q}}_1 = E[\mathbf{XZ}^T \boldsymbol{\alpha}_0(\tau) I(U > u_0)] f\{\mathbf{X}^T \boldsymbol{\beta}_0(\tau) | \mathbf{X}\}$.

The critical value can be obtained by simulating the asymptotic distribution of LRT_n under the null hypothesis; see Section 3.2 of Lee et al. (2011) for details. The limiting distribution of the LRT test statistic has a nonstandard form different from that of the SS test statistic, and this makes it difficult to compare two statistics explicitly. However, we can compare the asymptotic local powers of two methods analytically by simulating the limiting distributions of $T_n(\tau)$ and LRT_n . Our analysis in Section 4.2 shows that two methods have similar asymptotic local power for various designs considered.

3. Testing for threshold effects at multiple quantiles

In many cases, different magnitudes of changes may occur at different quantiles. Therefore, it will be more informative to incorporate multiple quantiles instead of only one quantile level to identify the threshold effects. We use $\mathcal{T} = [\omega_1, \omega_2]$ to represent a closed set of quantiles of interest. We focus on the following null and alternative

hypotheses:

$$H_0^* : \alpha_0(\tau) = 0 \text{ for any } u_0 \in (0, 1) \text{ and for all } \tau \in \mathcal{T},$$

against

$$H_1^* : \alpha_0(\tau) \neq 0 \text{ for some } u_0 \in (0, 1) \text{ and some } \tau \in \mathcal{T},$$

which imply that there is no threshold effect for any quantile in the set \mathcal{T} under the null while there exist threshold effects at some quantiles under the alternative. Similar as the method proposed in Section 2.1, we consider the following test statistic

$$T_n = \sup_{\tau \in \mathcal{T}} \sup_{u \in (0,1)} \|R_n\{u, \tau, \hat{\beta}(\tau)\}\|,$$

where $\hat{\beta}(\tau)$ is the coefficient estimate at the τ th quantile obtained under the null hypothesis. A large T_n indicates that there likely exist threshold effects for some $\tau \in \mathcal{T}$.

To establish the limiting distribution of T_n , we strength the condition A1 as follow.

A6 Assumption A1 holds uniformly in $\tau \in \mathcal{T}$.

Theorem 4 *Suppose that Assumptions A2-A6 hold. Under the null hypothesis H_0^* , we have*

$$T_n \Rightarrow \sup_{\tau \in \mathcal{T}} \sup_{u \in (0,1)} \|\mathbf{R}(u, \tau)\|, \text{ as } n \rightarrow \infty,$$

where $\mathbf{R}(u, \tau)$ is a q -vector of independent Gaussian processes with zero mean and covariance function

$$\begin{aligned} E\{\mathbf{R}(u, \tau)\mathbf{R}(u', \tau')\} &= \{\min(\tau, \tau') - \tau\tau'\} \{E(\mathbf{Z}_u\mathbf{Z}_{u'}) - E(\mathbf{Z}_u\mathbf{X}^T)\mathbf{S}^{-1}\mathbf{S}_z^T(u') \\ &\quad - \mathbf{S}_z(u)\mathbf{S}^{-1}E(\mathbf{X}\mathbf{Z}_{u'}^T) + \mathbf{S}_z(u)\mathbf{S}^{-1}E(\mathbf{X}\mathbf{X}^T)\mathbf{S}^{-1}\mathbf{S}_z^T(u')\}. \end{aligned} \quad (3.1)$$

Suppose the local alternative model (2.5) holds for all quantiles $\tau \in \mathcal{T}$, then

$$T_n \Rightarrow \sup_{\tau \in \mathcal{T}} \sup_{u \in (0,1)} \|\mathbf{R}(u, \tau) + \mathbf{q}\{u, \alpha_0(\tau)\}\|, \text{ as } n \rightarrow \infty,$$

where $\mathbf{q}\{u, \alpha_0(\tau)\}$ is the same as that defined in Theorem 2.

To obtain critical values of the test statistic across quantiles, the simulation method described in Section 2.3 can be slightly modified as follows. Let τ_1, \dots, τ_m be a grid

of quantile levels from \mathcal{T} . For the simulation method, we can generate the subgradient process as

$$\mathbf{R}_{nk}^*\{u, \tau_k, \boldsymbol{\beta}(\tau_k)\} = n^{-1/2} \sum_{i=1}^n \omega_i \psi_{\tau_k}\{e_i - Q_e(\tau_k)\} \{I(U_i \leq u) \mathbf{Z}_i - \mathbf{S}_{\mathbf{z},n}(u) \mathbf{S}_n^{-1} \mathbf{X}_i\},$$

for $k = 1, \dots, m$ and calculate $T_n^* = \max_{k=1, \dots, m} \sup_{u \in (0,1)} \|\mathbf{R}_{nk}^*\{u, \tau_k, \boldsymbol{\beta}(\tau_k)\}\|$, where e_i are *i.i.d.* random variables with the τ th quantile $Q_e(\tau)$, and ω_i are *i.i.d.* random variables independent of e_i with zero mean, unit variance and finite third moment.

4. Simulation Study

4.1. Setup

The simulated data sets are generated from the following model:

$$Y_i = 1 + X_i + U_i - \alpha_0 X_i I(U_i > 0.5) + \sigma(X_i, U_i) \epsilon_i, \quad (4.1)$$

where $X_i \sim Uniform(-2, 2)$, $U_i \sim Uniform(0, 1)$, ϵ_i are independent and identically distributed random errors, $\sigma(X_i, U_i)$ measures the heteroscedasticity, and α_0 controls the degree of departure from the null hypothesis with $\alpha_0 = 0$ representing no threshold effects in the model. We consider four different cases. In Case 1, $\sigma(X_i, U_i) = 1$ corresponds to a homoscedastic model. Cases 2-4 have heteroscedastic errors with $\sigma(X_i, U_i) = 1 + 0.3X_i$ in Cases 2 and 4, and $\sigma(X_i, U_i) = 1 + 0.3X_i + 0.3U_i$ in Case 3. The random errors ϵ_i are generated from $N(0, 1)$ in Cases 1-3 and from t_4 distribution in Case 4. The sample size is chosen to be $n = 200$ and 500 . For all scenarios, the simulation is repeated 500 times.

We consider two variations of the proposed testing procedure, *SIID* and *SNID*, for which the critical values are obtained through simulation by assuming *i.i.d.* and non-*i.i.d.* errors, respectively. For comparison, we include two variations of the sup-likelihood-ratio-type method, *LIID* and *LNID* for *i.i.d.* and non-*i.i.d.* errors, respectively. Different from *SIID*, the density function in *LIID* can not be cancelled out and thus the bandwidth parameter is still involved in the kernel estimation; see (3.8) and (3.9) in Lee et al. (2011) for more details. For all the methods considered, the lower and upper 10% quantiles of U are trimmed when maximizing over U , and we use the Gaussian kernel function for the *SNID*, *LIID* and *LNID* methods.

4.2. Type I error and sensitivity analysis

Table 4.1 summarizes the Type I errors of four testing procedures in Cases 1-4 with $n = 200$ and 500 at three quantile levels $\tau = 0.1, 0.5$ and 0.9 . The nominal significance level is set as 5% . For *SIID* and *SNID*, we also report the Type I errors for detecting the threshold effects at three quantiles jointly (referred to as “Multiple” in the table). The score-based methods *SIID* and *SNID* give slightly inflated Type I errors for the smaller sample size $n = 200$. However, for $n = 500$, the *SNID* method maintains the Type I errors close to the nominal level for all scenarios considered. Even though the *SIID* method assumes *i.i.d.* errors, the method is quite robust against the violation of this assumption. For $n = 500$, the Type I errors of *SIID* are close to the nominal level in Case 1 with homoscedastic errors, and they are also reasonable (slightly inflated for a few scenarios) in Cases 2-4 with heteroscedastic errors. Across all scenarios considered, the sup-likelihood-ratio-based methods *LIID* and *LNID* give deflated Type I errors for both $n = 200$ and $n = 500$, and this over-conservativeness was also observed in the Supplementary Material of Lee et al. (2011, page 19).

To assess the sensitivity of *SNID*, *LIID* and *LNID* to the bandwidth, we let $h_n = cn^{-1/5}\hat{\sigma}$ and plot the Type I errors of three methods against $c \in [0.1, 3]$ in Cases 1-4 at the single quantile level $\tau = 0.5$. The two LRT methods *LIID* and *LNID* are sensitive to the choice of the bandwidth. For all scenarios considered, the Type I errors of *LIID* fall below the nominal level across the entire region of c considered. The Type I errors of *LNID* are generally close to the nominal level for smaller c but they become deflated as c increases. Note that the acceptable region of c for the LRT methods appears to depend on n , the distribution and the quantile level of study. Therefore, in practice it would be difficult to find a constant c that works universally well for the LRT methods. The score-based test *SNID* is more robust against the choice of h_n , and it gives reasonable Type I errors across a wide range of c , especially for larger sample size $n = 500$.

4.3. Power analysis

For power analysis, we consider the local alternative model (4.1) with $\alpha_0 = n^{-1/2}\delta$, where $\delta = 1, 2, \dots, 9$. We showed in Section 4.2 that the LRT methods tend to have deflated Type I errors. For fairness of the power comparison, for both SS and LRT methods, we calculate the critical values as the 95th percentiles of the corresponding test statistics across 500 simulated data generated from the null hypothesis. This ensures that both SS and LRT tests have exactly 5% Type I error, but note that such adjustment is not feasible in real applications. For comparison, we also include the asymptotic




Figure 4.1: Type I errors of *SNID*, *LIID* and *LNID* at $\tau = 0.5$ against c , the constant involved in the bandwidth $h_n = cn^{-1/5}\hat{\sigma}$. The dashed horizontal line corresponds to the nominal level of 5%, and the shade area corresponds to the 95% confidence interval of Type I error of valid tests of size 5%. The solid, dashed and dotted lines represent Type I errors from the *SNID*, *LNID* and *LIID* methods, respectively.

local powers of the SS and LRT methods, which are obtained by simulating the limiting distributions of $T_n(\tau)$ and LRT_n using the results in Theorem 2 and Proposition 1, respectively.

Figure 4.2 plots the asymptotic and finite-sample Monte Carlo local powers of SS

Table 4.1: Type I errors of different testing procedures in Cases 1-4.

Sample size	τ	Methods	Case 1	Case 2	Case 3	Case 4	
n=200	$\tau = 0.1$	<i>SIID</i>	0.052	0.058	0.054	0.062	
		<i>SNID</i>	0.040	0.038	0.020	0.046	
		<i>LIID</i>	0.030	0.022	0.024	0.024	
		<i>LNID</i>	0.014	0.016	0.012	0.010	
	$\tau = 0.5$	<i>SIID</i>	0.082	0.086	0.082	0.074	
		<i>SNID</i>	0.068	0.072	0.070	0.058	
		<i>LIID</i>	0.016	0.022	0.018	0.022	
		<i>LNID</i>	0.008	0.014	0.006	0.006	
	$\tau = 0.9$	<i>SIID</i>	0.058	0.078	0.070	0.098	
		<i>SNID</i>	0.048	0.054	0.052	0.056	
		<i>LIID</i>	0.018	0.024	0.030	0.038	
		<i>LNID</i>	0.004	0.014	0.012	0.016	
	Multiple	<i>SIID</i>	0.084	0.084	0.084	0.076	
		<i>SNID</i>	0.068	0.072	0.080	0.056	
	n=500	$\tau = 0.1$	<i>SIID</i>	0.038	0.060	0.060	0.070
			<i>SNID</i>	0.036	0.046	0.038	0.038
<i>LIID</i>			0.018	0.022	0.020	0.028	
<i>LNID</i>			0.012	0.012	0.008	0.022	
$\tau = 0.5$		<i>SIID</i>	0.042	0.064	0.056	0.062	
		<i>SNID</i>	0.048	0.056	0.046	0.056	
		<i>LIID</i>	0.030	0.026	0.030	0.034	
		<i>LNID</i>	0.028	0.014	0.016	0.012	
$\tau = 0.9$		<i>SIID</i>	0.052	0.050	0.064	0.074	
		<i>SNID</i>	0.046	0.038	0.054	0.044	
		<i>LIID</i>	0.032	0.022	0.032	0.032	
		<i>LNID</i>	0.018	0.016	0.012	0.024	
Multiple		<i>SIID</i>	0.048	0.060	0.054	0.074	
		<i>SNID</i>	0.042	0.038	0.036	0.052	

and LRT tests across δ at $\tau = 0.5$, referred to as ASLP and MCLP, respectively. The finite-sample local powers are obtained through Monte Carlo simulation by running the test procedures for data generated from model (4.1) with $\alpha_0 = n^{-1/2}\delta$ and $n = 2000$. The results show that as δ increases, the local powers of both SS and LRT methods increase gradually to one. The MCLP and ASLP agree generally well with each other, and this validates the theoretical results. In addition, the asymptotic local powers of the SS and LRT methods appear close to each other for all four cases considered.

4.4. Comparison of computational efficiency

To compare the computational efficiency of the SS and LRT methods, we report in Table 4.2 the average computing time (in seconds) of each test procedure for analyzing one simulated data set at median. The *SIID* tends to be faster than *SNID* as it does not require estimating the unknown density function. Since SS test requires fitting the model only under null hypothesis, both *SIID* and *SNID* take less time than the LRT methods, and this computational efficiency is more obvious for larger samples.

In summary, the proposed SS method has similar power with the LRT method of Lee et al. (2011), when the Type I errors are controlled at the same level. However, in finite samples, the LRT method is sensitive to the choice of bandwidth, making it more difficult to maintain the Type I error close to the nominal level. In addition, the SS method is computationally more efficient than the LRT method, which requires fitting the model under both null and alternative hypotheses.

5. Blood Pressure and Body Mass Index Study

The relationship between blood pressure and body mass index has long been an important topic in public health studies. Many studies have reported positive associations between BP and BMI in different populations (He et al., 1994; Droyvold et al., 2005; Tesfaye et al., 2007). However, some researchers showed that a linear relationship may not hold across the entire BMI range (Bunker et al., 1995; Kaufman et al., 1997; Kerry et al., 2005).

We use the data from National Health and Nutrition Examination Survey (NHANES) to illustrate the performance of our proposed method. The NHANES program started in the early 1960s and is designed to assess the health and nutritional status of adults and children in the United States. We aim to study the relationship between systolic blood pressure (SBP) and BMI at different quantiles after accounting for gender and age ef-




Figure 4.2: The asymptotic local power (ASLP) and Monte Carlo local power (MCLP) obtained by simulation (500 repetitions) with $n = 2000$ from the SS and LRT methods against δ for the local alternative model with $\alpha_0 = n^{-1/2}\delta$ at $\tau = 0.5$. The lines with circles, squares, crosses and triangles represent $ASLP(LRT)$, $MCLP(LRT)$, $ASLP(SS)$, and $MCLP(SS)$ methods, respectively.

fects, and examine if the relationship is stable across the range of BMI. In this study, the analysis at high quantiles is of great interest since high blood pressure is known to be an important potential cause of heart and vascular diseases. We consider the survey data of non-hispanic black people collected in year 2009-2010, including 683 males and 695 females. For easier demonstration, we analyze data for females and males separately.

First we carry out hypothesis tests to assess if BMI has any threshold effects on the quantiles of BP. In the presence of threshold effects, we have the following quantile regression model

$$Q_{Y_i}(\tau|X_i, Z_i) = \begin{cases} a_1(\tau) + b_1(\tau)X_i + c_1(\tau)Z_i, & X_i \leq u_0(\tau), \\ a_2(\tau) + b_2(\tau)X_i + c_2(\tau)Z_i, & X_i > u_0(\tau), \end{cases} \quad (5.1)$$

where Y_i , X_i and Z_i represent the systolic blood pressure, BMI and age of the i th subject, respectively, and $u_0(\tau)$ is the unknown change point associated with the τ th quantile. We consider three quantile levels $\tau = 0.1, 0.5$ and 0.9 .

Table 5.3 summarizes the testing results from the proposed methods *SIID* and *SNID*, and the sup-likelihood-ratio-based methods *LIID* and *LNID* for males and females separately. For females, except the *LIID* method at $\tau = 0.5$, all four methods suggest that BMI has no significant threshold effect at three quantiles. For males, *SIID*, *SNID* and *LIID* suggest that BMI has significant threshold effects at three quantiles, but the *LNID* method only detects the threshold effect at $\tau = 0.1$ at the significance level of

Table 4.2: The average computing time (in seconds) of different testing procedures for analyzing one simulated data set at median.

	<i>SIID</i>	<i>SNID</i>	<i>LIID</i>	<i>LNID</i>
$n = 200$				
Case 1	0.60	16.45	33.28	36.38
Case 2	0.58	17.13	30.26	34.58
Case 3	0.54	17.45	30.21	34.16
Case 4	0.56	17.54	40.64	31.60
$n = 500$				
Case 1	1.51	21.65	115.31	159.57
Case 2	1.47	20.75	97.18	105.36
Case 3	1.34	21.27	94.66	110.42
Case 4	1.40	21.83	122.95	112.83

0.05. This agrees with the observation in the simulation study that the LRT methods tend to be more conservative than the proposed sup-score-based methods. In contrast, the likelihood-ratio test in Hansen (1999) for mean regression gives p -values 0.001 and 0.014 for the male and female groups, respectively. In the following, we focus our analysis on the male group.

Table 5.3: P -values from testing the existence of threshold effects based on different test procedures in the blood pressure and body mass index study. The last row provides the p -values from the joint testing at three quantile levels.

Quantile	SIID	SNID	LIID	LNID
Male				
$\tau = 0.1$	0.002	0.032	0.000	0.010
$\tau = 0.5$	0.008	0.044	0.020	0.064
$\tau = 0.9$	0.016	0.016	0.012	0.208
Multiple	0.002	0.044		
Female				
$\tau = 0.1$	0.434	0.206	0.546	0.686
$\tau = 0.5$	0.430	0.166	0.028	0.230
$\tau = 0.9$	0.186	0.098	0.094	0.912
Multiple	0.402	0.140		

Table 5.4: Estimation results in the study of BP and BMI association for the male group. The values in parentheses correspond to the bootstrap standard errors.

		First Segment			Second Segment			Change Point
		a_1	b_1	c_1	a_2	b_2	c_2	
$\tau = 0.1$	Est	64.291	1.310	0.341	113.105	-0.346	0.122	24.6
	SE	(8.570)	(0.474)	(0.063)	(4.935)	(0.164)	(0.056)	
$\tau = 0.5$	Est	62.433	1.855	0.590	102.842	0.157	0.393	21.7
	SE	(8.150)	(0.485)	(0.138)	(4.687)	(0.158)	(0.044)	
$\tau = 0.9$	Est	104.977	-0.086	1.129	115.623	0.168	0.589	22.4
	SE	(19.710)	(0.966)	(0.167)	(8.210)	(0.237)	(0.060)	
Mean	Est	74.958	1.126	0.667	107.728	0.060	0.377	22.4
	SE	(11.774)	(0.622)	(0.062)	(3.499)	(0.105)	(0.036)	

Note: Est and SE stand for Parameter estimate and Standard error, respectively.

For quantile regression model (5.1) with a threshold effect, let $\theta(\tau)$ be the collection of regression coefficients $\{a_k(\tau), b_k(\tau), c_k(\tau); k = 1, 2\}$ and the change point $u_0(\tau)$. Following Li et al. (2011), we can estimate $\theta(\tau)$ by

$$\hat{\theta}(\tau) = \arg \min \sum_{i=1}^n \rho_{\tau}\{Y_i - Q_{Y_i}(\tau|X_i, Z_i)\}. \quad (5.2)$$

We summarize the estimated parameters and corresponding standard errors (obtained by the paired bootstrap method with 200 bootstrap repetitions. in Table 5.4, and plot the estimated quantiles and mean functions of BP against BMI conditional on the average age in Figure 5.3. The estimated mean and median regression functions appear consistent with each other. The BMI cut point is around 22 kg/m^2 for both median and the upper quantile $\tau = 0.9$, while it is larger (24.6 kg/m^2) at the lower quantile $\tau = 0.1$, that is, for those with low blood pressures. Not surprisingly, age exhibits significantly positive effects on BP at all three quantiles. However, the effects of age become smaller for those males with BMI above the cut points. In addition, age tends to have larger effects on the upper quantile of the BP distribution. At lower and central quantiles, the effects of BMI are significantly positive before the change point, then become insignificant after the change point. At the upper quantile $\tau = 0.9$, the effects of BMI are not significant throughout the BMI range.

The consistency of BMI effects on BP has been examined in many medical studies for different populations. Past work suggested that the relationship between BMI and BP is likely nonlinear but this may vary between subgroups. By studying urban Nigerian civil servants in 1992, Bunker et al. (1995) suggested that BMI and BP were not correlated below the BMI threshold of 21.5 kg/m^2 but correlated above the threshold. In a study of low-BMI populations in Africa and the Caribbean, Kaufman et al. (1997) observed a threshold at 21 kg/m^2 in the relationship between BMI and BP for women but not for men. Kerry et al. (2005) found that for lean older and semi-urban women in West Africa, the effect of BMI on BP below the change point was greater than the effect above it.

Our analysis for US non-hispanic black subjects also suggests different BMI and BP relationships for different subgroups. The previous work all focus on mean regression analysis. Our proposed methods offer an alternative way to test for the existence of the BMI threshold effect at different tails of the BP distribution, and thus can help assess the BP and BMI association from more angles. For instance, our analysis suggests that at the




Figure 5.3: *Estimated conditional quantile functions of blood pressure against body mass index for the male group at the mean age.*

upper tails of the BP distribution, BP has weaker associations with BMI while stronger associations with age. Further investigation is needed to understand the reasons for this phenomenon and the health implications in obesity epidemic.

Acknowledgement

The authors would like to thank the Editor, the Associate Editor, and three anonymous reviewers for their thoughtful and constructive comments that have significantly improved the paper. Zhang's research was supported by NNSFC Grant 11171074. Dr. Wang's research was supported by NSF (National Science Foundation) CAREER Award DMS-1149355. Dr. Zhu's research is partially supported by NSFC 11271080.

Supplementary Material

Two remarks and all the technical proofs are provided in the Supplementary Material.

References

- Bai, J. (1996). Testing for parameter constancy in linear regressions: an empirical distribution function approach. *Econometrica* **64**, 597-622.
- Bunker, C. H., Ukoli, F. A., Matthews, K. A., Kriska, A. M., Huston, S. L. and Kuller, L. H. (1995). Weight threshold and blood pressure in a lean black population. *Hypertension* **26**, 616-623.
- Cai, Y. (2010). Forecasting for quantile self-exciting threshold autoregressive time series models. *Biometrika* **97**, 199-208.
- Cai, Y. and Stander, J. (2008). Quantile self-exciting threshold autoregressive time series models. *J. Time Ser. Anal.* **29**, 186-202.
- Caner, M. (2002). A note on least absolute deviation estimation of a threshold model. *Econometric Theory* **18**, 800-814.
- Chang, T. H., Huang, C. M. and Lee, M. C. (2009). Threshold effect of the economic growth rate on the renewable energy development from a change in energy price: evidence from oecd countries. *Energy Policy* **37**, 5796-5802.
- Droyvold, W., Midthjell, K., Nilsen, T. and Holmen, J. (2005). Change in body mass index and its impact on blood pressure: a prospective population study. *Int J. Obes (Lond)*, **29**, 650-655.
- Galvao, A.F., Montes-Rojas, G. and Olmo, J. (2011). Threshold quantile autoregressive models. *J. Time Ser. Anal.* **32**:253-267.
- Galvao A.F., Kato, K., Montes-Rojas, G. and Olmo, J. (2013). Testing linearity against threshold effects: uniform inference in quantile regression, *Ann. Inst. Stat. Math.* Forthcoming.
- Ghosh, P. and Vaida, F. (2007). Random changepoint modeling of HIV immunologic responses. *Stat. Med* **26**, 2074-2087.

- Hansen, B. E. (1996). Inference when a nuisance parameter is not identified under the null hypothesis. *Econometrica* **64**, 413-430.
- Hastie, T. and Tibshirani, R. (1993). Varying-coefficient models. *J. Roy. Statist. Soc. B* **55**, 757-796.
- He, J., Klag, M. J., Whelton, P. K., Chen, J. Y., Qian, M. C. and He, G. (1994). Body mass and blood pressure in a lean population in southwestern China. *Am J. Epidemiol* **139**, 380-389.
- Huang, S.C. (2009). Threshold and leverage effects of major asian stock markets based on stochastic volatility models. *International Research J. Finance Econ.* **27**, 106-116.
- Kato, K. (2009), Asymptotics for argmin processes: convexity arguments, *J. Multivariate Anal.*, **100**, 1816-1829.
- Kaufman, J., Asuzu, M., Mufunda, J., Forrester, T., Wilks, R., Luke, A., Long, A.E. and Cooper, R. S. (1997). Relationship between blood pressure and body mass index in lean populations. *Hypertension* **30**, 1511-1516.
- Kerry, S., Micah, F., Plange-Rhule, J., Eastwood, J. and Cappuccio, F. (2005). Blood pressure and body mass index in lean rural and semi-urban subjects in west Africa. *J. Hypertens* **23**, 1645-1651.
- Koenker, R. (2005). *Quantile regression*. Cambridge University Press, Cambridge.
- Koenker, R. and Bassett, Jr., G. (1978). Regression quantiles. *Econometrica* **46**, 33-50.
- Lee, S., Seo, M.H. and Shin, Y. (2011). Testing for threshold effects in regression models. *J. Amer. Statist. Assoc.* **106**, 220-231.
- Li, C., Wei, Y., Chappell, R. and He, X. (2011). Bent line quantile regression with application to an allometric study of land mammals' speed and mass. *Biometrics* **67**, 242-249.
- Oka, T. and Qu, Z. (2011). Estimating structural changes in regression quantiles. *J. Econometrics* **162**, 248-267.

- Pennell, M., Whitmore, G. and Lee, M. (2010). Bayesian random-effects threshold regression with application to survival data with nonproportional hazards. *Bio-statistics* **11**, 111-126.
- Qu, Z. (2008). Testing for structural change in regression quantiles. *J. Econometrics* **146**, 170-184.
- Silverman, B.W. (1986). *Density estimation for statistics and data analysis*. Chapman and Hall.
- Su, L. and Xiao, Z. (2008). Testing for parameter stability in quantile regression models. *Statist. and Probability L.* **78**, 2768-2775.
- Tesfaye, F., Nawi, N., Van, M.H., Byass, P., Berhane, Y., Bonita, R. and Wall, S. (2007). Association between body mass index and blood pressure across three populations in Africa and Asia. *J. Hum Hypertens* **21**, 28-37.

Department of Statistics, Fudan University

E-mail: 10110690012@fudan.edu.cn

Department of Statistics, North Carolina State University

E-mail: hwang3@ncsu.edu

Department of Statistics, Fudan University

E-mail: zhuzy@fudan.edu.cn