

STATISTICAL INFERENCE FOR STRUCTURALLY CHANGED THRESHOLD AUTOREGRESSIVE MODELS

Zhaoxing Gao and Shiqing Ling

University of Chicago and Hong Kong University of Science and Technology

Abstract: In this study, we examine the theory and methodology of statistical inferences of thresholds and change-points in threshold autoregressive models. We show that least squares estimators (LSEs) of thresholds and change-points are n -consistent, and that they converge weakly to the minimizer of a compound Poisson process and the location of minima of a two-sided random walk, respectively. When the magnitude of the change in the parameters of the state regimes or in the time horizon is small, we further show that these limiting distributions can be approximated by a class of known distributions. The LSEs of the slope parameters are \sqrt{n} -consistent and asymptotically normal. Furthermore, a likelihood-ratio based confidence set is given for the thresholds and change-points, respectively. A Simulation study is carried out to assess the performance of our procedure, and the proposed theory and methodology are illustrated using a tree-ring data set.

Key words and phrases: Brownian motion, change-point, compound poisson process, least squares estimation, threshold.

1. Introduction

Structural changes have been studied as important problems in econometrics, engineering, and statistics for a long time. They are ubiquitous in economic and financial time series and were widely recognized as early as the 1940s; see Haavelmo (1944). As a result, many approaches have been developed to detect whether structural changes exist in a statistical model. For example, see recent articles by Ling (2007), Aue et al. (2009), and Hidalgo and Seo (2013), as well as the references therein. Hinkley (1970) was the first to investigate the maximum likelihood estimator (MLE) of the change-point in a sequence of independent and identically distributed (i.i.d.) random variables, and showed that the MLE converges in distribution to the location of the maxima of a double-sided random walk. Except for the i.i.d. normal or binomial random variables, its limiting distribution does not have a closed form and is difficult to use in practice. When the magnitude of change is small, Yao (1987) showed that Hinkley's limiting distribution can be approximated by a very nice distribution. Picard (1985) studied

the MLE of change-points in autoregressive (AR) models, obtaining the same limiting distribution as that in Yao (1987) when the magnitude of the change in the parameters approaches zero as the sample size tends to ∞ . Following this result, Bai (1994, 1995) and Bai, Lumsdaine and Stock (1998) studied the estimated change-points in a multivariate AR model and a co-integrated time series model; see also Chong (2001). Saikkonen, Lütkepohl and Trenkler (2006) and Kejriwal and Perron (2008) used a similar method to estimate the change-points in vector AR models and co-integrated regression models, respectively. Davis, Lee and Rodriguez-Yam (2006) proposed a minimum description length principle to locate the change-points in AR models with multiple structural changes. In an AR setting, Chan, Yau and Zhang (2014) used the group lasso method to estimate clusters of parameters with identical values over time, and Qian and Su (2014) considered estimations in time series with endogenous regressors and an unknown number of breaks using the group-fused lasso method. Ling (2016) developed an asymptotic theory for the change-points in linear and nonlinear time-series models. Other contributions include the monograph of Csörgö and Horváth (1997) and Shao and Zhang (2012), among others.

Existing change-point problems often refer to statistical models with a structural change in the time horizon. However, in a dynamic system, structural changes may occur over state regimes. The threshold autoregressive (TAR) model proposed by Tong (1978) captures these phenomena, and has been applied extensively in many areas, including economics, finance, biological, and environmental sciences, among others; see Chan and Kutoyants (2010) and Tong (2011) for comprehensive reviews of the TAR models. The asymptotic theory of the least squares estimator (LSE) for a two-regime TAR model was established by Chan (1993) and Chan and Tsay (1998), and then extended by Li and Ling (2012) and Li, Ling and Li (2013) to include multiple-regime TAR models and TMA models, respectively. Hansen (2000) studied the LSE for two-regime TAR/regression models when the threshold effect is vanishingly small. Seo and Linton (2007) proposed a smoothed LSE for the TAR/regression model, and showed that the estimated threshold is asymptotically normal with a slower rate of convergence. Liu, Ling and Shao (2011) and Gao, Tjøstheim and Yin (2013) studied the LSE for nonstationary first-order TAR models and Chan, Yau and Zhang (2015) adopted the lasso method to estimate TAR models with multiple thresholds. Gao, Ling and Tong (2018) proposed a non-nested test for TAR models versus smooth TAR models. However, existing research on threshold models is limited to the cases without change-points over the time horizon. The

main difficulty with a change-point lies in the nonsmooth and nonlinear functions of the threshold models. The smoothness of the objective function in terms of the parameters is required in Ling (2016) to establish the asymptotic properties; hence, his results cannot be applied to threshold models. Recently, Yau, Tang and Lee (2015) constructed a genetic algorithm to estimate multiple-regime TAR models with structural breaks. However, they established the consistency results of the parameters under their setting only, without further studying the limiting distributions of the slope parameters, thresholds, and change-points. To the best of our knowledge, no limiting distributions have been obtained when a structural change occurs over both the time horizon and the state regimes, and no methodology is available for statistical inferences of the thresholds and change-points in this case.

On the other hand, empirical time series often exhibit complex patterns, which may include nonlinearity and nonstationarity. For example, Tong and Lim (1980) analyzed the Canadian lynx data using TAR models and found obvious limit cycles. They also found that the one-step-ahead prediction is better than using pure AR models in terms of the root mean square error. Today, many long time-series sequences possess nonstationarities and thus, it is no longer adequate to characterize data using a single stationary model; see, Shao and Zhang (2012) for examples of some applications. Therefore, it is important and interesting to combine thresholds and change-points in order to characterize the nonlinearity and nonstationarity of a time series.

This study establishes the theory and methodology for statistical inferences of thresholds and change-points in TAR models. We first study the LSE of a TAR model with a structural change. Here, we show that both the estimated threshold and the change-point are n -consistent, and that they converge weakly to the minimizer of a compound Poisson process and the location of the minima of a two-sided random walk, respectively. When the magnitude of the change in the parameters of the state regimes or in the time horizon is small, we further show that these limiting distributions can be approximated by a class of known distributions. Furthermore, a likelihood-ratio based confidence set is given for the thresholds and change-points, respectively. Similarly to Bai (1994) and Bai (1997), we find that other estimated slope parameters are \sqrt{n} -consistent and asymptotically normal. To illustrate the proposed method, we apply it to study the growth of tree rings in China, finding evidence of possible climate change during the period 1641–1663.

This paper proceeds as follows. Section 2 presents the model and the esti-

mation procedure. The asymptotic properties are presented in Section 3. Section 4 presents the approximating distribution of these limiting distributions. A likelihood-ratio-based inference method is discussed in Section 5. Section 6 reports simulation results. Lastly, Section 7 presents our analysis of the data on the growth of tree rings. Owing to space constraints, some tables and figures and all proofs of the theorems in this paper are given in the Supplementary Material.

Throughout this paper, $\|\cdot\|$ denotes the Euclidean norm of a matrix or vector, $O_p(1)$ (or $o_p(1)$) denotes a series of random variables that are bounded (or converge to zero) in probability, \triangleq means “is defined as”, and $\rightarrow_{\mathcal{L}}$ denotes convergence in distribution.

2. The Model and Least Squares Estimation

We consider the following TAR model with order p [TAR(p)]:

$$y_t = \Phi' Z_{t-1} I(q_{t-1} > r) + \Psi' Z_{t-1} I(q_{t-1} \leq r) + \varepsilon_t, \quad (2.1)$$

where p is some known positive integer, $\Phi = (\mu, \phi_1, \dots, \phi_p)'$, $\Psi = (\nu, \psi_1, \dots, \psi_p)'$, $Z_t = (1, y_t, \dots, y_{t-p+1})'$, $I(\cdot)$ is an indicator function with a univariate threshold variable $q_t = q(y_t, y_{t-1}, \dots, y_{t-p+1})$, and $\{\varepsilon_t\}$ are errors. Note that $q(x)$ is a known function from R^p to R . Previous studies often consider $q_{t-1} = y_{t-d}$, for some integer $1 \leq d \leq p$. Here we consider only the simple case, as Section 3 of Chan (1993), in which there is no threshold effect on ε_t , because the general case is similar. Denote $\theta = (\Phi', \Psi)'$. We assume that $\theta \in \Theta \subset R^{2p+2}$ and $r \in \Gamma = [\underline{r}, \bar{r}] \subset R$, where Θ is a compact set. We denote model (2.1) by $Y(\theta, r)$, with $\theta \in \Theta$ and $r \in \Gamma$; that is, we write $\{y_1, \dots, y_m\} \in Y(\theta, r)$, for some $m > 0$, in the sense that they are generated from model (2.1) with parameters θ and r .

Let $\{y_1, y_2, \dots, y_n\}$ be a random sample. We assume that the data before time k are generated from $Y(\theta_1, r_1)$, and those after time k are generated from $Y(\theta_2, r_2)$; that is,

$$\{y_1, \dots, y_k\} \in Y(\theta_1, r_1) \quad \text{and} \quad \{y_{k+1}, \dots, y_n\} \in Y(\theta_2, r_2),$$

with $(\theta_1', r_1)' \neq (\theta_2', r_2)'$ and $k \in \{1, 2, \dots, n-1\}$. Here, k is called the unknown change-point and its true value is k_0 . The true parameters of $(\theta_1', r_1)'$ and $(\theta_2', r_2)'$ are $(\theta_{10}', r_{10})'$ and $(\theta_{20}', r_{20})'$, respectively. We parameterize the unknown change-point k as $k = [n\tau]$, with $\tau \in (0, 1)$ and $k_0 = [n\tau_0]$, where $[x]$ is the integer part of x . For each k , we use the pre-sample to estimate $(\theta_{10}', r_{10})'$ and the post-sample to estimate $(\theta_{20}', r_{20})'$ by least squares estimation. Let

$$\ell_t(\theta, r) = [y_t - \Phi'Z_{t-1}I(q_{t-1} > r) - \Psi'Z_{t-1}I(q_{t-1} \leq r)]^2. \tag{2.2}$$

Then, the corresponding objective functions are as follows:

$$S_{1n}(\theta_1, r_1, k) = \sum_{t=1}^k \ell_t(\theta_1, r_1) \quad \text{and} \quad S_{2n}(\theta_2, r_2, k) = \sum_{t=k+1}^n \ell_t(\theta_2, r_2). \tag{2.3}$$

The objective function based on the complete sample is

$$S_n(\theta_1, \theta_2, r_1, r_2, k) = S_{1n}(\theta_1, r_1, k) + S_{2n}(\theta_2, r_2, k). \tag{2.4}$$

The minimizer $(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n)$ of $S_n(\theta_1, \theta_2, r_1, r_2, k)$ is called the LSE of the true parameters, that is,

$$(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n) = \underset{\substack{\theta_1, \theta_2 \in \Theta, r_1, r_2 \in \Gamma \\ 1 \leq k < n}}{\arg \min} S_n(\theta_1, \theta_2, r_1, r_2, k).$$

We obtain $(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n)$ as follows. Using a similar method to that in Chan (1993), for each $1 \leq k < n$ and fixed r_i , we first obtain

$$\hat{\theta}_{in}(r_i, k) = \arg \min_{\theta_i \in \Theta} S_{in}(\theta_i, r_i, k), \quad i = 1, 2,$$

where $\hat{\theta}_{in}(r, k)$ depends on r and k . Specifically, we have

$$\hat{\Phi}_{1n}(r_1, k) = \left[\sum_{t=1}^k Z_{t-1}Z'_{t-1}I(q_{t-1} > r_1) \right]^{-1} \sum_{t=1}^k Z_{t-1}I(q_{t-1} > r_1)y_t. \tag{2.5}$$

The remaining parameters are derived in a similar way. Then, we can obtain the minimizers $\hat{r}_{in}(k)$ for the fixed k as follows:

$$\hat{r}_{in}(k) = \arg \min_{r_i \in \Gamma} S_{in}(\hat{\theta}_{in}(r_i, k), r_i, k), \quad i = 1, 2.$$

Finally, substituting $\hat{\theta}_{in}(\hat{r}_{in}(k), k)$ and $\hat{r}_{in}(k)$ into (2.4) yields

$$\hat{k}_n = \arg \min_{1 \leq k < n} S_n(\hat{\theta}_{1n}(\hat{r}_{1n}(k), k), \hat{\theta}_{2n}(\hat{r}_{2n}(k), k), \hat{r}_{1n}(k), \hat{r}_{2n}(k), k),$$

$$\hat{r}_{in} = \hat{r}_{in}(\hat{k}_n) \quad \text{and} \quad \hat{\theta}_{in} = \hat{\theta}_{in}(\hat{r}_{in}, \hat{k}_n), \quad i = 1, 2.$$

The range $1 \leq k < n$ can be replaced by $\tilde{p} \leq k \leq n - \tilde{p}$, for some integer \tilde{p} , in practice. Let $\{q_{(1)}, q_{(2)}, \dots, q_{(k)}\}$ and $\{q_{(k+1)}, \dots, q_{(n)}\}$ be the corresponding order statistics of the subsamples $\{q_1, q_2, \dots, q_k\}$ and $\{q_{k+1}, \dots, q_n\}$. Then $S_{1n}(\hat{\theta}_{1n}(r_1, k), r_1, k)$ and $S_{2n}(\hat{\theta}_{2n}(r_2, k), r_2, k)$ are constants when $r_1 \in [q_{(i)}, q_{(i+1)})$ and $r_2 \in [q_{(j)}, q_{(j+1)})$. Thus, for each k , there exist two intervals, say $[q_{(i)}, q_{(i+1)})$ and $[q_{(j)}, q_{(j+1)})$, on which $S_{1n}(\hat{\theta}_{1n}(r_1, k), r_1, k)$ and $S_{2n}(\hat{\theta}_{2n}(r_2, k), r_2, k)$, respectively, achieve their global minima. We take $\hat{r}_{1n}(k) = q_{(i)}$ and $\hat{r}_{2n}(k) = q_{(j)}$. It is not difficult to show that $(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n)$ is the LSE of $(\theta_{10}, \theta_{20}, r_{10}, r_{20}, k_0)$.

We focus on model (2.1) with one change-point. If model (2.1) has m unknown change-points $(k_2, k_3, \dots, k_{m+1})$, such that $1 < k_2 < k_3 < \dots < k_{m+1} < n$, and the corresponding changed parameters are $\{(\theta_j, r_j) : j = 1, 2, \dots, m+1\}$, then the profile objective function becomes

$$S_n(k_2, \dots, k_{m+1}) = \sum_{j=1}^{m+1} \min_{\theta_j, r_j} \sum_{t=k_j+1}^{k_{j+1}} \ell_t(\theta_j, r_j), \quad (2.6)$$

where $k_1 = 0$ and $k_{m+2} = n$. The estimators of the true change-points are

$$(\hat{k}_2, \dots, \hat{k}_{m+1}) = \arg \min_{(k_2, \dots, k_{m+1}) \in \Lambda} S_n(k_2, \dots, k_{m+1}), \quad (2.7)$$

where Λ is some appropriate partition set. In this case, the dynamic programming algorithm of Bai and Perron (2003) can be used to search for the thresholds and change-points. However, determining the number of change-points m and establishing the asymptotic results remain challenging, and can be achieved by following the approach of Bai and Perron (1998). Yau, Tang and Lee (2015) proposed an efficient algorithm to estimate the number of change-points together with all other parameters using a different approach, but their asymptotic distributions are still unclear.

3. Asymptotic Properties

We first study the rates of convergence of the estimated threshold and change-point, which are essential in threshold and change-point problems. We first need to make several assumptions.

Assumption 1. $\varepsilon_t = \sigma_{10}u_t$ for $t \leq k_0$, and $\varepsilon_t = \sigma_{20}u_t$ for $t > k_0$, where u_t is a sequence of i.i.d. random variables with $Eu_1 = 0$, $Eu_1^2 = 1$ and has an absolutely continuous distribution with a uniformly continuous and positive density $f_u(x)$ on R . Furthermore, $E|u_t|^{2+\nu} < \infty$, for some $\nu \in (0, 1)$.

Assumption 2. q_t has an absolutely continuous distribution with a uniformly continuous and positive density function $\pi_i(r)$ on Γ , where $i = 1$ when $t \leq k_0$, and $i = 2$ when $t > k_0$.

Assumption 3. $(\theta'_{10}, r_{10})'$ and $(\theta'_{20}, r_{20})'$ are interior points in $\Theta \times \Gamma$, and $\Phi_{10} \neq \Psi_{10}$, $\Phi_{20} \neq \Psi_{20}$, and $(\theta'_{10}, r_{10})' \neq (\theta'_{20}, r_{20})'$.

Assumption 1 allows for a variance change on the errors across the change-point. The conditions of the density function are the same as those in Condition 2 in Chan (1993). This implies Assumption 2 and entails that $\pi(\cdot)$ is bounded,

continuous, and positive on Γ if we take $q_{t-1} = y_{t-d}$ as a special case; see (ii) in Remark B of Chan (1993). Assumption 3 guarantees the identification of r_{10} , r_{20} , and k_0 . In order to obtain the consistency of $\hat{\theta}_{in}$ and \hat{r}_{in} , for $i = 1, 2$, we need an assumption similar to Condition 1 in Chan (1993). Let $\mathcal{Z}_{it} = (y_t, \dots, y_{t-p+1}) \in Y(\theta_{i0}, r_{i0})$, for $i = 1, 2$. Then, \mathcal{Z}_{it} is a Markov chain. Denote its l -step transition probability by $P^l(x, A)$, where $x \in R$ and A is a Borel set of R .

Assumption 4. \mathcal{Z}_{it} admits a unique invariant measure $\Pi_i(\cdot)$ such that there exist a $K > 0$ and $\rho \in [0, 1)$, and for any $x \in R$ and any positive integer l , $\|P^l(x, \cdot) - \Pi_i(\cdot)\|_v \leq K(1 + \|x\|)\rho^l$, where $\|\cdot\|_v$ and $\|\cdot\|$ denote the total variation norm and the Euclidean norm, respectively.

When $t \leq k_0$, $\ell_t(\theta, r) = \ell(\theta, r, y_t, \dots, y_1, \mathcal{Z}_{10})$, and when $t > k_0$, $\ell_t(\theta, r) = \ell(\theta, r, y_t, \dots, y_{k_0+1}, \mathcal{Z}_{2k_0})$, where $\ell(\cdot)$ is a measurable function of $\{y_t\}$ with parameters θ and r . That is, there are two processes $\{y_{1t}\} \in Y(\theta_{10}, r_{10})$ and $\{y_{2t}\} \in Y(\theta_{20}, r_{20})$ and we observe that $y_t = y_{1t}$ when $t \leq k_0$, and $y_t = y_{2t}$ when $t > k_0$. A set of sufficient conditions for Assumption 4 is given by

$$\max_{i=1,2} \left\{ \sum_{j=1}^p |\phi_{ij}|, \sum_{j=1}^p |\psi_{ij}| \right\} < 1$$

and Assumption 1; see Chan and Tong (1985) and Chan (1989). When $p = 1$, the above coefficient condition can be weakened to $\phi_{i1} < 1$, $\psi_{i1} < 1$, and $\phi_{i1}\psi_{i1} < 1$, for $i = 1, 2$. Under Assumption 4, $\{\mathcal{Z}_{it}\}$ is V -uniformly ergodic with $V(x) = K(1 + \|x\|)$; see Meyn and Tweedie (2009). This condition is stronger than geometric ergodicity and the strong mixing (α -mixing) condition; see the definition in Rosenblatt (1956) and the discussion in Hansen (2000). If the initial values \mathcal{Z}_{10} and \mathcal{Z}_{2k_0} are from the distributions $\Pi_1(\cdot)$ and $\Pi_2(\cdot)$, respectively, then Assumption 4 implies that $\{y_t\}_{t=1}^{k_0}$ and $\{y_t\}_{k_0+1}^n$ are two strictly stationary and ergodic sequences. In practice, the initial values \mathcal{Z}_{10} and \mathcal{Z}_{2k_0} are replaced by some chosen constants. We discuss their effect on the estimators in Section 5.

Our first result is stated as follows.

Theorem 1. *If Assumptions 1–4 hold, then*

- (a) $\hat{\theta}_{in} = \theta_{i0} + o_p(1)$ and $\hat{r}_{in} = r_{i0} + o_p(1)$, $i = 1, 2$,
- (b) $\hat{k}_n = k_0 + O_p(1)$.

From Theorem 1, we have that the thresholds and the slope parameters are all consistent. Similar results are obtained in Theorem 2 of Yau, Tang and Lee (2015). We can write $\hat{k}_n = [n\hat{\tau}_n]$, in which case, $\hat{\tau}_n$ is an estimator of τ_0 .

Theorem 1 implies that the rate of convergence of $\hat{\tau}_n$ is n , which is the same as those in Bai and Perron (1998) and Yau, Tang and Lee (2015), but is faster than those in Picard (1985) and Bai, Lumsdaine and Stock (1998) for AR models because their convergence rate is essentially nd_n^2 , where d_n is the magnitude of the change as $d_n \rightarrow 0$. To obtain the rate of convergence of \hat{r}_{in} , we need two further assumptions. Define $M_i(r) = E(Z_t Z_t' | q_t = r)$, where $i = 1$ when $t \leq k_0$, and $i = 2$ when $t > k_0$.

Assumption 5. (i) $M_i(r)$ is continuous at $r = r_{i0}$; (ii) $(\Psi_{i0} - \Phi_{i0})' M_i(r_{i0}) (\Psi_{i0} - \Phi_{i0}) > 0$; $i = 1, 2$.

We say that model (2.1) has a discontinuous AR function if Assumption 5 is satisfied. For Assumption 5 to hold, we require the threshold variable to have a continuous distribution; see Assumption 1.5 of Hansen (2000) or Assumption 2.2 of Gao, Ling and Tong (2018) for further details. Assumption 5 (ii) is natural for a positive-definite matrix $M_i(r_{i0})$. Note that if we choose $q_{t-1} = y_{t-d}$, as in Chan (1993), Assumption 5(ii) implies Condition 4 in Chan (1993) and the threshold r_{i0} becomes the jump-point of the AR function. For simplicity, let

$$M_i(r^+) = E Z_t Z_t'(r^+), M_i(r^-) = E Z_t Z_t'(r^-) \text{ and } M_i(r_1, r_2) = E Z_t Z_t'(r_1, r_2),$$

where $i = 1$ when $1 \leq t \leq k_0$, $i = 2$ when $k_0 + 1 \leq t \leq n$, $Z_t(r^+) = Z_t I(q_t > r)$, $Z_t(r^-) = Z_t I(q_t \leq r)$, and $Z_t(r_1, r_2) = Z_t I(r_1 < q_t \leq r_2)$. Under Assumptions 1–3, it is not difficult to show that $M_i(r^+) > 0$, $M_i(r^-) > 0$, and $M_i(r_1, r_2) > 0$, for all $r, r_1, r_2 \in \Gamma$ with $r_1 < r_2$, $i = 1, 2$. By applying the techniques in Chan (1993) to each segment of $\{y_t\}$, we have the convergence rate of \hat{r}_{in} and the asymptotic normality of $\hat{\theta}_{in}$, as follows.

Theorem 2. *If Assumptions 1–5 hold, then*

$$\begin{aligned} (a) \quad & n(\hat{r}_{in} - r_{i0}) = O_p(1), \\ (b) \quad & \sqrt{n}(\hat{\theta}_{in} - \theta_{i0}) \rightarrow_{\mathcal{L}} N(0, \sigma_{i0}^2 \Sigma_i^{-1}), \end{aligned}$$

for $i = 1, 2$, as $n \rightarrow \infty$, where $\Sigma_1 = \tau_0 \text{diag}\{M_1(r_{10}^+), M_1(r_{10}^-)\}$ and $\Sigma_2 = (1 - \tau_0) \text{diag}\{M_2(r_{20}^+), M_2(r_{20}^-)\}$. Furthermore, $n(\hat{r}_{in} - r_{i0})$ is asymptotically independent of $\sqrt{n}(\hat{\theta}_{in} - \theta_{i0})$, which is always asymptotically normal, and $\hat{\theta}_{1n}$ and $\hat{\theta}_{2n}$ are asymptotically independent of each other.

This theorem shows that the rate of convergence of \hat{r}_{in} is the same as those in Chan (1993) and Yau, Tang and Lee (2015), but is faster than those in Hansen (2000) and Seo and Linton (2007) because their convergence rate is essentially $n^{1-2\tilde{\alpha}}$, for some $\tilde{\alpha} \in (0, 1/2)$, by assuming a vanishingly small threshold effect.

Whereas Yau, Tang and Lee (2015) obtained only the \sqrt{n} -consistency of the slope parameters, we show that these parameters have the same asymptotic properties as those in Picard (1985) and Bai (1997), among others; that is, they are not affected by the threshold parameters.

In Theorem 2, τ_0 , σ_{i0}^2 , and Σ_i are not known, in practice, but they can be replaced by consistent estimates from $(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n)$. For example, σ_{i0}^2 can be estimated by

$$\hat{\sigma}_{1n}^2 = \frac{1}{\hat{k}_n} \sum_{t=1}^{\hat{k}_n} \ell_t(\hat{\theta}_{1n}, \hat{r}_{1n}), \quad \hat{\sigma}_{2n}^2 = \sum_{t=\hat{k}_n+1}^n \ell_t(\hat{\theta}_{2n}, \hat{r}_{2n}), \quad (3.1)$$

and $M_i(r_{i0}^\pm)$ can be estimated by

$$\hat{M}_1(\hat{r}_{1n}^\pm) = \frac{1}{\hat{k}_n} \sum_{t=1}^{\hat{k}_n} Z_t Z_t'(\hat{r}_{1n}^\pm), \quad \hat{M}_2(\hat{r}_{2n}^\pm) = \frac{1}{n - \hat{k}_n} \sum_{t=\hat{k}_n+1}^n Z_t Z_t'(\hat{r}_{2n}^\pm), \quad (3.2)$$

respectively. Hence, Σ_i can be estimated by

$$\hat{\Sigma}_{1n} = \hat{\tau}_n \text{diag}\{\hat{M}_1(\hat{r}_{1n}^+), \hat{M}_1(\hat{r}_{1n}^-)\} \quad \text{and} \quad \hat{\Sigma}_{2n} = (1 - \hat{\tau}_n) \text{diag}\{\hat{M}_2(\hat{r}_{2n}^+), \hat{M}_2(\hat{r}_{2n}^-)\}. \quad (3.3)$$

Using Theorems 1–2, it is not difficult to show that the above sample estimators are all consistent with their corresponding true values and, hence, can be used for statistical inferences.

To study the limiting distributions of \hat{r}_{in} , we consider the profile objective function

$$\tilde{S}_{in}(z, \hat{k}_n) \equiv S_{in}\left(\hat{\theta}_{in}\left(r_{i0} + \frac{z}{n}, \hat{k}_n\right), r_{i0} + \frac{z}{n}, \hat{k}_n\right) - S_{in}(\hat{\theta}_{in}(r_{i0}, \hat{k}_n), r_{i0}, \hat{k}_n), \quad (3.4)$$

where $z \in R$, $i = 1, 2$. According to the procedure for \hat{r}_{in} , we have that

$$n(\hat{r}_{in} - r_{i0}) = \arg \min_{z \in R} \tilde{S}_{in}(z, \hat{k}_n). \quad (3.5)$$

Now, we define two jump processes, as follows:

$$\tilde{S}_{1n}(k_0, \theta_{10}, r_{10}, r_1) = \begin{cases} \sum_{t=1}^{k_0} \xi_{1t} I(r_1 < q_{t-1} \leq r_{10}), & r_1 < r_{10}, \\ 0, & r_1 = r_{10}, \\ \sum_{t=1}^{k_0} \eta_{1t} I(r_{10} < q_{t-1} \leq r_1), & r_1 > r_{10}, \end{cases} \quad (3.6)$$

$$\tilde{S}_{2n}(k_0, \theta_{20}, r_{20}, r_2) = \begin{cases} \sum_{t=k_0+1}^n \xi_{2t} I(r_2 < q_{t-1} \leq r_{20}), & r_2 < r_{20}, \\ 0, & r_2 = r_{20}, \\ \sum_{t=k_0+1}^n \eta_{2t} I(r_{20} < q_{t-1} \leq r_2), & r_2 > r_{20}, \end{cases} \quad (3.7)$$

where $\xi_{it} = (\Phi_{i0} - \Psi_{i0})' Z_{t-1} Z_{t-1}' (\Phi_{i0} - \Psi_{i0}) - 2(\Phi_{i0} - \Psi_{i0})' Z_{t-1} \varepsilon_t$ and $\eta_{it} = (\Psi_{i0} -$

$\Phi_{i0})'Z_{t-1}Z'_{t-1}(\Psi_{i0} - \Phi_{i0}) - 2(\Psi_{i0} - \Phi_{i0})'Z_{t-1}\varepsilon_t$, for $i = 1, 2$. We reparameterize r_i as $r_{i0} + z/n$ in (3.6) and (3.7). On the event $\{|z| \leq B, |\hat{k}_n - k_0| \leq M\}$ for any fixed $B, M \in (0, \infty)$, $\tilde{\mathcal{S}}_{in}(z, \hat{k}_n)$ can be approximated by $\tilde{\mathcal{S}}_{in}(k_0, \theta_{i0}, r_{i0}, r_{i0} + z/n)$ in $D(R)$, the space of all càdlàg functions on R equipped with the Skorokhod metric, as follows:

$$\tilde{\mathcal{S}}_{in}(z, \hat{k}_n) = \tilde{\mathcal{S}}_{in}\left(k_0, \theta_{i0}, r_{i0}, r_{i0} + \frac{z}{n}\right) + o_p(1), \quad i = 1, 2; \tag{3.8}$$

see the proof of Lemma 4 in the Supplementary Material. We show that $\tilde{\mathcal{S}}_{in}(k_0, \theta_{i0}, r_{i0}, r_{i0} + z/n)$ weakly converges to a two-sided compound Poisson process $\mathcal{P}_1(\tau_0 z)$ for $i = 1$, and to $\mathcal{P}_2((1 - \tau_0)z)$ for $i = 2$, which are defined as follows:

$$\mathcal{P}_i(z) = I(z < 0) \sum_{t=1}^{N_1^i(-z)} Y_{it} + I(z \geq 0) \sum_{t=1}^{N_2^i(z)} Z_{it}, \quad z \in R, i = 1, 2, \tag{3.9}$$

where $\{N_1^i(z), z \geq 0\}$ and $\{N_2^i(z), z \geq 0\}$ are two independent Poisson processes with $N_1^i(0) = 0$ and $N_2^i(0) = 0$ a.s. and with the same jump rate $\pi_i(r_{i0})$, where $\pi_i(\cdot)$ is the density function of $\{q_t\}$; see Assumption 2. $\{Y_{it}, t \geq 1\}$ are i.i.d. random variables with a distribution function $F_{1i}(\cdot|r_{i0})$, and $\{Z_{it}, t \geq 1\}$ are i.i.d. random variables with a distribution function $F_{2i}(\cdot|r_{i0})$. Here, $F_{1i}(\cdot|x)$ and $F_{2i}(\cdot|x)$ are the conditional distribution functions of ξ_{it} and η_{it} , respectively, given $q_{t-1} = x$, and $\{Y_{it}, t \geq 1\}$ and $\{Z_{it}, t \geq 1\}$ are mutually independent. Clearly, $\mathcal{P}_i(z) \rightarrow +\infty$ a.s. when $|z| \rightarrow \infty$ because $EY_{it} = EZ_{it} > 0$ from Assumptions 3–4. Therefore, there exists a unique random interval $[M_-^{(i)}, M_+^{(i)})$ on which the process $\mathcal{P}_i(z)$ attains its global minimum a.s.. That is,

$$[M_-^{(i)}, M_+^{(i)}) = \arg \min_{z \in R} \mathcal{P}_i(z), \quad i = 1, 2. \tag{3.10}$$

To obtain the limiting distribution of \hat{k}_n , we define a two-sided random walk as follows:

$$W(k, \theta_{10}, \theta_{20}, r_{10}, r_{20}) = \begin{cases} \sum_{t=k}^{-1} [\ell_t(\theta_{20}, r_{20}) - \ell_t(\theta_{10}, r_{10})], & k < 0, \\ 0, & k = 0, \\ \sum_{t=1}^k [\ell_t(\theta_{10}, r_{10}) - \ell_t(\theta_{20}, r_{20})], & k > 0, \end{cases} \tag{3.11}$$

where $\ell_t(\theta, r)$ is defined in (2.2). Here, $y_t \in Y(\theta_{10}, r_{10})$ when $t < 0$, and $y_t \in Y(\theta_{20}, r_{20})$ when $t > 0$. Now we can state our next theorem.

Theorem 3. *If Assumptions 1–5 hold, then \hat{k}_n , \hat{r}_{1n} , and \hat{r}_{2n} are asymptotically independent of each other, and*

$$(a) \quad n(\hat{r}_{1n} - r_{10}) \xrightarrow{\mathcal{L}} \frac{1}{\tau_0} M_-^{(1)}$$

$$\text{and } n(\hat{r}_{2n} - r_{20}) \xrightarrow{\mathcal{L}} \frac{1}{1 - \tau_0} M_-^{(2)},$$

$$(b) \quad \hat{k}_n - k_0 \xrightarrow{\mathcal{L}} \arg \min_k W(k, \theta_{10}, \theta_{20}, r_{10}, r_{20}),$$

as $n \rightarrow \infty$.

From this theorem, we have that the limiting distribution of \hat{r}_{in} is the same as that in Chan (1993), subject to a scale τ_0 or $1 - \tau_0$. The limiting distribution of the estimated change-point corresponds to that of the MLE in Ling (2016) and is related to a two-sided random walk, even if the objective function is not continuous. On the other hand, this theorem can be treated as a complement to the results in Yau, Tang and Lee (2015) because they obtained only the consistency results, without investigating the limiting distributions of the thresholds and the change-point.

4. Approximating the Limiting Distributions

Except for the MLE of the threshold in a simple regression model with i.i.d. data in Yu (2012), we do not have a closed-form solution for the estimated threshold, in general. The distributions in Theorem 3 are difficult to use directly for statistical inferences. Li and Ling (2012) proposed a numerical method to simulate the limiting distribution of the estimated thresholds when the threshold effect is fixed, whereas Hansen (2000) adopted a different approach, obtaining a closed form for the distribution by assuming the threshold effect is vanishingly small as the sample size $n \rightarrow \infty$. In this section, we use the method of Hansen (2000) to obtain the approximating distributions of r_{10} and r_{20} . By borrowing a similar idea to those in Bai (1997) and Hansen (2000), we also obtain the approximating distribution of $W(k, \theta_{10}, \theta_{20}, r_{10}, r_{20})$. Our discussions are separated into two subsections.

4.1. Approximating distributions of the estimated r_{10} and r_{20}

Let $\bar{\xi}_{it} = \delta'_{in} Z_{t-1} Z'_{t-1} \delta_{in} + 2\delta'_{in} Z_{t-1} \varepsilon_t$ and $\bar{\eta}_{it} = \delta'_{in} Z_{t-1} Z'_{t-1} \delta_{in} - 2\delta'_{in} Z_{t-1} \varepsilon_t$, for $i = 1, 2$. Similarly to (3.6)–(3.7), we first define the following two processes:

$$\bar{S}_{1n}(k_0, \theta_{10}, r_{10}, r_1) = \begin{cases} \sum_{t=1}^{k_0} \bar{\xi}_{1t} I(r_1 < q_{t-1} \leq r_{10}), & r_1 < r_{10}, \\ 0, & r_1 = r_{10}, \\ \sum_{t=1}^{k_0} \bar{\eta}_{1t} I(r_{10} < q_{t-1} \leq r_1), & r_1 > r_{10}, \end{cases} \quad (4.1)$$

and

$$\bar{S}_{2n}(k_0, \theta_{20}, r_{20}, r_2) = \begin{cases} \sum_{t=k_0+1}^n \bar{\xi}_{2t} I(r_2 < q_{t-1} \leq r_{20}), & r_2 < r_{20}, \\ 0, & r_2 = r_{20}, \\ \sum_{t=k_0+1}^n \bar{\eta}_{2t} I(r_{20} < q_{t-1} \leq r_2), & r_2 > r_{20}. \end{cases} \quad (4.2)$$

The difference between the processes in (3.6)–(3.7) and those in (4.1)–(4.2) lies in (ξ_{it}, η_{it}) and $(\bar{\xi}_{it}, \bar{\eta}_{it})$. Here, y_t in (ξ_{it}, η_{it}) depends on Φ_{i0} and Ψ_{i0} , and hence is changing if $\Psi_{i0} - \Phi_{i0} \rightarrow 0$. However, y_t in $(\bar{\xi}_{it}, \bar{\eta}_{it})$ is irrelevant to δ_{in} , and hence is still stationary and ergodic when $\delta_{in} \rightarrow 0$. This is why we introduce the two new processes. We first make one additional assumption.

Assumption 6. $\delta_{in} = c_i n^{-\beta}$, where $\beta \in (0, 1/2)$, and c_1 and c_2 are nonzero constant vectors.

Define

$$\hat{z}_{in} = \arg \min_{z \in R} \bar{S}_{in} \left(k_0, \theta_{i0}, r_{i0}, r_{i0} + \frac{z}{n} \right), \quad i = 1, 2, \quad (4.3)$$

and $B(r)$ is a standard Brownian motion on $(-\infty, \infty)$.

From Assumptions 1–5, 6, and the proof of Theorem 1 in Hansen (2000), we easily obtain the following theorem.

Theorem 4. *If Assumptions 1–5 and 6 hold and $E y_t^4 < \infty$, then*

$$n^{-2\beta} \hat{z}_{in} \xrightarrow{\mathcal{L}} w_i \arg \min_{-\infty < r < \infty} \left[\frac{|r|}{2} + B(r) \right], \quad i = 1, 2, \quad (4.4)$$

where $w_1 = \sigma_{10}^2 / [\tau_0 c_1' M_1(r_{10}) c_1 \pi_1(r_{10})]$, $w_2 = \sigma_{20}^2 / [(1 - \tau_0) c_2' M_2(r_{20}) c_2 \pi_2(r_{20})]$, $M_1(r) = E(Z_t Z_t' | q_t = r)$, with $t \leq k_0$, and $M_2(r) = E(Z_t Z_t' | q_t = r)$, with $t > k_0$.

Remark 1. The finite fourth moment in Theorem 4 is inherited from Hansen (2000). In case of a threshold effect in the variance in model (2.1) as Chan (1993), Li and Ling (2012) discussed the limiting distribution of the estimated threshold. Here, the approximation in Theorem 4 should be asymmetric. For a further discussion, see Yu (2012) and Yu (2015).

Let $T = \arg \min_{-\infty < r < \infty} [|r|/2 + B(r)]$ and let $\Phi(x)$ denote the cumulative standard normal distribution function. Then, for $x \geq 0$,

$$P(T \leq x) = 1 + \sqrt{\frac{x}{2\pi}} \exp\left(-\frac{x}{8}\right) + \frac{3}{2} \exp(x) \Phi\left(-\frac{3\sqrt{x}}{2}\right) - \left(\frac{x+5}{2}\right) \Phi\left(-\frac{\sqrt{x}}{2}\right).$$

and for $x < 0$, $P(T \geq x) = 1 - P(T \geq -x)$; see Yao (1987). By Theorem 4, if $\Psi_{i0} - \Phi_{i0} \approx \delta_{in}$, then we can use the following approximation:

$$n^{1-2\beta} (\hat{r}_{in} - r_{i0}) / w_i \approx_d \arg \min_{-\infty < r < \infty} \left[\frac{|r|}{2} + B(r) \right], \quad i = 1, 2, \quad (4.5)$$

where $n^{1-2\beta}/w_i$ can be replaced by $\sigma_{10}^{-2}[(\Psi_{10} - \Phi_{10})'M_1(r_{10})(\Psi_{10} - \Phi_{10})\pi_1(r_{10})]k_0$ when $i = 1$, and by $\sigma_{20}^{-2}[(\Psi_{20} - \Phi_{20})'M_2(r_{20})(\Psi_{20} - \Phi_{20})\pi_2(r_{20})](n - k_0)$ when $i = 2$. In practice, k_0 can be replaced by $[n\hat{\tau}_n]$, other true values can be estimated consistently in a similar way to (3.1)–(3.2), $\pi_i(r_{i0})$ can be estimated by its kernel density estimator, and $M_i(r_{i0})$ can be estimated by a polynomial regression, as in Hansen (2000). Thus, we can use the distribution of T to make statistical inferences for the thresholds r_{10} and r_{20} .

4.2. Approximating distribution of the estimated k_0

In this subsection, we investigate the limiting distribution of $W(k, \theta_{10}, \theta_{20}, r_{10}, r_{20})$ in Theorem 3. From (S2.24) in Section S2 of the Supplementary Material, we have

$$\hat{k}_n = \arg \min_{1 \leq k < n} \left\{ I(k < k_0) \sum_{t=k+1}^{k_0} (A_{1t}^2 + 2A_{1t}\varepsilon_t) + I(k \geq k_0) \sum_{t=k_0+1}^k (A_{2t}^2 + 2A_{2t}\varepsilon_t) + o_p(1) \right\}, \tag{4.6}$$

where

$$A_{1t} = (\Phi_{10} - \Phi_{20})'Z_{t-1}I(q_{t-1} > r_{20}) + (\Psi_{10} - \Psi_{20})'Z_{t-1}I(q_{t-1} \leq r_{10}) + (\Phi_{10} - \Psi_{20})'Z_{t-1}I(r_{10} < q_{t-1} \leq r_{20}) \tag{4.7}$$

and $A_{2t} = -A_{1t}$ if $r_{20} > r_{10}$, and

$$A_{1t} = (\Phi_{10} - \Phi_{20})'Z_{t-1}I(q_{t-1} > r_{10}) + (\Psi_{10} - \Psi_{20})'Z_{t-1}I(q_{t-1} \leq r_{20}) + (\Psi_{10} - \Phi_{20})'Z_{t-1}I(r_{20} < q_{t-1} \leq r_{10}) \tag{4.8}$$

and $A_{2t} = -A_{1t}$ if $r_{10} > r_{20}$. We only consider the case of $r_{20} > r_{10}$ in (4.6) because it is similar to the case when $r_{10} > r_{20}$. Now, we define the following process:

$$\bar{W}_n(k) = I(k < k_0) \sum_{t=k+1}^{k_0} (\bar{A}_{1t}^2 + 2\bar{A}_{1t}\varepsilon_t) + I(k \geq k_0) \sum_{t=k_0+1}^k (\bar{A}_{2t}^2 + 2\bar{A}_{2t}\varepsilon_t), \tag{4.9}$$

where

$$\bar{A}_{1t} = \kappa'_{1n}Z_{t-1}I(q_{t-1} > r_{20}) + \kappa'_{2n}Z_{t-1}I(q_{t-1} \leq r_{10}) + \kappa'_{3n}Z_{t-1}I(r_{10} < q_{t-1} \leq r_{20}) \tag{4.10}$$

and $\bar{A}_{2t} = -\bar{A}_{1t}$. Here, y_t in A_{it} depends on Φ_{i0} and Ψ_{i0} , and hence is changing if $\Phi_{20} - \Phi_{10} \rightarrow 0$, $\Psi_{20} - \Psi_{10} \rightarrow 0$, and $\Psi_{20} - \Phi_{10} \rightarrow 0$. However, y_t in \bar{A}_{it} is irrelevant to κ_{in} and is stationary and ergodic when $\kappa_{in} \rightarrow 0$. Next, we make one

more assumption.

Assumption 7. $\kappa_{1n} = c_3 n^{-\beta}$, $\kappa_{2n} = c_4 n^{-\beta}$, and $\kappa_{3n} = c_5 n^{-\beta}$, where $\beta \in (0, 1/2)$, c_3, c_4 , and c_5 are nonzero constant vectors.

We parameterize k as $k = k_0 + [n^{2\beta} s]$ in (4.9). From Assumptions 1–5 and Theorem 1 in the Supplementary Material, we can show that the sum of \bar{A}_{it}^2 goes to $\varpi_i |s|$ in probability and uniformly on any compact set of s , for $i = 1, 2$, where

$$\varpi_i = c_3' M_i(r_{20}^+) c_3 + c_4' M_i(r_{10}^-) c_4 + c_5' M_i(r_{10}, r_{20}) c_5.$$

By Theorem A.1 in Li, Ling and Zhang (2016), the sum of $\bar{A}_{it} \varepsilon_t$ weakly converges to a Gaussian process $G(s)$ with covariance kernel $\varpi_i \sigma^2 (|s| \wedge |t|)$, $i = 1, 2$, respectively. Now we can state the following theorem.

Theorem 5. *If Assumptions 1–5 and 7 hold, then*

$$\sigma_{10}^{-2} \varpi_1 \arg \min_{-\infty < s < \infty} \bar{W}_n \left(k_0 + [n^{2\beta} s] \right) \rightarrow_{\mathcal{L}} T_{\phi, \xi},$$

where $\phi = \varpi_2 \sigma_{20}^2 / (\varpi_1 \sigma_{10}^2)$, $\xi = \varpi_2 / \varpi_1$, and

$$T_{\phi, \xi} = \arg \min_{s \in R} \left\{ \left[\frac{\xi}{2} |s| + \sqrt{\phi} B(s) \right] I(s \geq 0) + \left[\frac{1}{2} |s| + B(s) \right] I(s < 0) \right\},$$

where $M_i(r_{i0}^+)$ and $M_i(r_{i0}^-)$ are defined as in Theorem 2.

Remark 2. Under Assumption 7, the third term in (4.10) and in ϖ_i vanishes as $r_{20} - r_{10} \rightarrow 0$, and the assumption on κ_{3n} is redundant if $r_{20} = r_{10}$. Alternatively, if κ_{3n} is fixed and $r_{20} - r_{10}$ is vanishingly small, such that it matches the convergence rate of s in Theorem 5, we obtain a similar result.

The distribution of $T_{\phi, \xi}$ can be found in Bai (1997) and its density is asymmetric unless $\phi = \xi = 1$. We can use the cumulative distribution functions in Appendix B of Bai (1997) to construct confidence intervals once we know the values of ϕ and ξ .

If $\Phi_{20} - \Phi_{10} \approx \kappa_{1n}$, $\Psi_{20} - \Psi_{10} \approx \kappa_{2n}$, and $\Phi_{10} - \Psi_{20} \approx \kappa_{3n}$, by Theorem 5, (4.6) and (4.9) and parameterizing $k = k_0 + [n^{2\beta} s]$, we have

$$\frac{\varpi_1}{\sigma_{10}^2 n^{2\beta}} (\hat{k}_n - k_0) \approx \frac{\varpi_1}{\sigma_{10}^2} \arg \min_{-\infty < s < \infty} \bar{W}_n(k_0 + [n^{2\beta} s]) \approx_d T_{\phi, \xi}. \tag{4.11}$$

Here, we can approximate ϕ , ξ , and $\varpi_1 / (\sigma^2 n^{2\beta})$ as follows:

$$\phi \approx \frac{d_2 \sigma_{20}^2}{d_1 \sigma_{10}^2}, \quad \xi \approx \frac{d_2}{d_1}, \quad \text{and} \quad \frac{\varpi_1}{\sigma_{10}^2 n^{2\beta}} \approx \frac{d_1}{\sigma_{10}^2}, \tag{4.12}$$

where $d_i = (\Phi_{20} - \Phi_{10})' M_i(r_{20}^+) (\Phi_{20} - \Phi_{10}) + (\Psi_{20} - \Psi_{10})' M_i(r_{10}^-) (\Psi_{20} - \Psi_{10}) + (\Phi_{10} - \Psi_{20})' M_i(r_{10}, r_{20}) (\Phi_{10} - \Psi_{20})$, for $i = 1, 2$. In practice, d_i can be estimated

by

$$\begin{aligned} \hat{d}_{in} &= (\hat{\Phi}_{2n} - \hat{\Phi}_{1n})' \hat{M}_i(\hat{r}_{2n}^+) (\hat{\Phi}_{2n} - \hat{\Phi}_{1n}) + (\hat{\Psi}_{2n} - \hat{\Psi}_{1n})' \hat{M}_i(\hat{r}_{1n}^-) (\hat{\Psi}_{2n} - \hat{\Psi}_{1n}) \\ &\quad + (\hat{\Phi}_{1n} - \hat{\Psi}_{2n})' \hat{M}_i(\hat{r}_{1n}, \hat{r}_{2n}) (\hat{\Phi}_{1n} - \hat{\Psi}_{2n}), \quad i = 1, 2, \end{aligned} \tag{4.13}$$

where $\hat{M}_i(\hat{r}_{1n}, \hat{r}_{2n})$ is defined in a similar way to $\hat{M}_i(\hat{r}_{in}^\pm)$ in (3.2). Let $\hat{\phi}_n = \hat{d}_{2n} \hat{\sigma}_{2n}^2 / (\hat{d}_{1n} \hat{\sigma}_{1n}^2)$ and $\hat{\xi}_n = \hat{d}_{2n} / \hat{d}_{1n}$. Then it is not difficult to show that

$$\hat{\phi}_n \rightarrow_p \phi, \quad \hat{\xi}_n \rightarrow_p \xi \quad \text{and} \quad \frac{\hat{d}_{1n}}{\hat{\sigma}_{1n}^2} \rightarrow_p \frac{d_1}{\sigma_{10}^2}. \tag{4.14}$$

In practice, we do not know whether $r_{20} > r_{10}$. If $\hat{r}_{2n} > \hat{r}_{1n}$, we use (4.13) to obtain consistent estimates for (4.12). Otherwise, based on the expression of A_{1t} in (4.8) when $r_{10} > r_{20}$, we replace $\hat{\Phi}_{1n} - \hat{\Psi}_{2n}$ with $\hat{\Psi}_{1n} - \hat{\Phi}_{2n}$ and interchange the positions of \hat{r}_{2n} and \hat{r}_{1n} in (5.13). Thus, we can use $T_{\hat{\phi}_n, \hat{\xi}_n}$ to make statistical inferences for the change-point k_0 . Note that $T_{\hat{\phi}_n, \hat{\xi}_n}$ is asymmetric and differs from the symmetric T , which is commonly used to approximate the distribution of the estimated change-point in the literature. See Bai (1994), Chong (2001), and Ling (2016), among others.

5. Likelihood-ratio-based Inference for Thresholds and Change-points

The simulation method of Li and Ling (2012) for inferences of thresholds may not be accurate when the threshold effect is small. At the same time, the confidence interval based on the approximating method in Section 4 has a coverage rate below the nominal level when the threshold/structural change effect is large. See the discussion in Hansen (2000) for the threshold scenario, and those in Elliott and Müller (2007), Eo and Morley (2015), and Elliott, Müller and Watson (2015) for the change-point scenario, and they also commented that the likelihood-ratio test is asymptotically pivotal when the threshold/structural change effect is small and the confidence region based on the inverted likelihood-ratio test is asymptotically valid, even if the threshold/structural change effect is relatively large. In the regression model for the i.i.d. data, the nonparametric approach of Yu (2015) seems to work well when the threshold effect is relatively strong, but undercovers when the threshold effect is weak. However, it is not clear whether his method can be applied to a threshold model with time-series dependence. In this section, we consider likelihood-ratio-based confidence sets for the thresholds and the change-change, respectively.

We first investigate the likelihood-ratio test statistics for the thresholds. Following Hansen (2000), we consider the likelihood-ratio statistic $LR_{in}(r)$ for

$H_{i0} : r = r_{i0}$ as

$$LR_{in}(r) = \frac{1}{\hat{\sigma}_{in}^2} [S_{in}(\hat{\theta}_{in}(r, \hat{k}_n), r, \hat{k}_n) - S_{in}(\hat{\theta}_{in}, \hat{r}_{in}, \hat{k}_n)], \quad i = 1, 2, \quad (5.1)$$

where $\hat{\sigma}_{in}^2$ and S_{in} are defined in (3.1) and (2.3), respectively. Under H_{i0} , by (3.4), (3.5), and (5.1), we have

$$LR_{in}(r_{i0}) = \frac{1}{\hat{\sigma}_{in}^2} \max_{z \in R} \left[-\tilde{S}_{in} \left(k_0, \theta_{i0}, r_{i0}, r_{i0} + \frac{z}{n} \right) \right] + o_p(1), \quad (5.2)$$

where $\tilde{S}_{in}(k_0, \theta_{i0}, r_{i0}, r)$ are defined in (3.6) and (3.7) for $i = 1, 2$, respectively. Using a similar argument to that in Section 4.1, we use \bar{S}_{in} in (4.1)–(4.2) to approximate \tilde{S}_{in} for $i = 1, 2$, respectively. Using Theorem 4, it is not difficult to show that

$$\begin{aligned} \frac{1}{\hat{\sigma}_{in}^2} \max_{z \in R} \left[-\bar{S}_{in} \left(k_0, \theta_{i0}, r_{i0}, r_{i0} + \frac{z}{n} \right) \right] &= \frac{1}{\hat{\sigma}_{in}^2} \max_{v \in R} \left[-\bar{S}_{in} \left(k_0, \theta_{i0}, r_{i0}, r_{i0} + \frac{v}{n^{1-2\beta}} \right) \right] \\ &\Rightarrow \max_{v \in R} [2B(v) - |v|]. \end{aligned} \quad (5.3)$$

Then, if $\Psi_{i0} - \Phi_{i0} \approx \delta_{in}$, we can use the following approximation:

$$LR_{in}(r_{i0}) \approx_d \Delta := \max_{v \in R} [2B(v) - |v|], \quad (5.4)$$

where the distribution of Δ is $P(\Delta \leq x) = (1 - e^{-x/2})^2$; see Hansen (2000). From (5.4), we find that the asymptotic distribution of $LR_{in}(r_{i0})$ is free of nuisance parameters because the errors $\{\varepsilon_t\}$ are homoskedastic on each segment, by Assumption 1; see Theorem 2 in Hansen (2000) and the discussion thereafter. We use the distribution of Δ to solve for the critical value $c_{1-\alpha}$ (e.g., $\alpha = 0.05$), and a $1 - \alpha$ likelihood-ratio-based confidence set for r_{i0} is given by

$$\Gamma_{1-\alpha}^i = \{r : LR_{in}(r) \leq c_{1-\alpha}\}. \quad (5.5)$$

Next, we study the likelihood-ratio-based confidence set for the change-point. Following Eo and Morley (2015), we define the likelihood-ratio test statistic for $H_0 : k = k_0$ as follows:

$$LR_n(k) = S_n \left(\hat{\theta}_{1n}(\hat{r}_{1n}, k), \hat{\theta}_{2n}(\hat{r}_{2n}, k), \hat{r}_{1n}, \hat{r}_{2n}, k \right) - S_n \left(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n \right), \quad (5.6)$$

where S_n is defined as in (2.4). From our estimation procedure, the second term of (5.6) is $S_n(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, \hat{k}_n) = \min_{1 \leq k < n} S_n(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, k)$. Then, under H_0 , and using the results in Sections 3–4 and (S3.4) in the Supplementary Material, it is not difficult to show that

$$LR_n(k_0) = \max_{1 \leq k < n} \left[S_n \left(\hat{\theta}_{1n}(\hat{r}_{1n}, k_0), \hat{\theta}_{2n}(\hat{r}_{2n}, k_0), \hat{r}_{1n}, \hat{r}_{2n}, k_0 \right) \right]$$

$$\begin{aligned}
 & -S_n \left(\hat{\theta}_{1n}, \hat{\theta}_{2n}, \hat{r}_{1n}, \hat{r}_{2n}, k \right) \\
 &= \max_{1 \leq k < n} \left\{ I(k < k_0) \sum_{t=k+1}^{k_0} [\ell_t(\theta_{10}, r_{10}) - \ell_t(\theta_{20}, r_{20})] \right. \\
 & \quad \left. + I(k \geq k_0) \sum_{t=k_0+1}^k [\ell_t(\theta_{20}, r_{20}) - \ell_t(\theta_{10}, r_{10})] \right\} + o_p(1) \\
 & \triangleq \max_{1 \leq k < n} \{-W_n(k)\} + o_p(1). \tag{5.7}
 \end{aligned}$$

Without loss of generality, we assume $r_{20} > r_{10}$. Later, we show that this assumption does not affect our limiting theory. Using a similar argument to that in Section 4.2, we use $\bar{W}_n(k)$ in (4.9) to approximate $W_n(k)$ in (5.7). We only consider the case when $k < k_0$ in (4.9) because the other case is similar. From (4.10) and Assumption 7, we first define

$$\bar{A}_t = c'_3 Z_{t-1} I(q_{t-1} > r_{20}) + c'_4 Z_{t-1} I(q_{t-1} \leq r_{10}) + c'_5 Z_{t-1} I(r_{10} < q_{t-1} \leq r_{20}) \tag{5.8}$$

and, hence, $\bar{A}_{1t} = n^{-\beta} \bar{A}_t$. Under Assumption 7, we have

$$\begin{aligned}
 \max_{1 \leq k < n} \sum_{t=k+1}^{k_0} (-\bar{A}_{1t}^2 - 2\bar{A}_{1t}\varepsilon_t) &= \max_{s \in R} \sum_{t=k_0 + [n^{2\beta}s] + 1}^{k_0} (-\bar{A}_{1t}^2 - 2\bar{A}_{1t}\varepsilon_t) \\
 &\Rightarrow \max_{s \in R} \left[2\sqrt{E\bar{A}_t^2 \sigma_{10}^2} B(s) - E\bar{A}_t^2 |s| \right] \\
 &=_d \max_{v \in R} \{ \sigma_{10}^2 [2B(v) - |v|] \}, \tag{5.9}
 \end{aligned}$$

where we make a change of variables $s = v\sigma_{10}^2/E\bar{A}_t^2$ in the last step of (5.9). Then, we conclude that

$$\max_{1 \leq k < n} \{-\bar{W}_n(k)\} \Rightarrow \tilde{\Delta} = \max_v \begin{cases} \sigma_{10}^2 [2B(v) - |v|] & \text{for } v \in (-\infty, 0), \\ \sigma_{20}^2 [2B(v) - |v|] & \text{for } v \in [0, \infty), \end{cases} \tag{5.10}$$

and

$$LR_n(k_0) \approx_d \tilde{\Delta}, \tag{5.11}$$

where the distribution function of $\tilde{\Delta}$ is

$$P(\tilde{\Delta} \leq x) = \left(1 - \exp\left(-\frac{x}{2\sigma_{10}^2}\right) \right) \left(1 - \exp\left(-\frac{x}{2\sigma_{20}^2}\right) \right); \tag{5.12}$$

see, for example, Eo and Morley (2015). Then, we use (5.12) to solve for the critical value $\tilde{c}_{1-\alpha}$ and a $1 - \alpha$ likelihood-ratio-based confidence set for k_0 is given by

$$C_{1-\alpha} = \{k : LR_n(k) \leq \tilde{c}_{1-\alpha}\}. \tag{5.13}$$

In general, the different scaling factors σ_{10}^2 and σ_{20}^2 make the distribution of (5.12) asymmetric. In practice, σ_{10}^2 and σ_{20}^2 are replaced by consistent estimates from (3.1), and the calculation of the critical value using (5.12) is straightforward.

6. Simulation Study

This section examines the performance of our asymptotic results in finite samples using Monte Carlo experiments; all tables are provided in the Supplementary Material. We use sample sizes of $n = 400, 800,$ and $1,200$ with true change-points $k_0 = 200, 400,$ and $600,$ respectively. The data are generated from the following TAR(1) model with a change-point k :

$$y_t = \begin{cases} (\mu_1 + \phi_1 y_{t-1})I(y_{t-1} > r_1) \\ + (\nu_1 + \psi_1 y_{t-1})I(y_{t-1} \leq r_1) + \varepsilon_t, & \text{if } t \leq k, \\ (\mu_2 + \phi_2 y_{t-1})I(y_{t-1} > r_2) \\ + (\nu_2 + \psi_2 y_{t-1})I(y_{t-1} \leq r_2) + \varepsilon_t, & \text{if } t > k, \end{cases} \quad (6.1)$$

with the true values given as follows:

$$\begin{aligned} (\theta'_{10}, r_{10}) &= (-1, -0.6, 1, 0.4, 0.8) - \gamma(-1, -1, 1, 1, 0), \\ (\theta'_{20}, r_{20}) &= (-0.8, -0.9, 0.7, 0.6, 0.5) - \gamma(-1, -1, 1, 1, 0), \end{aligned}$$

and $\gamma = 0, 0.2,$ and $0.4,$ respectively, where $\varepsilon_t \sim N(0, 1)$; that is, $\sigma_{10} = \sigma_{20} = 1$. Clearly, the AR functions are not continuous over two thresholds $\{0.8, 0.5\}$ in all cases. We use 1,000 replications for each case. Table S1 summarizes the averages of the bias, empirical standard deviation (ESD), asymptotic standard deviation (ASD), and estimated asymptotic standard deviation (EASD) when $\gamma = 0$. The results are similar for the other cases and, hence, are not reported. Here, the ASDs of $\hat{\theta}_{in}$ are computed using the true $\sigma_{i0}^2 \Sigma_i$ in Theorem 2, where Σ_i is calculated from (3.2) and (3.3) using the true thresholds r_{i0} and the change-point k_0 . The ASDs of \hat{r}_{in} are obtained using the simulation method in Section 4 of Li and Ling (2012) with the true τ_0 in Theorem 3. The EASDs of $\hat{\theta}_{in}$ are computed using the estimators in (3.1)–(3.3), and the EASDs of \hat{r}_{in} are obtained using the simulation method of Li and Ling (2012), replacing τ_0 by $\hat{\tau}_n$ in Theorem 3. Table S1 shows that the bias is very close to zero for large n , but that it is not strictly decreasing as n increases because the average of the empirical bias also depends on the variance of the estimator. In addition, the larger the sample size, the closer the ESDs, ASDs, and EASDs are, on the whole. We also find that the convergence rate of the thresholds is n ; for example, the ESDs of \hat{r}_{in} for the sample of size 800 are half of those of the sample of size 400, and the ESDs of \hat{r}_{in}

in sample of size 1,200 are one-third of those of the sample of size 400, for $i = 1, 2$. Similarly, we find that the convergence rates of the other parameters are lower than those of the thresholds. These findings are similar to those reported in Li and Ling (2012). Furthermore, Table S1 shows that all estimated thresholds have a negative bias. This is because we used the left end-point of the interval on which (2.4) achieves its minimum. This negative bias can be overcome by using the middle-point; see Yu (2012) and Yu (2015). Note that we do not pursue this issue here because the bias is negligible for a large sample size and because we use the left end-point $M_-^{(i)}$ to make statistical inferences in Theorem 3 and the simulation studies below.

We now examine the coverage probabilities of r_{i0} and the performance of the approximating distributions and the likelihood-ratio-based confidence sets in Section 4.1 and Section 5, respectively. To do so, we first simulate the quantiles of $M_-^{(1)}$ and $M_-^{(2)}$ with 10,000 replications. Based on these quantiles and those in Table 1 of Hansen (1997), the coverage probabilities of r_{i0} are reported in Table S2 when $\gamma = 0, 0.2, \text{ and } 0.4$, where $\|\delta_{in}\| = \|\Psi_{i0} - \Phi_{i0}\|$, for $i = 1, 2$. The results show that the coverage probabilities based on $M_-^{(1)}$ and $M_-^{(2)}$ are relatively accurate in all cases, on the whole, but tend to undercover when the threshold effect $\|\delta_{in}\|$ is small. The coverage probabilities based on the quantiles of T are often relatively worse than those of the other two methods, especially when $\gamma = 0$ (i.e., $\|\delta_{in}\|$ is very large), but their accuracy improves as $\|\delta_{in}\|$ becomes smaller. The coverage probabilities based on LR_{in} tend to exceed the nominal levels for large threshold effects $\|\delta_{in}\|$ and decrease with $\|\delta_{in}\|$; similar results can be found in Hansen (2000), who also found that the likelihood-ratio-based method may undercover for very small threshold effects and small sample sizes. Overall, when the threshold effect is large, the simulation method of Li and Ling (2012) is relatively accurate, the approximating method in Section 4.1 undercovers and the likelihood-ratio-based method overcovers and is quite conservative. When the threshold effect is small, the simulation method of Li and Ling (2012) undercovers and the methods based on the approximation and the likelihood-ratio may be more accurate. In practice, we do not know the exact magnitude of the threshold effect. Thus, we recommend using the likelihood-ratio method and the simulation method when making statistical inferences because they tend to be more accurate.

We next examine the performance of the approximating distribution in Theorem 5 and the likelihood-ratio-based confidence sets in Section 5 for the estimated change-points in finite samples. We examine both large and small structural

change effects in our experiments. To determine the finite-sample performance when the structural change effect is relatively large, we first fix the true parameter $(\theta'_{10}, r_{10}) = (-1, -0.6, 1, 0.4, 0.8)$ and let the true parameter

$$(\theta'_{20}, r_{20}) = (-\theta'_{10}, r_{10}) - \tilde{\gamma}_1(1, 1, -1, -1, 1), \quad (6.2)$$

with $\tilde{\gamma}_1 = 0, 0.1, 0.2,$ and 0.4 . It is easy to see that the structural change effects are relatively large for the choices of $\tilde{\gamma}_1$ because the vectors θ_{10} and θ_{20} have different signs. When $\tilde{\gamma}_1$ increases, the structural change effect decreases. The number of replications is 1,000 for each case in this experiment. Table S3 summarizes the mean, ESD, ASD, and the estimators of d_{i0} , ϕ , and ξ for (6.2). Here, $\kappa_n = \|\Phi_{20} - \Phi_{10}\| + \|\Psi_{20} - \Phi_{10}\| + \|\Psi_{10} - \Phi_{20}\|$, representing the structural change effect, the ESD is calculated based on the 1,000 estimators, the ASD is computed using the approximating distribution in (5.9), and \hat{d}_{in} is based on (4.13). From the expressions of $\hat{\phi}_n$ and $\hat{\xi}_n$ in (4.14), our simulation results show that $\hat{\phi}_n$ and $\hat{\xi}_n$ are almost the same, because we assume $\sigma_{10} = \sigma_{20} = 1$ in our experiment. Using the true parameters and (4.12) to obtain d_{i0} , ϕ , and ξ , the results are similar to those in Table S3 and, hence, are not reported here. For reference purposes, we report only the values of $(\hat{d}_{10}, \hat{d}_{20})$ corresponding to different $\tilde{\gamma}_1$, which are calculated using (4.12) with the true parameters and the sample estimates of $M_i(r_{i0}^\pm)$. From Table S3, we find that the means of the estimated change-points are close to the true change-points in all cases. The ASDs are smaller than the ESDs in all cases, and tend to become closer as the structural change effect decreases. This is reasonable because our approximating distribution is based on a small change effect. We also find that the estimated d_{i0} , ϕ , and ξ are almost the same for different sample sizes with fixed $\tilde{\gamma}_1$.

Based on the results in Table S3 and the density function of $T_{\phi, \xi}$ in Bai (1997), we examine the coverage probabilities of the estimated change-points. The results are reported in Table S4. From Table S4, we can see that the likelihood-ratio-based confidence sets of LR_n overcover at all three nominal levels, while the approximating distribution $T_{\phi, \xi}$ undercovers significantly for relatively large structural change effects in (6.2). These findings are similar to those in Eo and Morley (2015), who found that the coverage rate based on the likelihood-ratio approach is more precise than those of Bai (1997) and Elliott and Müller (2007).

Now, we study the finite-sample performance when the structural change effect is relatively small. We set (θ'_{10}, r_{10}) as before, and let the true parameter

$$(\theta'_{20}, r_{20}) = (\theta'_{10}, r_{10}) - \tilde{\gamma}_2(-1, -1, 1, 1, 1), \quad (6.3)$$

with $\tilde{\gamma}_2 = 0.1, 0.2, 0.3,$ and 0.5 . It is easy to see that the structural change

effects are relatively small for the choices of $\tilde{\gamma}_2$. As $\tilde{\gamma}_2$ increases, the structural change effect increases. Table S5 summarizes the results. The results shown in Table S5 are similar to those in Table S3, although the ESDs and ASDs are larger than those reported in Table S3. This is reasonable because it is not easy to locate the change-point when the structural change effect is small. Based on the estimators in Table S5 and the density function of $T_{\phi,\xi}$ in Bai (1997), we examine the coverage probabilities of the estimated change-points. The results are reported in Table S6, which shows that the likelihood-ratio-based confidence sets of LR_n undercover at all three nominal levels. In addition, the approximating distribution $T_{\phi,\xi}$ outperforms the likelihood-ratio-based method for relatively small structural change effects in (6.3).

Overall, when the structural change effect is large, the likelihood-ratio-based confidence sets of LR_n overcover and are somewhat conservative. When the structural change effect is very small, the approximating method in Section 5.2 may be more accurate. In practice, it is not easy to evaluate the magnitude of the change, especially when the two thresholds are different. Based on our limited simulation experience, we recommend using the likelihood-ratio-based method because it exhibits better performance in general.

7. A Real-Data Example

Yau, Tang and Lee (2015) applied the TAR model with structural breaks to U.S. GNP data and found breaks associated with substantial changes in the U.S. economy. This section uses model (2.1) with a structural change to study a long time series of annual tree-ring widths (Figure S1(a)). All measurements are taken from a Qilian Juniper tree in the northeastern Tibetan Plateau of China. The time series spans the period from 1079 to 2009 and it was obtained from the NOAA paleoclimatology database, available at <https://www.ncdc.noaa.gov/paleo/study/16645>. Tree rings provide important records of past climates and, hence, can be used to study climate change; see Cook (1985). In the past decades, TAR models have been recognized as an important nonlinear time series model for studying climate changes. Ellis and Post (2004) and Tong (2011) demonstrate the merits of using TAR models rather than linear AR models.

Let x_t denote the original data and y_t denote the continuously annualized average growth rate; that is, $y_t = \log(x_t/x_{t-1})$. Figure S1(b) shows the time plot of y_t . We can see large fluctuations in Figure S1(a)–(b). Figure S2 shows the autocorrelation function (ACF) and the partial ACF (PACF) of $\{y_t\}$, which

indicate that $\{y_t\}$ is a sequence of dependent time series. Next, we explain how to build a TAR model with a structural change to $\{y_t\}$.

Step 1. We first perform the threshold nonlinearity test. There are many ways to do this in the literature; see Tsay (1989) and Chan (1991), among others. Here, we adopt the likelihood-ratio approach of Chan (1991), and use the corresponding package TSA in R. See also Cryer and Chan (2008) for details. For each null model AR(p), we choose the threshold variable $q_{t-1} = y_{t-d}$ with $1 \leq d \leq p$. Table S7 reports the p -values when performing the threshold nonlinearity test under possible linear AR models. The choice of the AR order p is based on the PACF in Figure S2(b), as suggested by Tsay (1989), when testing for threshold nonlinearity. From Table S7, we can see that most of the p -values are close to zero except for some cases when $d = 2$, where the p -values are only slightly larger than 5%. We conclude that there is likely a threshold effect in the data $\{y_t\}$. In other words, it is better to use a threshold model to fit the data than to use a pure AR model.

Step 2. We now fit a TAR(p) model to $\{y_t\}$ with a threshold y_{t-d} , where $1 \leq d \leq p$. Define

$$\text{AIC}(p, d) = n \log(\hat{\sigma}_n^2) + 2(p + 1) \quad \text{and} \quad \text{BIC}(p, d) = n \log(\hat{\sigma}_n^2) + (p + 1) \log(n), \quad (7.1)$$

where $\hat{\sigma}_n^2$ is defined similarly to (3.1), using the whole sample. Here, (7.1) is slightly different from that in Li and Ling (2012) because we do not allow a threshold effect in the error term of (2.1). To simplify the model, based on the PACF in Figure S2(b), we set $1 \leq p \leq 12$ and $1 \leq d \leq p$. The results of the AICs and BICs are summarized in Table S8. From Table S8, we can see that the AIC selects the model TAR(12) with $d = 10$, and the BIC selects TAR(8) with $d = 8$. For ease of exposition, we choose the simpler model TAR(8). Thus, the fitted model is as follows:

$$y_t = (\mu + \sum_{i=1}^8 \phi_i y_{t-i}) I(y_{t-8} > 0.1252) + (\nu + \sum_{i=1}^8 \psi_i y_{t-i}) I(y_{t-8} \leq 0.1252) + 0.247 u_t, \quad (7.2)$$

where the standard deviation 0.247 is calculated using the residual sum of squares and the other parameters are summarized in Table S9. We now use the portmanteau test in (15.8.3) of Cryer and Chan (2008, p.412) to check whether model (7.2) is adequate. Figure S3 displays the p -values of the test, showing that they

are all larger than 5%. Therefore, model (7.2) is adequate for the data $\{y_t\}$.

Step 3. We use the Sup-likelihood-ratio test statistic in Andrews (1993) (see also Davis, Huang and Yao (1995)), $\sup_{\tau \in [0.05, 0.95]} LR(\tau)$, to test whether a structural change exists in model (7.2). Because the estimate of the threshold is super-efficient with convergence rate of n under the null hypothesis of no change-point, the limiting distribution in Andrews (1993) or Davis, Huang and Yao (1995) is still applicable to the TAR(p) model with degrees of freedom $2(p+1)$. We find that $\sup_{\tau \in [0.1, 0.9]} LR(\tau) = 89.77$, which exceeds the critical value of 46.69 at the 0.01 significance level; see Table 1 in Andrews (1993). Hence, model (7.2) most likely has a structural change during this period. Note that this does not contradict the finding in Step 2 that model (7.2) is adequate for the data because the likelihood-ratio test uses a different criterion for the model selection in this step.

Step 4. Based on the estimation procedure in Section 2, a TAR(8) model with $d = 8$ and a structural change is used to fit the data. The result is as follows:

$$y_t = \begin{cases} (\mu_1 + \sum_{i=1}^8 \phi_{1i} y_{t-i}) I(y_{t-8} > -0.1744) \\ + (\nu_1 + \sum_{i=1}^8 \psi_{1i} y_{t-i}) I(y_{t-8} \leq -0.1744) + 0.241 u_t, & t \leq 578, \\ (\mu_2 + \sum_{i=1}^8 \phi_{2i} y_{t-i}) I(y_{t-8} > 0.0972) \\ + (\nu_2 + \sum_{i=1}^8 \psi_{2i} y_{t-i}) I(y_{t-8} \leq 0.0972) + 0.232 u_t, & t > 578, \end{cases} \quad (7.3)$$

where the two standard deviations 0.241 and 0.232 are calculated from (3.1) and the other coefficients are reported in Table S10. The standard deviations are given in parentheses, and some parameters are not significant at the 5% level. We use the method of Cryer and Chan (2008, p.412) for the model diagnostic checking. Figure S4 displays the p -values of the test in the two segments of (7.3), and again shows that model (7.3) is adequate for $\{y_t\}$. Note that there appear to be at least two change-points in the data from Figure S1(b), but that our method finds the most visually obvious point. For multiple change-points, interested readers may consult the approach provided in Section 2 on page 9.

From the model, we can see that almost all of the AR coefficients are negative. This is reasonable because a higher growth rate in the current year will result in a lower growth rate in the subsequent year, and vice versa. Furthermore, the AR coefficients before and after the change-point $\hat{k}_n = 578$ change significantly, and almost all of their absolute values after the change-point are larger than their counterparts before the change-point. Therefore, the dependence of the growth rates grows stronger after the change-point. Based on the methods of Li and Ling (2012), the approximation in Section 4.1, and the

likelihood-ratio method in Section 5, the 95% confidence intervals for r_{10} are $[-0.189, -0.166]$, $[-0.213, -0.135]$, and $[-0.174, -0.014]$, respectively, and those of r_{20} are $[0.084, 0.119]$, $[-0.179, 0.374]$, and $[-0.028, 0.301]$, respectively. We can see that the simulation method provides rather tight confidence intervals for the thresholds, but the other two methods tend to yield wider confidence intervals. The likelihood-ratio-based method gives similar intervals to those of the simulation one for r_{10} , but provides a much wider interval for r_{20} . Thus, we suggest using the intervals produced by the simulation-based method because they are relatively tight in this case. The 95% confidence interval of k_0 based on the approximation method is $[563, 585]$. This tight interval indicates that the estimator \hat{k}_n is very accurate.

After checking the data, we find that \hat{k}_n represents the year 1656. The 95% confidence intervals show that there was most likely a significant change in the climate in the period 1641–1663. From Figure S1(a), we can see that the growth of the tree rings declined from the 1600s onwards, indicating that temperatures might have changed rapidly around this period. Historical records indicate that there were many disasters, such as dry weather and crop failures, in this period, and that the Ming dynasty collapsed in 1644. There are no climate records for this period in China. Many historians suspect that the bad weather was the result of climate change. Our findings provide evidence that supports this view and may be useful for future study of Chinese history.

In the Supplementary Material, we further examine some of the steps in this section and demonstrate the merits of model (7.3) with a change-point by focusing on the forecasting errors. See Section S5 in the Supplementary Material for details.

Supplementary Materials

Owing to space constraints, we provide a new SLLN, the proofs of all theorems, the effect of the initial values, and some tables and figures in the Supplementary Material.

Acknowledgment

We are grateful to the editors and the anonymous referees for their insightful comments and suggestions that have substantially improved the presentation and the content of this paper. This project is supported by Hong Kong Research Grants Commission Grants GRF 16500915, GRF 16307516, GRF 16500117, and

the fifth batch of excellent talent support program for Chongqing Colleges and Universities.

References

- Andrews, D. W. (1993). Tests for parameter instability and structural change with unknown change point. *Econometrica* **61**, 821–856.
- Aue, A., Hörmann, S., Horváth, L. and Reimherr, M. (2009). Break detection in the covariance structure of multivariate time series models. *The Annals of Statistics* **37**, 4046–4087.
- Bai, J. (1994). Least squares estimation of a shift in linear processes. *Journal of Time Series Analysis* **15**, 453–472.
- Bai, J. (1995). Least absolute deviation estimation of a shift. *Econometric Theory* **11**, 403–436.
- Bai, J. (1997). Estimation of a change point in multiple regression models. *Review of Economics and Statistics* **79**, 551–563.
- Bai, J., Lumsdaine, R. L. and Stock J. H. (1998). Testing for and dating common breaks in multivariate time series. *The Review of Economic Studies* **65**, 395–432.
- Bai, J. and Perron, P. (1998). Estimating and testing linear models with multiple structural changes. *Econometrica*, 47–78.
- Bai, J., and Perron, P. (2003). Computation and analysis of multiple structural change models. *Journal of Applied Econometrics* **18**, 1–22.
- Chan, K. -S. (1989). A note on the geometric ergodicity of a Markov chain. *Advances in Applied Probability* **21**, 702–704.
- Chan, K. S. (1991). Percentage points of likelihood ratio tests for threshold autoregression. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)* **53**, 691–696.
- Chan, K. -S. (1993). Consistency and limiting distribution of the least squares estimator of a threshold autoregressive model. *The Annals of Statistics* **21**, 520–533.
- Chan, K. -S. and Tong, H. (1985). On the use of the deterministic Lyapunov function for the ergodicity of stochastic difference equations. *Advances in Applied Probability* **17**, 666–678.
- Chan, K. -S. and Tsay, R. S. (1998). Limiting properties of the least squares estimator of a continuous threshold autoregressive model. *Biometrika* **85**, 413–426.
- Chan, N. H. and Kutoyants, Y. A. (2010). Recent developments of threshold estimation for nonlinear time series. *Journal of the Japan Statistical Society* **40**, 277–303.
- Chan, N. H., Yau, C. Y. and Zhang, R. -M. (2014). Group Lasso for structural break time series. *Journal of the American Statistical Association* **109**, 590–599.
- Chan, N. H., Yau, C. Y. and Zhang, R. -M. (2015). LASSO estimation of threshold autoregressive models. *Journal of Econometrics* **189**, 285–296.
- Chong, T. T. -L. (2001). Structural change in AR(1) models. *Econometric Theory* **17**, 87–155.
- Cook, E. (1985). *A Time Series Analysis Approach to Tree Ring Standardization*. Thesis. The University of Arizona.
- Cryer, D. and Chan, K. S. (2008). *Time Series Analysis With Applications in R*. Springer.
- Csörgö, M. and Horváth, L. (1997). *Limit Theorems in Change-Point Analysis*, Volume 18. John Wiley & Sons Inc.
- Davis, R. A., Huang, D. and Yao, Y. -C. (1995). Testing for a change in the parameter values and order of an autoregressive model. *The Annals of Statistics* **23**, 282–304.

- Davis, R. A., Lee, T. C. M. and Rodriguez-Yam, G. A. (2006). Structural break estimation for nonstationary time series models. *Journal of the American Statistical Association* **101**, 223–239.
- Elliott, G. and Müller, U. K. (2007). Confidence sets for the date of a single break in linear time series regressions. *Journal of Econometrics* **141**, 1196–1218.
- Elliott, G., Müller, U. K. and Watson, M. W. (2015). Nearly optimal tests when a nuisance parameter is present under the null hypothesis. *Econometrica* **83**, 771–811.
- Ellis, A. M. and Post, E. (2004). Population response to climate change: linear vs. non-linear modeling approaches. *BMC Ecology* **4**, 2.
- Eo, Y. and Morley, J. (2015). Likelihood-ratio-based confidence sets for the timing of structural breaks. *Quantitative Economics* **6**, 463–497.
- Gao, J., Tjøstheim, D. and Yin, J. (2013). Estimation in threshold autoregressive models with a stationary and a unit root regime. *Journal of Econometrics* **172**, 1–13.
- Gao, Z., Ling, S. and Tong, H. (2018). Tests for TAR models vs. STAR models—a separate family of hypotheses approach. *Statistica Sinica* **28**, 2857–2883.
- Haavelmo, T. (1944). The probability approach in econometrics. *Econometrica*, 1–118.
- Hansen, B. E. (1997). Inference in tar models. *Studies in Nonlinear Dynamics & Econometrics* **2**, 119–131.
- Hansen, B. E. (2000). Sample splitting and threshold estimation. *Econometrica* **68**, 575–603.
- Hidalgo, J. and Seo, M. H. (2013). Testing for structural stability in the whole sample. *Journal of Econometrics* **175**, 84–93.
- Hinkley, D. V. (1970). Inference about the change-point in a sequence of random variables. *Biometrika* **57**, 1–17.
- Kejriwal, M. and Perron, P. (2008). The limit distribution of the estimates in cointegrated regression models with multiple structural changes. *Journal of Econometrics* **146**, 59–73.
- Li, D. and Ling, S. (2012). On the least squares estimation of multiple-regime threshold autoregressive models. *Journal of Econometrics* **167**, 240–253.
- Li, D., Ling, S. and Li, W. K. (2013). Asymptotic theory on the least squares estimation of threshold moving-average models. *Econometric Theory* **29**, 482–516.
- Li, D., Ling, S. and Zhang R. M. (2016). On a threshold double autoregressive model. *Journal of Business & Economic Statistics* **34**, 68–80.
- Ling, S. (2007). Testing for change points in time series models and limiting theorems for ned sequences. *The Annals of Statistics* **35**, 1213–1237.
- Ling, S. (2016). Estimation of change-points in linear and nonlinear time series models. *Econometric Theory* **32**, 402–430.
- Liu, W., Ling, S. and Shao, Q. -M. (2011). On non-stationary threshold autoregressive models. *Bernoulli* **17**, 969–986.
- Meyn, S. P. and Tweedie, R. L. (2009). *Markov Chains and Stochastic Stability*. Cambridge Univ Press.
- Picard, D. (1985). Testing and estimating change-points in time series. *Advances in Applied Probability*, 841–867.
- Qian, J. and Su, L. (2014). Structural change estimation in time series regressions with endogenous variables. *Economics Letters* **125**, 415–421.
- Rosenblatt, M. (1956). A central limit theorem and a strong mixing condition. *Proceedings of*

- the National Academy of Sciences of the United States of America* **42**, 43–47.
- Saikkonen, P., Lütkepohl, H. and Trenkler, C. (2006). Break date estimation for var processes with level shift with an application to cointegration testing. *Econometric Theory* **22**, 15–68.
- Seo, M. H. and Linton, O. (2007). A smoothed least squares estimator for threshold regression models. *Journal of Econometrics* **141**, 704–735.
- Shao, X. and Zhang, X. (2012). Testing for change points in time series. *Journal of the American Statistical Association* **105**, 1228–1240.
- Tong, H. (1978). On a threshold model. In *Pattern Recognition and Signal Processing* (Edited by C. H. Chen), 101–141. Sijthoff & Noordhoff.
- Tong, H. and Lim, K. S. (1980). Threshold autoregression, limit cycles and cyclical data-with discussion. *Journal of the Royal Statistical Society, Series B (Statistical Methodology)* **42**, 245–292.
- Tong, H. (2011). Threshold models in time series analysis—30 years on. *Statistics and its Interface* **4**, 107–118.
- Tsay, R. S. (1989). Testing and modeling threshold autoregressive processes. *Journal of the American Statistical Association* **84**, 231–240.
- Yao, Y. -C. (1987). Approximating the distribution of the maximum likelihood estimate of the change-point in a sequence of independent random variables. *The Annals of Statistics* **15**, 1321–1328.
- Yau, C. Y., Tang, C. M. and Lee, T. C. (2015). Estimation of multiple-regime threshold autoregressive models with structural breaks. *Journal of the American Statistical Association* **110**, 1175–1186.
- Yu, P. (2012). Likelihood estimation and inference in threshold regression. *Journal of Econometrics* **167**, 274–294.
- Yu, P. (2015). Adaptive estimation of the threshold point in threshold regression. *Journal of Econometrics* **189**, 83–100.

Booth School of Business, University of Chicago , Chicago, IL 60637, USA.

E-mail: mazxgao@gmail.com

Department of Mathematics, Hong Kong University of Science and Technology Clear Water Bay, Hong Kong.

E-mail: maling@ust.hk

(Received July 2017; accepted January 2018)