

BAYESIAN MODELS FOR DETECTING DIFFERENCE BOUNDARIES IN AREAL DATA

Pei Li, Sudipto Banerjee, Timothy A. Hanson and Alexander M. McBean

*Medtronic Inc., University of California, Los Angeles,
University of South Carolina and University of Minnesota*

Abstract: With increasing accessibility to Geographical Information Systems (GIS) software, researchers and administrators in public health routinely encounter *areal* data compiled as *aggregates* over areal regions, such as counts or rates across counties in a state. Spatial models for areal data attempt to deliver smoothed maps by accounting for high variability in certain regions. Subsequently, inferential interest is focused upon formally identifying the “difference edges” or “difference boundaries” on the map that delineate adjacent regions with vastly disparate outcomes, perhaps caused by latent risk factors. We propose nonparametric Bayesian models for areal data that can formally identify boundaries between disparate neighbors. After elucidating these models and their estimation methods, we resort to simulation experiments to assess their effectiveness, and subsequently analyze Pneumonia and Influenza hospitalization maps from the SEER-Medicare program in Minnesota, where we detect and report highly disparate neighboring counties.

Key words and phrases: Areal data, conditional autoregressive model, difference boundary, Dirichlet process, stick-breaking process, wombling.

1. Introduction

With increasing accessibility to Geographical Information Systems (GIS), researchers and administrators in public health are increasingly encountering *areal* datasets that are *aggregated* as case counts or rates over *areal* units or regions (e.g. counties, census-tracts or ZIP codes). This is common practice in public health to protect patient privacy.

Statistical models for areal data can adjust for known causes of variability in the data and also for sparsely sampled regions by smoothing across and borrowing information from its spatial neighbors (see, e.g., Anselin (1988); Le Sage and Pace (2009); Banerjee, Carlin, and Gelfand (2004)). An especially pertinent issue is to ascertain statistically significant differences among neighboring regions, hence identifying the *spatial barriers* or *difference boundaries* that delineate them. Ultimately, the underlying influences responsible for these boundaries or barriers are typically of scientific and administrative interest. This ‘boundary’ detection problem is often referred to as “wombling”, after a foundational

article by Womble (1951). While statistical boundary analysis has been applied extensively to point-referenced and gridded (or lattice) data (see, e.g., Banerjee and Gelfand (2006)), formal statistical inference in areal contexts present unique challenges that we outline later.

Deterministic areal wombling is often carried out using algorithms (Jacquez and Greiling (2003a,b)) that are fast and straightforward to implement but fail to account for sources of uncertainty, such as extremes in counts and rates corresponding to thinly populated regions. Li, Banerjee, and McBean (2011) proposed statistical learning for boundaries using the Bayesian Information Criterion. In hierarchical model based approaches, Lu and Carlin (2005), Lu et al. (2007) and Ma, Carlin, and Banerjee (2010) investigated estimation of the adjacency matrix within a hierarchical framework using priors on the edges. However, inference from these models are usually highly sensitive to prior specifications on certain parameters.

Our primary contribution is a method to deliver inference for areally aggregated health outcome data, including assessment of difference boundaries, using classes of more flexible and robust non-parametric Bayesian hierarchical models. Section 2 offers a brief exposition to models for areally referenced count data. Section 3 elucidates the key issues in areal boundary analysis and our Bayesian nonparametric modeling approaches. Sections 4 and 5 discuss, respectively, a simulation study and the analysis of a Minnesota Pneumonia & Influenza (P & I) dataset to detect spatial health barriers between neighboring counties in Minnesota. Finally, Section 6 concludes the article with an eye towards future work.

2. Hierarchical Models for Areal Data

Areal data can be analyzed using Bayesian hierarchical models that incorporate geographical effects. For example, let Y_i (random) be the observed number of patients who underwent a specific preventive or clinical outcome in areal unit i , $i = 1, \dots, n$, and let E_i (fixed) be the expected number of outcomes for that unit. A commonly used likelihood is

$$Y_i \stackrel{ind.}{\sim} \text{Poisson}(E_i e^{\mu_i}), \quad i = 1, \dots, n, \quad (2.1)$$

where $\mu_i = \mathbf{x}_i' \boldsymbol{\beta} + \phi_i$ represents the log-relative risk, estimates of which are often based on the departures of observed from expected counts, \mathbf{x}_i includes explanatory, region-level covariates or predictors for region i and $\boldsymbol{\beta}$ are the corresponding regression coefficients.

The ϕ_i represent the *spatial random effect* associated with region i ; they are often modeled using *Markov random fields* (e.g., Cressie (1993);

Banerjee, Carlin, and Gelfand (2004, Chap. 3)) that imply a joint distribution for $\boldsymbol{\phi} = (\phi_1, \phi_2, \dots, \phi_n)'$:

$$\boldsymbol{\phi} \sim N_n(\mathbf{0}, \sigma^2 (D - \rho W)^{-1}) , \quad (2.2)$$

where N_n denotes the n -dimensional normal distribution, D is a $n \times n$ diagonal matrix with diagonal elements m_i equal to the number of neighbors of area i , and $W = \{w_{ij}\}$ is the adjacency matrix with $w_{ii} = 0$, and $w_{ij} = 1$ if i is adjacent to j and 0 otherwise. In the joint distribution (2.2), σ^2 is the spatial dispersion parameter, and ρ is a spatial autocorrelation parameter. We denote this distribution concisely as $CAR(\rho, \sigma^2)$. A sufficient condition for $D - \rho W$ to be positive definite is that $\rho \in (1/\lambda_{(1)}, 1)$, where $\lambda_{(1)}$ is the minimum eigenvalue of W (Banerjee, Carlin, and Gelfand (2004)).

The CAR model has been especially popular in Bayesian inference as its conditional specification is convenient for Gibbs sampling and MCMC schemes. The distribution in (2.2) reduces to the well-known intrinsic conditionally autoregressive (ICAR) prior if $\rho = 1$, or an independence model if $\rho = 0$. The ICAR model induces “local” smoothing by borrowing strength from the neighbors, while the independence model assumes independence of spatial rates and induces “global” smoothing. The smoothing parameter ρ in the CAR prior (2.2) controls the strength of spatial dependence among regions, though it is well-appreciated that a fairly large ρ may be required to deliver significant spatial correlation.

3. Bayesian Nonparametric Models for Areal Data

3.1. Modelling considerations for areal boundary analysis

Areal boundary analysis can be approached from different perspectives. For example, Li, Banerjee, and McBean (2011) treat the problem as one of statistical learning for the edges, where each model represents a different *boundary hypothesis*. More generally, one can consider models varying in their specification of the neighborhood matrix W that controls spatial smoothing. This requires sophisticated MCMC model composition or MC³ algorithms or other types of stochastic variable selection algorithms for selecting models (see, e.g., Hoeting et al. (1999)); these methods are computationally intensive in relatively large maps. Li et al. (2012) reformulate the problem as one of Bayesian hypothesis testing within a class of spatial moving average models and adjust multiple tests using false discovery rates. The method, though still computationally intensive is competitive, and provides a benchmark in our simulation studies. Another approach seeks to estimate the adjacency matrix within a hierarchical framework using priors on the adjacency relationships, see Ma, Carlin, and Banerjee (2010).

We explore an alternative stochastic mechanism that allows us detect difference boundaries by considering probabilities such as $P(\phi_i = \phi_j | i \sim j)$. Clearly, continuous priors for the ϕ_i 's do not work as they render $P(\phi_i = \phi_j | i \sim j) = 0$. A nonparametric Bayesian framework that models the spatial effects as almost surely discrete realizations of some distribution comes to mind – the Dirichlet process (Ferguson (1973)) presents itself as a natural choice, but how do we accommodate spatial (areal) dependence? We address this issue in the subsequent sections.

3.2. Dirichlet process mixture (DPM) models for clustered data

In the context of (2.1) and (2.2), a Dirichlet process mixture (DPM) prior specifies $\phi_i \sim G$, where $G \sim DP(\alpha, G_0)$ is some unknown distribution modeled as a Dirichlet process (DP) with baseline measure G_0 (e.g., Ferguson (1973)). Blackwell and MacQueen (1973) related the Dirichlet process to a generalized *Polya urn scheme* that leads to effective sampling strategies if given an explicit and simple prediction rule.

The stick-breaking representation of the DP (Sethuraman (1994)) says that a draw from the Dirichlet process can be written as $G(\cdot) = \sum_{i=1}^{\infty} p_i \delta_{\theta_i}(\cdot)$ *a.s.*, where δ_{θ_i} is the Dirac measure (point mass) located at θ_i , each θ_i is a random draw from the base distribution G_0 , and $p_i = V_i \prod_{l=1}^{i-1} (1 - V_l)$ with $p_1 = V_1$, where each $V_i \stackrel{i.i.d.}{\sim} \text{Beta}(1, \alpha)$. The p_i 's are called the “stick-breaking” weights (their infinite sum equals 1) and the θ_i 's are called atoms. In practice, the infinite sum is often replaced by the sum of the first N ($N \leq n$) terms, since the probability mass in each term decays rapidly. We can simply let $V_N = 1$ to truncate the sum to finite terms (Ishwaran and Zarepour (2000)). Many authors simply choose N to be a number large enough that there exists some empty components during the MCMC run or by examining the size of the last weight p_N under the prior. Following Reich and Fuentes (2007), we choose N according to the latter (see Section 4). For concerns regarding truncation bias, exact sampling can be executed using slice-sampling (Kalli, Griffin, and Walker (2011)).

The stick-breaking representation is an extremely rich framework that subsumes DP's and other extensions (e.g., MacEachern (2001)) such as Dependent Dirichlet processes (DDP)'s. In fact, for every conceivable joint distribution on the stick-breaking weights and the atoms, there is associated a stick-breaking stochastic process. Introducing dependence is now natural. For example, the DDP introduces dependence through the stick-breaking weights and the atoms. De Iorio et al. (2004) used the dependent Dirichlet process to define the desired dependence across the related random distributions in any ANOVA-type models. Gelfand, Kottas, and Maceachern (2005) used DDP on geostatistical data

and introduced spatial dependence through an underlying base measure, where $G \sim DP(\alpha, G_0^{(n)})$ and $G_0^{(n)}$ is a Gaussian process with a given covariance structure. Alternatively, Griffin and Steel (2006) proposed an order-based DDP which include dependence on predictors by permutation of elements in stick-breaking priors. Variants include kernel stick-breaking processes by Dunson and Park (2008) and the probit stick-breaking process by Chung and Dunson (2009). Spatial Dirichlet process (SDP) mixture models (Gelfand, Kottas, and Maceachern (2005)) are defined on a space of surfaces that yields almost surely discrete realizations with countable support. Duan, Guindani, and Gelfand (2007) extended the SDP by allowing different surface selection at different sites. Reich and Fuentes (2007) develop a spatial stick-breaking prior (SSB) to analyze hurricane surface wind fields.

Our need to move into DP's (and their extensions) is fundamental: to accommodate non-zero probability masses for spatial random effects without sacrificing richness for areal models is problematic otherwise. The local Dirichlet process (Chung and Dunson (2011)) offers an approach to the localized spatial "sharing" of atoms and weights that could conceivably be extended to the areal setting through a suitable definition of what a neighborhood is at each areal location; also see Theorem 4 in Dunson, Pillai, and Park (2007) for a related idea. The models we propose below correspond to a subclass of stick-breaking process priors that includes the DP and the SDP as special cases. In particular, we construct an areally-referenced stick-breaking process (ARSB) and an areally-referenced Dirichlet Process (ARDP) that will serve well for areal data allowing formal boundary analysis.

More recent developments in Dependent Product Partition Models (DPPM), which refers to classes of predictor-dependent product partition models, encourage clustering among subjects with like covariates. Both Müller, Quintana, and Rosner (2011) and Park and Dunson (2010) require a "similarity function" $g(x_1, \dots, x_k)$ that adjusts the (typically Dirichlet process) cohesion function, giving larger values for sets of covariates x_1, \dots, x_k that are "similar." Park and Dunson's (2010) model development proceeds through the consideration of similarity functions that place probability distributions on the covariates x_1, x_2, \dots , treating them as *continuous*; specifically, they consider the DPM of normals proposed by Müller, Erkanli, and West (1996). Müller, Quintana, and Rosner (2011) consider normal models for continuous covariates, and other choices for ordinal and nominal categorical predictors. For our model and application, we require a cohesion function that gives larger values for proximal counties, so our "predictor" is a categorical variable with spatial information. Here we propose incorporating areal spatial information directly into the stick-breaking weights (ARSB) or through a copula-type formulation (ARDP).

3.3. Areally-referenced spatial stick-breaking prior

We adapt the point-referenced spatial stick-breaking approach of Reich and Fuentes (2007) to areal data by incorporating spatial dependence in the DP by introducing additional weights that borrow strength across the neighbors using CAR priors (Section 2)

The spatial random effects are assigned a stick-breaking prior, whose weight parameters $p_{i1} = w_{i1}V_1, p_{ik} = w_{ik}V_k \prod_{l=1}^{k-1}(1 - w_{il}V_l), i = 1, \dots, n, k = 1, 2, \dots,$ depend not only on the V_k s, but also on “location” weight parameters w_{ik} . These weights satisfy $0 < p_{ik} < 1$. Since the CAR distribution has support over the entire real line, we introduce a transformation $\text{logit}(w_{ik}) = z_{ik}$ and allow the z_{ik} ’s to be distributed as CAR. Of course, any other link mapping the unit interval to the real line could be used. For each k , we let $\{z_{ik}\}_{i=1}^n$ be distributed as a CAR distribution yielding a Markov random field (MRF) on the location weights and allowing the desired smoothing across neighbors. Usually larger values of ρ induce greater smoothing and setting $\rho = 1$ yields the popular ICAR prior. This prior is improper as $D - W$ is singular, but for a map without islands this issue can be resolved by imposing the additional constraint $\sum_{i=1}^n z_{ik} = 0$.

The ARSB model, truncated to m terms for the stick-breaking representation, with a Poisson likelihood is

$$\begin{aligned}
 Y_i | \boldsymbol{\beta}, \phi_i &\sim \text{Poisson}(E_i e^{\mu_i}), \mu_i = \mathbf{x}'_i \boldsymbol{\beta} + \phi_i; \phi_i \sim G^{(i)}; \\
 G^{(i)}(\cdot) &= \sum_{k=1}^m p_{ik} \delta_{\theta_k}(\cdot), \theta_k \sim N(0, \sigma_s^2); p_{i1} = w_{i1}V_1, \\
 p_{ik} &= w_{ik}V_k \prod_{l=1}^{k-1} (1 - w_{il}V_l), V_k \stackrel{i.i.d.}{\sim} \text{Beta}(1, \alpha), \{z_{ik}\} \sim \text{CAR}(\rho, \sigma_k^2),
 \end{aligned} \tag{3.1}$$

where $\text{logit}\{w_{ik}\} = z_{ik}$ for $k = 1, \dots, m$. The α parameter stochastically controls the number of distinct values among the n observations, and the covariance between dependent variables Y_i and Y_j is induced by the covariance of the spatial random effects ϕ_i and ϕ_j .

The ARSB model incorporates dependence between the discrete distributions on different regions but does not yield identical marginal distributions on the ϕ_i . Duan, Guindani, and Gelfand (2007) introduced random distributions for the spatial effects associated with point-referenced data allowing different surface selection at different sites while ensuring that the marginal distribution of the effect at each sites still comes from a Dirichlet process. Here we propose an areal alternative, which we call the areally-referenced Dirichlet process (ARDP). The ARDP maintains the marginal distribution of each spatial random effect to be a regular univariate DP while incorporating the spatial dependence between these DPs.

Consider spatial random effects $\phi_i, i = 1, \dots, n$, arising marginally from an identical random measure G , where $G \sim DP(\alpha, G_0)$. We introduce spatial dependence between these DPs by constructing dependent uniform $(0, 1)$ random variables. Suppose $\gamma_1, \dots, \gamma_n$ are jointly distributed as a $CAR(\rho, \sigma_\gamma)$, and $F^{(1)}(\cdot), \dots, F^{(n)}(\cdot)$ denote the cumulative distribution functions of the marginal distributions of each component of the CAR random vector. Marginally, each $F^{(i)}(\gamma_i)$ is uniform $(0, 1)$ but they are *dependent* through $\gamma_1, \dots, \gamma_n$.

We formulate our hierarchical areally-referenced Dirichlet process (ARDP) model as follows. We use a Poisson likelihood for the first stage model, though this could be replaced by any discrete distribution in the exponential family. Here,

$$\begin{aligned}
 Y_i | \boldsymbol{\beta}, \phi_i &\sim \text{Poisson}(E_i e^{\mu_i}), \mu_i = \mathbf{x}'_i \boldsymbol{\beta} + \phi_i; \boldsymbol{\phi} = \{\phi_i\}_{i=1}^n \sim G_n; \\
 G_n &= \sum_{u_1, \dots, u_n} \pi_{u_1, \dots, u_n} \delta_{\theta_{u_1}} \cdots \delta_{\theta_{u_n}}; \\
 \pi_{u_1, \dots, u_n} &= P\left(\sum_{k=1}^{u_1-1} p_k < F^{(1)}(\gamma_1) < \sum_{k=1}^{u_1} p_k, \dots, \sum_{k=1}^{u_n-1} p_k < F^{(n)}(\gamma_n) < \sum_{k=1}^{u_n} p_k\right); \\
 \theta_k &\stackrel{i.i.d.}{\sim} N(0, \sigma_s^2); p_1 = V_1; p_j = V_j \prod_{k < j} (1 - V_k); V_j \stackrel{i.i.d.}{\sim} \text{Beta}(1, \alpha); \\
 \boldsymbol{\gamma} &= \{\gamma_i\}_{i=1}^n \sim N_n(\mathbf{0}, \boldsymbol{\Sigma}), \tag{3.2}
 \end{aligned}$$

where $k = 1, 2, \dots, K$ truncates the stick breaking function to K terms, $\boldsymbol{\Sigma} = \sigma_\gamma^2(D - \rho W)^{-1}$ is the covariance matrix of a *proper* CAR distribution. Using the cumulative distribution function of the γ_i 's to model the weights is an adaptation of the Hybrid Dirichlet Process (Petrone, Guindani, and Gelfand (2009)), where copulas are used to model weights. The distinction is that we model areal dependence using Markov random fields, while the hybrid DP models dependence using continuous spatial processes for inference on uncountable sets. Both methods ensure that the marginal distribution of $G^{(i)}(\phi_i)$, for each i , follows an identical DP

$$G^{(i)}(\phi_i) = \sum_{k=1}^K \sum_{u_1, \dots, u_i=k, \dots, u_n} \pi_{u_1, \dots, u_i=k, \dots, u_n} \delta_{\theta_{u_1}} \cdots \delta_{\theta_{u_i=k}} \cdots \delta_{\theta_{u_n}} = \sum_{k=1}^K p_k \delta_{\theta_k}, \tag{3.3}$$

where $p_k = \sum_{i=1}^K P\left(\sum_{t=1}^{k-1} p_t < F^{(i)}(\gamma_i) < \sum_{t=1}^k p_t\right)$. How the covariance between ϕ_i and ϕ_j depends upon the probabilities p_1, \dots, p_K can be seen from

$$\text{Cov}(\phi_i, \phi_j) = \sigma_s^2 \sum_{l=1}^K P(u_i = u_j = l)$$

$$\begin{aligned}
 &= \sigma_s^2 \sum_{l=1}^K P \left(\sum_{k=1}^{l-1} p_k < F^{(i)}(\gamma_i) < \sum_{k=1}^l p_k, \sum_{k=1}^{l-1} p_k < F^{(j)}(\gamma_j) < \sum_{k=1}^l p_k \right) \\
 &= \sigma_s^2 \sum_{l=1}^K p_l P \left(F^{(i)-1} \left(\sum_{k=1}^{l-1} p_k \right) < \gamma_i < F^{(i)-1} \left(\sum_{k=1}^l p_k \right) \right. \\
 &\quad \left. \left| F^{(j)-1} \left(\sum_{k=1}^{l-1} p_k \right) < \gamma_j < F^{(j)-1} \left(\sum_{k=1}^l p_k \right) \right), \tag{3.4}
 \end{aligned}$$

where (γ_i, γ_j) follows a bivariate normal distribution with covariance specified by the CAR model. Posterior inference for the ARSB and ARDP models are based upon Markov chain Monte Carlo simulations (e.g., Gelman et al. (2004); Carlin and Louis (2008)). The details are outlined in the Supplement.

3.4. A practical FDR-based method to select difference boundaries

To obtain a threshold for detecting difference boundaries, our approach treats the spatial boundary analysis problem as one of multiple hypothesis testing. For each pair of adjacent regions, say i and j , we test $\phi_i = \phi_j$ against $\phi_i \neq \phi_j$. This produces as many hypothesis as there are edges. Recently, several authors have advocated the use of the false discovery rate (FDR) to adjust for multiplicities in hypothesis testing problems (see, e.g., Benjamini and Hochberg (1995); Efron et al. (2001); Storey (2002, 2003)). Li et al. (2012) used the FDR on Smoothed Moving Average (SMA) models involving some awkward constraints on the random effects for model fitting. We adapt this approach to the ARDP and ARSB models free of such constraints.

We identify a boundary (i, j) as a difference boundary if the posterior probability that $P(\phi_i = \phi_j | Y)$ exceeds a certain threshold t , where $Y = \{Y_1, \dots, Y_n\}$ is the entire collection of observed outcomes. For each pair of neighboring regions, we construct $A_{(i,j)}(Y; t) = \{Y : P(\phi_i \neq \phi_j | Y) > t\}$, a *critical region* that indicates evidence in favor of (i, j) being a difference boundary. The choice of t will control the FDR below a level $\delta = 0.05$. If $Z_{(i,j)} = I(\phi_i = \phi_j)$ and $v_{(i,j)} = P(Z_{(i,j)} = 0 | Y)$, then the FDR is

$$FDR = \frac{\sum_{i \sim j} Z_{(i,j)} I(v_{(i,j)} > t)}{\sum_{i \sim j} I(Z_{(i,j)} > t)}, \text{ where } i \sim j \text{ if } w_{ij} \neq 0. \tag{3.5}$$

Estimation of (3.5) is straightforward. It is obtained as the posterior expectation

$$\widehat{FDR} = E[FDR | Y] = \frac{\sum_{i \sim j} (1 - v_{(i,j)}) I(v_{(i,j)} > t)}{\sum_{i \sim j} I(v_{(i,j)} > t)}, \tag{3.6}$$




Figure 1. A map of the simulated data with the grey-scales showing the six different clusters, each having its own mean. There are 47 boundary segments that separate regions with different means (shades). The percentages reflect the quantiles for the distribution of the outcomes.

where $v_{(i,j)}$ is computed as a Monte Carlo mean of the posterior samples for Z_{ij} . Rejection rules can be then constructed to bound the FDR at target level δ : reject if $v_{(i,j)} > t$, where

$$t = \sup \left\{ u : \frac{\sum_{i \sim j} I(v_{(i,j)} > u)(1 - v_{(i,j)})}{\sum_{i \sim j} I(v_{(i,j)} > u)} \leq \delta \right\}.$$

4. A Simulation Study

To evaluate our methods, we conducted a simulation study using the template of a Minnesota county map in Li, Banerjee, and McBean (2011). There are $n = 87$ counties in Minnesota, and 211 pairs of neighboring counties. We simulated 50 datasets on a map of Minnesota, where the state was divided into six regions. Each dataset was generated from (2.1), where μ_i was one of five different means corresponding to the five different shades mapped on Figure 1; the darker shades correspond to higher means. To add some irregularity, we also included one county (Sherburne county shaded white in Figure 1) that has all its boundaries as true difference boundaries. This resulted in “six” different clusters on the map and 47 “true difference boundaries” delineating the clusters with substantially different means.

For every pair of geographical neighbors (i, j) , we computed the posterior probability $P(\phi_i \neq \phi_j | Y)$ and chose the $T = 35, 40, 45, 50$ and 55 edges with the highest posterior probabilities. As there are 47 true difference boundaries, these choices encompass settings where we could, theoretically have obtained 100% accuracy (when $T = 35, 40, 45$) and also where we are assured of a few false positives (when $T = 50, 55$).

The prior specification and computational details of the ARDP model are in the Web Supplement. The parameter α can be fixed based upon the expected number of clusters *a priori*. In this case, six clusters suggest a value of α around 1.25. We experimented with α ranging from 0.25 to 1.75 and obtained very robust inference. The results presented here correspond to $\alpha = 0.5$, which leads to an expected number of clusters of around 3, which is half the number of true clusters. We also fixed $\rho = 0.98$ in the ARDP model (ICAR is inappropriate since the covariance matrix must be proper and nonsingular). Customarily, higher values of ρ (≈ 1) yield sufficient smoothing, while values lower than 0.95 tend not to (e.g., Banerjee, Carlin, and Gelfand (2004)). For ARSB, we used the ICAR model (with $\rho = 1$) along with the sum-to-zero constraint, which yields legitimate posterior samples. We assumed a regression structure with only an intercept (i.e., $\mathbf{x}_i \equiv 1$) and placed a flat prior on the corresponding β . A weakly informative prior $\Gamma(0.01, 0.01)$ was specified for the precision parameters τ_s and τ_γ . In both models, the stick-breaking prior was computed using about 15 terms.

We compared the performance of DPM, ARSB, ARDP with three existing methods: the deterministic Boundary Likelihood Value (BLV) algorithm of Jacquez and Greiling (2003a,b) using the `BoundarySEER` software (see <http://www.biomedware.com>) with default thresholds set from a BLV histogram; the model-based approach of Lu and Carlin (2005), which we call the “LC method”; a class of discrete Spatial Moving Average (SMA) models outlined in Li, Banerjee, and McBean (2011). We ran these models within the R statistical software environment running three parallel chains for each model and dataset. Convergence was diagnosed after 12,000 iterations of burn-in using Gelman-Rubin diagnostics and autocorrelation plots from the `coda` package in R. A subsequent $5,000 \times 3 = 15,000$ samples were used for posterior inference. On a workstation using a Intel dual core 4 GHz processor, each model took less than five hours of CPU time to deliver its entire inferential output for all the 50 simulated datasets. Minnesota Pneumonia and Influenza data consumed less than fifteen minutes of CPU time.

Table 1 presents the average detection rates for these different methods applied to the 50 simulated datasets. The DPM and the BLV methods do not explicitly borrow strength across neighbors, while the other four methods in Table 1 exploit the adjacency structure of the underlying map. There seems to be little to choose between the ARDP and ARSB, but both methods seem to be slightly outperforming the other methods in both sensitivity and specificity under all five scenarios. In addition, while the performance of the SMA model is perhaps comparable, it is computationally onerous and less robust to prior assumptions (Li, Banerjee, and McBean (2011)) than ARDP or ARSB.

The LC method is based upon a parametric CAR model that does not render itself to probabilistic boundary analysis (since $P(\phi_i = \phi_j)$ will always be zero).

Table 1. Sensitivity and specificity in the simulation study (50 datasets generated on a Minnesota map) for the ARDP, ARSB, DPM, LC and BLV methods.

T	Method	Sensitivity	Specificity	T	Method	Sensitivity	Specificity
35	ARDP	0.768	0.998	40	ARDP	0.822	0.990
	ARSB	0.771	0.991		ARSB	0.821	0.989
	DPM	0.737	0.989		DPM	0.791	0.991
	BLV	0.711	0.990		BLV	0.778	0.979
	LC	0.702	0.989		LC	0.767	0.976
	SMA	0.740	0.998		SMA	0.818	0.991
45	ARDP	0.881	0.971	50	ARDP	0.927	0.962
	ARSB	0.878	0.972		ARSB	0.930	0.968
	DPM	0.870	0.968		DPM	0.897	0.952
	BLV	0.831	0.964		BLV	0.869	0.944
	LC	0.813	0.959		LC	0.859	0.941
	SMA	0.872	0.975		SMA	0.901	0.955
55	ARDP	0.940	0.943				
	ARSB	0.941	0.940				
	DPM	0.895	0.915				
	BLV	0.891	0.920				
	LC	0.881	0.917				
	SMA	0.925	0.930				

However, one could fit parametric CAR models and use the posterior expectation of the absolute differences of the rates $E(\|\eta_i - \eta_j\| | Y)$, where $\eta_i = \mu_i/E_i$ acts as a boundary difference score. Higher values indicate spatial barriers between units i and j . The DPM, ARSB and ARDP models not only yield estimates of η_i , as in the “LC” method, but they also deliver nonzero posterior probabilities $P(\phi_i = \phi_j | Y)$. Therefore, we used the posterior expectation metric to compare its performance. The SMA model does not deliver posterior estimates of spatial effects from a single model. Hence, we exclude it from this comparison. we compare our models with the deterministic Boundary Likelihood Value (BLV) algorithm of Jacquez and Greiling (2003a,b).

Table 2 presents the results for four of the methods. The deterministic BLV method detects 89.6% of the boundaries. The promise of our stochastic models is evident from the superior performances of the ARDP and the ARSB models. Since we know the true boundaries in Figure 1, we can assess the performances of these approaches in detecting the true boundaries. Using direct posterior estimates 47 difference boundaries. We find that the DPM, ARDP and the ARSB models are each able to detect about 90% of the true boundaries. The ARDP model performs slightly better than the other two. Using the posterior expectation metric, we again find that the proposed ARSB and ARDP models

Table 2. Assessment of the true wombling boundaries with those produced by LC, ARDP and ARSB based on $P(\phi_i = \phi_j | Y)$ and $E(\|\eta_i - \eta_j\| | Y)$ in the simulation study.

	Assessment using $P(\phi_i = \phi_j Y)$	Assessment using $E(\ \eta_i - \eta_j\ Y)$
LC	-	78.7%
DPM	89.3%	82.2%
ARDP	91.4%	88.3%
ARSB	89.1%	83.3%

outperforming the LC method; they outperform the DPM model as well in terms of the posterior expectation metric.

5. Analysis of Minnesota *P&I* Dataset

We applied our method to the Minnesota Pneumonia and Influenza (*P&I*) diagnosis dataset. *P* & *I* rank as the eighth leading cause of death in the United States and the sixth leading cause in people over 65 years of age, with Pneumonia consistently accounting for the overwhelming majority of deaths of the two. Together, they cost the U.S. economy in 2005 an estimated \$40.2 billion. Identifying difference boundaries that perform well with regard to sensitivity and specificity can help identify so-called “health barriers” more accurately and buttress an active surveillance program for an influenza-like illness.

We analyzed a dataset consisting of Minnesota residents above 65 years of age who were enrolled in the Medicare fee-for-service program as of December 31, 2001. The Medicare Denominator file for 2001 was used to define the cohort. The Medicare Provider Analysis and Review (MedPAR) manages patient records based on date of discharge and supplied information regarding hospitalizations resulting from *P&I*. Rates of *P&I* hospitalization are traditional measures of the impact of influenza virus in the elderly population. We identify the ‘boundaries’ that separate the more affected areas from the less affected areas.

If Y_i and O_i are the observed number of hospitalizations and the population in county i respectively, then $E_i = (\sum_{k=1}^n Y_k / \sum_{k=1}^n O_k) O_i$ is the expected number of cases (under the assumption of no spatial variation in rates), where n is the total number of counties. The choropleth map of the raw data is shown in Figure 2. The high-valued SMR (standard mortality ratio) counties are scattered over the map, with a clump in the southwest and some isolated regions surrounded by sparsely inhabited counties that also have lower counts.

We employed the models in Section 4 to detect boundaries on the *P&I* hospitalization map. The same prior specification and model settings were applied here as the simulation study, except we took $\alpha = 1$, a customary choice when one does not seek a prior distribution on this parameter (Escobar and West




Figure 2. Choropleth map of the SMR in MN (P&I) dataset. The percentages reflect the quantiles for the distribution of the SMR.

(1995)) or has no *a priori* information about the number of clusters. Three parallel MCMC chains were executed on the same computing environment as described in Section 4. Convergence was diagnosed after 10,000 iterations of burn-in using Gelman-Rubin diagnostics and autocorrelation plots and a subsequent $5,000 \times 3 = 15,000$ samples were used for posterior inference. Each model consumed less than ten minutes of CPU time to produce its entire inferential output for the Minnesota Pneumonia and Influenza dataset with very little difference between the ARSB, ARDP and the (non-spatial) DPM model.

Health administrators may prefer to use a “top bracket” of most likely difference boundaries for policy formulation. The top 50 difference boundaries detected by each model are highlighted in Figure 3. Table 3 presents a comprehensive “lookup table” containing the names of adjacent counties that have been ranked in decreasing order according to $1 - P(\phi_i = \phi_j | Data)$ from the ARDP model. Instead of selecting this “bracket” arbitrarily, statisticians may prefer a threshold obtained by controlling the FDR. Setting $\delta = 5\%$ yields Numbers 1-33 as difference boundaries, while setting $\delta = 10\%$ detects Numbers 1-42 as difference boundaries. This table offers an easy reference for health administrators and officials to identify the more substantial spatial health barriers in the state.

About 90% of the boundaries listed in Table 3 are detected by all four models. As a specific example consider Cook and Koochiching county. The outcome variable in the former is substantially higher than its only neighbor, Lake, while Koochiching county is separated from all its neighbors due to its extremely high *P&I* SMR, even after being smoothed by the model. Among the 50 difference boundaries detected by the ARDP model, 47 are also detected by the ARSB model. The three county-pairs that went undetected by ARSB were Goodhue and Olmsted, Freeborn and Steele, and Big Stone and Traverse. The ARSB




Figure 3. Difference boundaries detected by various models in the Minnesota (P&I) dataset.

model detected boundaries between counties Becker and Wadena, Cotton Wood and Jackson, and Cook and Lake.

The map in Figure 2 does not display clustering as pronounced as did the simulation example. It does, however, reflect well on our models that the rankings in Table 3 are very consistent with competing methods. The agreement between the ARDP and the SMA in terms of identifying the difference boundaries using FDR-based thresholds is very strong with over 95% agreement in boundary selection.

6. Conclusion and Future Work

The paper presented a class of nonparametric Bayesian hierarchical models for detecting difference boundaries on maps. An advantage of the new approach is that it permits the probabilistic estimation of an edge as a difference boundary, and improves the percentage of true detection. A disadvantage is that the model cannot be easily fit into any existing commercial software. We fit these models in R (www.r-project.org), and we hope to collect these models in an R package in the near future.

The ARDP and ARSB models in conjunction with the FDR controlled threshold selection provide a major improvement over earlier work by Li et al.

Table 3. Names of adjacent counties that have significant boundary effects from the ARDP model. The numbers in the first column are the ranks according to $P(\phi_i = \phi_j | Y)$

1	Beltrami , Koochiching	26	Koochiching, Lake of the Woods
2	Cass, Wadena	27	Isanti, Mille Lacs
3	Douglas , Pope	28	Chippewa, Renville
4	Freeborn , Steele	29	Murray, Pipestone
5	Goodhue , Olmsted	30	Becker ,Mahnomen
6	Itasca , Koochiching	31	Rice , Waseca
7	Kandiyohi, Pope	32	Blue Earth, Brown
8	Koochiching, St. Louis	33	Dodge, Olmsted
9	Pope, Stearns	34	Chisago , Isanti
10	Anoka , Isanti	35	Redwood , Yellow Medicine
11	Dakota, Goodhue	36	Pennington, Polk
12	Lincoln, Pipestone	37	Goodhue, Wabasha
13	Murray, Redwood	38	Pope, Swift
14	Steele, Waseca	39	Morrison, Todd
15	Renville, Yellow Medicine	40	Fillmore, Olmsted
16	Cottonwood, Murray	41	Cook , Lake
17	Jackson , Martin	42	Douglas, Grant
18	Kandiyohi, Swift	43	Mahnomen,Norman
19	Pope, Stevens	44	Grant, Wilkin
20	Todd, Wadena	45	Mahnomen, Polk
21	Lyon, Redwood	46	Jackson, Nobles
22	Murray, Nobles	47	Morrison, Todd
23	Isanti, Sherburne	48	Dodge , Olmsted
24	Otter Tail , Todd	49	Big Stone , Traverse
25	Clay, Otter Tail	50	Morrison, Stearns

(2012). However, issues related to optimal selection of boundaries warrants further investigation, especially regarding the sensitivity of the inference to FDR-based cutoffs and to prior specifications. Further extensions can be formulated by incorporating classes of loss functions, as discussed by Müller, Parmigiani, and Rice (2006), for a more comprehensive decision-theoretic framework. Such developments may, in turn, lead to more definitive conclusions regarding the performance of these models in maps that display weaker clustering patterns.

The ARDP and ARSB models render themselves to multivariate extensions, where multiple health outcomes need to be modeled jointly. Analogous to the univariate ARSB model, we can construct a multivariate areally referenced stick breaking (MARSB) model such that the sticking breaking weights \mathbf{p} are correlated through another type of “weight” parameter that scales the V_i 's. We place a multivariate CAR (MCAR) prior on these “weight” parameters so as to capture both spatial correlation and inter-variable correlation (Jin, Carlin, and Banerjee

(2005, 2007)). For the ARDP, the distribution of the spatial random effects can be modeled by constructing multivariate areally referenced Dirichlet processes (MARDP). These pursuits constitute natural extensions of our current work.

Acknowledgement

The authors are grateful to two reviewers, an associate editor, and the Co-Editor for their insightful comments and suggestions which have improved the manuscript significantly. This work was supported by federal grants NSF/DMS-1106609 and NIH/NIGMS 1-RC1-GM092400-01.

References

- Anselin, L. (1988). *Spatial Econometrics: Methods and Models*. Kluwer Academic Publishers, Boston, MA.
- Banerjee, S., Carlin, B. P. and Gelfand, A. E. (2004). *Hierarchical Modeling and Analysis for Spatial Data*. Chapman and Hall/CRC Press, Boca Raton, FL.
- Banerjee, S. and Gelfand, A. E. (2006). Bayesian Wombling: Curvilinear gradients assessment under spatial process models. *J. Amer. Statist. Assoc.* **101**, 1487-1501.
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B* **57**, 289-300.
- Blackwell, D. and MacQueen, J. B. (1973). Ferguson distributions via Polya urn schemes. *Ann. Statist.* **1**, 353-355.
- Carlin, B. P. and Louis, T. A. (2008). *Bayesian Methods for Data Analysis*. 3rd edition. Chapman and Hall/CRC, Boca Raton, FL.
- Chung, Y. and Dunson, D. B. (2009). Nonparametric Bayes conditional distribution modeling with variable selection. *J. Amer. Statist. Assoc.* **104**, 1646-1660. .
- Chung, Y. and Dunson, D. B. (2011). The local Dirichlet process. *Ann. Instit. Statist. Math.* **63**, 59-80.
- Cressie, N. (1993). *Statistics for Spatial Data*. Revised edition. Wiley, Hoboken, NJ.
- De Iorio, M., Mueller, P., Rosner, G. L. and MacEachern, S. N. (2004). An ANOVA model for dependent random measures. *J. Amer. Statist. Assoc.* **99**, 205-215.
- Duan, J., Guindani, M. and Gelfand, A. E. (2007). Generalized spatial Dirichlet process models. *Biometrika* **94**, 809-825.
- Dunson, D. B. and Park, J.-H. (2008). Kernel stick-breaking processes. *Biometrika* **95**, 307-323.
- Dunson, D. B., Pillai, N. S. and Park, J.-H. (2007). Bayesian density regression. *J. Roy. Statist. Soc. Ser. B* **69**, 163-183.
- Efron, B., Tibshirani, R., Storey, J. D. and Tusher, V. (2001). Empirical Bayes analysis of a microarray experiment. *J. Amer. Statist. Assoc.* **96**, 1151-1160.
- Escobar, M. D. and West, M. (1995). Bayesian density estimation and inference using mixtures. *J. Amer. Stat. Assoc.* **90**, 577-588.
- Ferguson, T. S. (1973). A Bayesian analysis of some nonparametric problems. *Ann. Statist.* **1**, 209-230.
- Gelfand, A. E., Kottas, A. and Maceachern, S. N. (2005). Bayesian nonparametric spatial modelling with Dirichlet process mixing. *J. Amer. Statist. Assoc.* **100**, 1021-1035.

- Gelman, A., Carlin, J. B., Stern, H. S. and Rubin, D. B. (2004). *Bayesian Data Analysis*. Chapman and Hall/CRC, Boca Raton, FL.
- Griffin, J. E. and Steel, M. F.J. (2006). Order-based dependent Dirichlet processes. *J. Amer. Statist. Assoc.* **101**, 179-194.
- Hoeting, J. A., Madigan, D., Raftery, A. E. and Volinsky, C. T. (1999). Bayesian model averaging: A tutorial. *Statist. Sci.* **14**, 382-401.
- Ishwaran, H. and James, L. F. (2001). Gibbs sampling methods for stick-breaking priors. *J. Amer. Statist. Assoc.* **96**, 161-173.
- Ishwaran, H. and Zarepour, M. (2000). Markov chain Monte Carlo in approximate Dirichlet and beta two-parameter process hierarchical models. *Biometrika* **87** 371-390.
- Jacquez, G. M. and Greiling, D. A. (2003a). Local clustering in breast, lung and colorectal cancer in Long Island, New York. *Int. J. Health. Geogr.* **2**, 3.
- Jacquez, G. M. and Greiling, D. A. (2003b). Geographic boundaries in breast, lung and colorectal cancers in relation to exposure to air toxics in Long Island, New York. *Int. J. Health. Geogr.* **2**, 4.
- Jin, X., Carlin, B. P. and Banerjee, S. (2005). Generalized hierarchical multivariate CAR models for areal data. *Biometrics* **61**, 950-961.
- Jin, X., Carlin, B. P. and Banerjee, S. (2007). Order-free co-regionalized areal data models with application to multiple-disease mapping. *J. Roy. Statist. Soc. Ser. B* **65**, 817-838.
- Kalli, M., Griffin, J. E. and Walker, S.G. (2011). Slice sampling mixture models. *Statist. Comp.* **21**, 93-105.
- Le Sage, J. and Pace, K. (2009). *Introduction to Spatial Econometrics*. Chapman and Hall/CRC, Boca Raton, FL.
- Li, P., Banerjee, S. and McBean, A. M. (2011). Mining edge effects in areally referenced spatial data: A Bayesian model choice approach. *Geoinformatica* **15**, 435-454.
- Li, P., Banerjee S., McBean A.M. and Carlin, B.P. (2012). Bayesian areal wombling using false discovery rates. *Statist. and its Interface* **5**, 149-158.
- Lu, H. and Carlin, B. P. (2005). Bayesian areal wombling for geographical boundary analysis. *Geogr. Anal.* **37**, 265-285.
- Lu, H., Reilly, C., Banerjee, S. and Carlin, B.P. (2007). Bayesian areal wombling via adjacency modeling. *Environ. Ecol. Statist.* **14**, 433-452.
- Ma, H., Carlin, B. P. and Banerjee, S. (2010). Hierarchical and joint site-edge methods for Medicare hospice service region boundary analysis. *Biometrics* **66**, 355-364..
- MacEachern, S. N. (2001). Decision theoretic aspects of dependent nonparametric processes. In *Bayesian Methods with Applications to Science, Policy and Official Statistics* (Edited by E. George), 551-560. Eurostat.
- Müller, P., Erkanli, A. and West, M. (1996). Bayesian curve fitting using Multivariate normal mixtures. *Biometrika* **83**, 67-79.
- Müller, P., Parmigiani, G. and Rice, K. (2006). FDR and Bayesian multiple comparisons rules. In *Bayesian Statistics 8*. (Edied by J. M. Bernardo, S. Bayarri, J. O. Berger, A. P. Dawid, D. Heckerman, A. F. M. Smith and M. West). Oxford University Press.
- Müller, P., Quintana, F. A. and Rosner, G. L. (2011). A product partition model with regression on covariates. *J. Comput. Graph. Statist.* **20**, 260-278.
- Park, J.-H. and Dunson, D. B. (2010). Bayesian generalized product partition model. *Statist. Sinica*, **20**, 1203-1226.

- Petrone, S., Guindani, M. and Gelfand, A. E. (2009). Hybrid Dirichlet mixture models for functional data. *J. Roy. Statist. Soc. Ser. B.* **71**, 755-782.
- Reich, B. and Fuentes, M. (2007). A multivariate semiparametric Bayesian spatial modeling framework for hurricane surface wind fields. *Ann. Appl. Statist.* **1**, 249-264.
- Sethuraman, J. (1994). A constructive definition of Dirichlet priors. *Statist. Sinica* **4**, 639-650.
- Storey, J. D. (2002). A direct approach to false discovery rates. *J. Roy. Statist. Soc. Ser. B* **64**, 479-498.
- Storey, J. D. (2003). The positive false discovery rate: A Bayesian interpretation and the q -value. *Ann. Statist.* **31**, 2013-2035.
- Womble, W. H. (1951). Differential systematics. *Science* **114**, 315-322.

Medtronic Incorporated, 710 Medtronic Pkwy NE, Minneapolis, MN 55432, USA.

E-mail: pei.li@medtronic.com

Department of Biostatistics, Suite 51-254 CHS UCLA Fielding School of Public Health, 650 Charles E. Young Drive South Los Angeles, CA 90095-1772, USA.

E-mail: sudipto@ucla.edu

Division of Biostatistics, University of Minnesota, 420 Delaware Street SE, Minneapolis, MN 55455, USA.

E-mail: baner009@umn.edu

Division of Health Policy & Management, School of Public Health, University of Minnesota, Twin Cities 420 Delaware Street SE, A369-1 Mayo MMC 729 Minneapolis MN 55455, USA.

E-mail: mcbea002@umn.edu

(Received August 2013; accepted May 2014)